

Coupled McKean-Vlasov diffusions

Duong, Manh Hong; Tugaut, J.

DOI:

[10.1080/17442508.2019.1677663](https://doi.org/10.1080/17442508.2019.1677663)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Duong, MH & Tugaut, J 2019, 'Coupled McKean-Vlasov diffusions: wellposedness, propagation of chaos and invariant measures', *Stochastics: an international journal of probability and stochastic processes*.
<https://doi.org/10.1080/17442508.2019.1677663>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is an Accepted Manuscript of an article published by Taylor & Francis in *Stochastics: an international journal of probability and stochastic processes* on 22/10/2019, available online: <https://www.tandfonline.com/doi/full/10.1080/17442508.2019.1677663>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Coupled McKean-Vlasov diffusions: wellposedness, propagation of chaos and invariant measures

Manh Hong Duong* Julian Tugaut†

July 15, 2019

Abstract

In this paper, we study a two-species model in the form of a coupled system of nonlinear stochastic differential equations (SDEs) that arises from a variety of applications such as aggregation of biological cells and pedestrian movements. The evolution of each process is influenced by four different forces, namely an external force, a self-interacting force, a cross-interacting force and a stochastic noise where the two interactions depend on the laws of the two processes. We also consider a many-particle system and a (nonlinear) partial differential equation (PDE) system that associate to the model. We prove the wellposedness of the SDEs, the propagation of chaos of the particle system, and the existence and (non)-uniqueness of invariant measures of the PDE system.

Key words and phrases. interacting particle systems, McKean-Vlasov dynamics, propagation of chaos, invariant measures.

AMS subject classification. 60H10; 35K55; 60J60; 60G10.

1 Introduction

In this paper, we study a two-species model in the form of a coupled system of nonlinear stochastic differential equations

$$dX_t = -\nabla V_1(X_t) dt - a\nabla F_{11} * \mu_t(X_t) dt - (1-a)\nabla F_{12} * \nu_t(X_t) dt + \sigma dW_t, \quad (1a)$$

$$dY_t = -\nabla V_2(Y_t) dt - a\nabla F_{21} * \mu_t(Y_t) dt - (1-a)\nabla F_{22} * \nu_t(Y_t) dt + \sigma d\widehat{W}_t, \quad (1b)$$

$$\mathbb{P}(X_t \in dx) = \mu_t(x) dx, \quad \mathbb{P}(Y_t \in dx) = \nu_t(x) dx. \quad (1c)$$

Here $0 \leq a \leq 1$ and $\sigma > 0$ are given constants; V_1, V_2 are two external potentials; F_{11}, F_{22} are self-interacting potentials describing the interactions among individuals of the same species; F_{12}, F_{21} are cross-interacting potentials representing the interactions between individuals belonging to different species; σ is the diffusion intensity; $(W_t, t \geq 0)$ and $(\widehat{W}_t, t \geq 0)$ are independent Wiener processes and finally $*$ denotes the standard convolution operator: for a function G and a measure γ , the convolution between G and γ , $G * \gamma$, is given by

$$(G * \gamma)(x) = \int G(x-y)\gamma(y) dy.$$

In (1) the evolution of X_t and Y_t depend on their own laws, $\{\mu_t, t \geq 0\}$ and $\{\nu_t, t \geq 0\}$ respectively, that are unknown. Using Itô formula one can show that $\{\mu_t, t \geq 0\}$ and $\{\nu_t, t \geq 0\}$ satisfy the following system of nonlinear nonlocal partial differential equations

$$\partial_t \mu_t = \operatorname{div} \left(\left(\nabla V_1 + a(\nabla F_{11} * \mu_t) + (1-a)(\nabla F_{12} * \nu_t) \right) \mu_t \right) + \frac{\sigma^2}{2} \Delta \mu_t, \quad (2a)$$

*School of Mathematics, University of Birmingham, Birmingham B15 2TT, UK (hduong@bham.ac.uk)

†Université Jean Monnet, Institut Camille Jordan, 23, rue du docteur Paul Michelon, CS 82301, 42023 Saint-Étienne Cedex 2, France (tugaut@math.cnrs.fr)

$$\partial_t \nu_t = \operatorname{div} \left((\nabla V_2 + a(\nabla F_{21} * \mu_t) + (1-a)(\nabla F_{22} * \nu_t)) \nu_t \right) + \frac{\sigma^2}{2} \Delta \nu_t, \quad (2b)$$

$$\mu_0(dx) = \mathbb{P}(X_0 \in dx), \quad \nu_0(dx) = \mathbb{P}(Y_0 \in dx). \quad (2c)$$

System (1) naturally generalizes the one-specie McKean-Vlasov dynamics

$$dZ_t = -\nabla V(Z_t) dt - \nabla F * \zeta_t(Z_t) dt + \sigma dW_t, \quad (3)$$

where ζ_t is the law of Z_t that solves the following (nonlocal nonlinear) PDE

$$\partial \zeta_t = \operatorname{div} \left[(\nabla V + \nabla F * \zeta_t) \zeta_t \right] + \frac{\sigma^2}{2} \Delta \zeta_t. \quad (4)$$

Systems of (multi-species, interacting) nonlinear stochastic differential equations and nonlocal nonlinear PDEs of the type (1)-(4) arise in a plethora of applications such as mathematical biology (bacteria chemotaxis [KS71, KO03, ESV10, CEV11, KRZ18], aggregation of biological cells [EK16, EFK17]), plasma physics and galactic dynamics [BT08], statistical mechanics and granular materials [CMV03, CMV06], pedestrian movements [CLM12, CLM13], risk management [GPY13] and opinion formation [GPY17]. The mathematical analysis of such systems has been getting a lot of attention over the last two decades both in the probability and in the PDE community. In particular, the McKean-Vlasov dynamics has been investigated from various aspects. Existence and uniqueness of solutions of (3) under fairly general assumptions on the external potential V and interacting potential F has been proved [McK66, Fun84, Szn91, M  l96, HIP08, BRTV98, CGM08]. The propagation of chaos, which was introduced by Kac [Kac56] and further developed by Sznitman [Szn91], for the McKean-Vlasov was also proved [BRTV98, Mal03, CGM08]. That is, as n gets large, the n interacting processes

$$dZ_t^i = -\nabla V(Z_t^i) dt - \frac{1}{n} \sum_{j=1}^n \nabla F(Z_t^i - Z_t^j) dt + \sigma dW_t^i, \quad i = 1, \dots, n, \quad (5)$$

behave more and more like the n independent processes

$$dZ_t^i = -\nabla V(Z_t^i) dt - \nabla F * \zeta_t(Z_t^i) dt + \sigma dW_t^i, \quad i = 1, \dots, n,$$

where $(W_t^i)_{t \geq 0}$ are independent Wiener processes and each particle's distribution tends to $\zeta_t(dx) = \zeta_t(x)dx$ where ζ_t solves (4). In addition, the empirical measure $\rho_t^n := \frac{1}{n} \sum_{j=1}^n \delta_{Z_t^j}$ converges in law, on the space $C([0, T], \mathbb{R})$, to $\zeta_t(dx)$. Thus both (3) and (4) can be numerically approximated by simulating the particle system (5) for large n . We also refer the reader to [BGM10, Duo15, JW16, Mon17] for similar results for the Vlasov-Fokker-Planck equation, to [MM13, HM14, MMW15] for analytical approach to propagation of chaos and to recent papers and surveys [JW17a, JW17b, DEGZ18] for further discussions on this interesting topic. Another important aspect of the McKean-Vlasov dynamics, namely the existence and (non)uniqueness of invariant measures and convergence to an invariant measure, also was studied by many authors using different techniques [BRV98, CMV03, Mal03, CMV06, CGM08, BGG13]. One interesting question that still largely remains open in this direction is to characterise the relative basins of attraction of the equilibria of the McKean-Vlasov equation when there are multiple invariant measures [Daw83, Shi87, Tug14a].

In contrast to the McKean-Vlasov equation, the coupled McKean-Vlasov dynamics is less understood although some initial attempts have been made. Herrmann [Her03] obtained results for three aforementioned issues for a special case of (1) where $V_1 = V_2 = 0, F_{11} = F_{22}$ and $F_{21} = F_{12}$, see also [DMR17] for some formal computations regarding the hydrodynamics limit for this case. Another special case, where $V_1 = V_2 = 0$ and $\sigma = 0$, has been studied by several authors: [FF13] established a systematic existence and uniqueness theory of weak measure solutions for system (2) while its equilibrium properties were investigated in [EFK17, DFF16]. More recently, [CL16, Lab17] proved, using a discrete variational approximation scheme à la Jordan-Kinderlehrer-Otto, existence and uniqueness results for a class of parabolic systems with nonlinear diffusion

and nonlocal interaction that includes the PDE system (2). We also refer the reader to recent works [LM17, CJ17, CDJ18, FEF18, CHS18] on similar multi-species systems where a (nonlinear) cross-diffusion is also included.

The aim of the present paper is to study the well-posedness, propagation of chaos phenomenon and the existence of (multiple) invariant measures of the coupled McKean-Vlasov system (1)-(2). We generalize some of the aforementioned results for special cases to the full system and obtain new results. Our method works for systems of finitely many species, but we proceed in this paper for two-species models for simplifying of the presentation.

Well-posedness of (1). Proving the existence and uniqueness of solutions of interacting (multi-species) systems such as (1) is highly nontrivial because of its nonlocality and nonlinearity. When both the confining and interaction potentials are globally Lipschitz, the well-posedness of (1) can be established using the by now standard techniques [McK66, Fun84, Szn91, M  l96]. When either of the potentials is non-Lipschitz, it is a more intricate problem. The following theorem, which is our first result, generalizes similar results of [BRTV98, HIP08, Tug10] for the McKean-Vlasov equation and of [Her03] for the special case of (1) (where $V_1 = V_2 = 0, F_{11} = F_{22}$ and $F_{21} = F_{12}$ as mentioned in a previous paragraph) to the general coupled system (1).

Theorem 1.1. Suppose that Assumption 2.1 holds and that X_0 and Y_0 are such that $\mathbb{E}(|X_0|^{8q^2}) < \infty$ and $\mathbb{E}(|Y_0|^{8q^2}) < \infty$ where $q > 0$ is defined in (H7) of Assumption 2.1. The system (1) admits a unique strong solution on \mathbb{R}_+ . In other words, given a probability space with two Brownian motions, there exists a solution to the system with these Brownian motions.

Propagation of chaos. To describe our result on propagation of chaos for the coupled McKean-Vlasov system, we take two sequences of integers, $(M_n)_{n \in \mathbb{N}}$ and $(N_n)_{n \in \mathbb{N}}$, that go to infinity as n tends to infinity and consider the following system of interacting particles

$$\begin{aligned} dX_t^i &= -\nabla V_1(X_t^i) dt - \frac{1}{N_n + M_n} \sum_{j=1}^{N_n} \nabla F_{11}(X_t^i - X_t^j) dt \\ &\quad - \frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \nabla F_{12}(X_t^i - Y_t^k) dt + \sigma dW_t^i; \quad i = 1, \dots, N_n; \end{aligned} \quad (6a)$$

$$\begin{aligned} dY_t^i &= -\nabla V_2(Y_t^i) dt - \frac{1}{N_n + M_n} \sum_{j=1}^{N_n} \nabla F_{21}(Y_t^i - X_t^j) dt \\ &\quad - \frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \nabla F_{22}(Y_t^i - Y_t^k) dt + \sigma d\widetilde{W}_t^i, \quad i = 1, \dots, M_n. \end{aligned} \quad (6b)$$

Note that the convolution operators $\nabla F_{ij} * \gamma$ ($i, j \in \{1, 2\}, \gamma \in \{\mu, \nu\}$) in (1) are replaced by the average sums in (6). These sums can also be viewed as convolutions between ∇F_{ij} with the empirical measures, μ_t^n and ν_t^n , instead of the laws μ_t and ν_t where

$$\mu_t^n := \frac{1}{M_n + N_n} \sum_{j=1}^{N_n} \delta_{X_t^j} \quad \text{and} \quad \nu_t^n := \frac{1}{M_n + N_n} \sum_{k=1}^{M_n} \delta_{Y_t^k}.$$

We will show that the propagation of chaos phenomenon holds for the system (6), that is, for all $(p, q) \in \mathbb{N}^2$, $(X_t^1, \dots, X_t^p, Y_t^1, \dots, Y_t^q)$ converges as n tends to infinity to $\otimes_{i=1}^p \mu_t \otimes_{j=1}^q \nu_t$, where μ_t, ν_t are respectively the laws of X_t and Y_t that are solutions of (1). This result is the consequence of the following theorem, that is our second result and extends [Her03] to the general case,

Theorem 1.2. Under the same assumption as in Theorem 1.1, for $T < \infty$, we have

$$\lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{t \in [0, T]} (X_t^i - \widehat{X}_t^i)^2 \right] = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \mathbb{E} \left[\sup_{t \in [0, T]} (Y_t^i - \widehat{Y}_t^i)^2 \right] = 0, \quad (7)$$

where $(\widehat{X}_t^i, \widehat{Y}_t^i)$ is a solution to the following system

$$\begin{aligned} d\widehat{X}_t^i &= -\nabla V_1(\widehat{X}_t^i) dt - a(\nabla F_{11} * \mu_t)(\widehat{X}_t^i) dt \\ &\quad - (1-a)(\nabla F_{12} * \nu_t)(\widehat{X}_t^i) dt + \sigma dW_t^i, \quad i = 1, \dots, N_n; \end{aligned} \quad (8a)$$

$$\begin{aligned} d\widehat{Y}_t^i &= -\nabla V_2(\widehat{Y}_t^i) dt - a(\nabla F_{21} * \mu_t)(\widehat{Y}_t^i) dt \\ &\quad - (1-a)(\nabla F_{22} * \nu_t)(\widehat{Y}_t^i) dt + \sigma d\widetilde{W}_t^i, \quad i = 1, \dots, M_n, \end{aligned} \quad (8b)$$

with $(W_t^i, \widetilde{W}_t^i)$ being independent Wiener processes. Note that $\{\widehat{X}_t^i\}_{i=1}^{N_n}$ ($\{\widehat{Y}_t^k\}_{k=1}^{M_n}$ resp.) are identically independent copies of X_t (Y_k resp.).

Existence and non-uniqueness of invariant measures in non-convex landscapes. It is by now well-known that when the confining potential V is not convex the McKean-Vlasov equation exhibits a phase transition phenomenon, that is it may have a unique stationary solution or several ones when the diffusion coefficient (i.e., the temperature) is above or below a critical value [Daw83, Tam84, Shi87, Tug14b, BnCD16]. Similar results of nonuniqueness of the stationary state at low temperatures have been also obtained for McKean-Vlasov equations modeling opinion formation [WLEC17, CP10], for the Desai-Zwanzig model in a two-scale potential [GP18] as well as for the McKean-Vlasov equations on the torus [CP10, CGPS18]. Our third result is the following existence and non-uniqueness of invariant measures. This is significantly different from [Her03] where there is a unique invariant measure.

Theorem 1.3. Suppose that $F_{ij}(x) = \frac{\alpha_{ij}x^2}{2}$ for $i, j \in \{1, 2\}$ and that V_1 and V_2 have a common unique minimizer m^* . Then for any ρ such that

$$\rho \geq \max \left\{ \frac{|V_1^{(3)}(m^*)|}{4V_1''(m^*)(V_1''(m^*) + a\alpha_{11} + (1-a)\alpha_{12})}, \frac{|V_2^{(3)}(m^*)|}{4V_2''(m^*)(V_2''(m^*) + a\alpha_{21} + (1-a)\alpha_{22})} \right\}$$

system (2) has an invariant measure (μ, ν) whose mean values belong to $[m^* - \rho\sigma^2, m^* + \rho\sigma^2] \times [m^* - \rho\sigma^2, m^* + \rho\sigma^2]$. In addition, if V_1 and V_2 are symmetrical, then there is a unique symmetrical invariant measure (μ^0, ν^0) whose mean values are zeros.

This implies that if $V_1 = V_2 = V$ where V is a double-wells landscape, then there are at least three invariant probabilities.

Organisation of the paper. The rest of the paper is organised as follows. In Section 2, we prove Theorem 1.1 on the wellposedness of (1). In Section 3 we study the propagation of chaos phenomenon and establish Theorem 1.2. Finally, in Section 4 we prove Theorem 1.3 on the existence and nonuniqueness of invariant measures.

2 Existence and uniqueness of strong solutions

In this section, we prove Theorem 1.1 establishing the existence and uniqueness of strong solutions of the system (1). We adapt the proof of [BRTV98] for the existence and uniqueness of strong solutions of the McKean-Vlasov dynamics (3), see also [Her03, HIP08, Tug10]. To this end, we transform (1) into a fixed point problem of a map Γ on a functional space Λ , we then show that Γ is a contraction map on a subspace $\Lambda_T \subset \Lambda$ proving the existence and uniqueness of strong solutions over a finite time interval $[0, T]$. The local solution is then extended to become a global one by controlling its moments.

Assumption 2.1. We make the following assumptions.

(H1) The coefficients $\nabla V_1, \nabla V_2, \nabla F_{ij}$ are locally Lipschitz for any $i, j \in \{1, 2\}$.

(H2) The functions V_1, V_2 and F_{ij} are continuously differentiable for any $i, j \in \{1, 2\}$.

(H3) There exist $\theta_1 > 0$ and $\theta_2 > 0$ such that

$$(\nabla V_1(x) - \nabla V_1(y))(x - y) \geq -\theta_1 |x - y|^2 \quad \text{and} \quad (\nabla V_2(x) - \nabla V_2(y))(x - y) \geq -\theta_2 |x - y|^2 \quad \forall x, y. \quad (9)$$

(H4) $xV_1'(x) \geq C_4 x^4 - C_2 x^2$ with $C_2, C_4 > 0$. The same holds with V_2 .

(H5) The potential V_1 is convex at infinity: $\lim_{|x| \rightarrow +\infty} \nabla^2 V_1(x) = +\infty$. The same holds with V_2 .

(H6) There exist $m \in \mathbb{N}$ and $C > 0$ such that $|\nabla V_1(x)| + |\nabla V_2(x)| \leq C|x|^{2m-1}$ and $m \geq 2$.

(H7) ∇F_{11} and ∇F_{22} are odd and increasing with polynomial growth functions, the degree being $2q - 1$.

(H8) ∇F_{12} and ∇F_{21} are Lipschitz.

(H9) $\nabla F_{11}(0) = \nabla F_{12}(0) = \nabla F_{21}(0) = \nabla F_{22}(0) = 0$.

We now need to introduce some functional spaces.

Definition 2.1. On the space of functions from $\mathbb{R}_+ \times \mathbb{R}$ to \mathbb{R} , we introduce the norm

$$\|b\|_T := \sup_{0 \leq s \leq T} \sup_{x \in \mathbb{R}} \left(\frac{|b(s, x)|}{1 + |x|^{2q}} \right).$$

We now introduce the functional space that will be used in the following.

Definition 2.2. We consider the space

$$\Lambda_T := \Lambda_T^1 \cap \Lambda_T^2 \cap \Lambda_T^3,$$

where the three spaces of functions Λ_T^1 , Λ_T^2 and Λ_T^3 are defined by

$$\Lambda_T^1 := \{b : [0; T] \times \mathbb{R} \longrightarrow \mathbb{R} \mid x \mapsto b(s, x) \text{ is locally Lipschitz uniformly in } s\},$$

where the parameter of Lipschitz may depend on b ;

$$\Lambda_T^2 := \{b : [0; T] \times \mathbb{R} \longrightarrow \mathbb{R} \mid x \mapsto b(s, x) \text{ is increasing and } b(s, x) - b(s, y) \geq \xi_1(x - y) + \xi_0\},$$

where $\xi_1 > 0$, $\xi_0 \in \mathbb{R}$ and $x \geq y$; and

$$\Lambda_T^3 := \{b : [0; T] \times \mathbb{R} \longrightarrow \mathbb{R} \mid \|b\|_T < \infty\}.$$

The space Λ_T is equipped with the norm $\|\cdot\|_T$.

Definition 2.3. We finally put $F_T := \Lambda_T \times \Lambda_T \times \Lambda_T \times \Lambda_T$ equipped with the norm

$$\|b\|_T^F := \sum_{i=1}^4 \|b_i\|_T,$$

where $b := (b_1, b_2, b_3, b_4)$.

We will also use a transformation in order to apply a fixed point theorem.

Definition 2.4. We consider Γ from F_T to F_T defined by its coordinates:

$$\begin{aligned} p_1 \circ \Gamma(b)(x) &:= a\mathbb{E} [\nabla F_{11}(x - X_t^b)], \quad p_2 \circ \Gamma(b)(x) := (1 - a)\mathbb{E} [\nabla F_{12}(x - Y_t^b)], \\ p_3 \circ \Gamma(b)(x) &:= a\mathbb{E} [\nabla F_{21}(x - X_t^b)] \quad \text{and} \quad p_4 \circ \Gamma(b)(x) := (1 - a)\mathbb{E} [\nabla F_{22}(x - Y_t^b)], \end{aligned}$$

where p_i is the i th projection on the space F_T and X_t^b (resp. Y_t^b) is solution of the SDE

$$dX_t^b = \sigma dB_t - \nabla V_1(X_t^b) dt - b_1(t, X_t^b) dt - b_2(t, X_t^b) dt, \quad (10)$$

respectively

$$dY_t^b = \sigma d\widetilde{B}_t - \nabla V_2(Y_t^b) dt - b_3(t, Y_t^b) dt - b_4(t, Y_t^b) dt. \quad (11)$$

The fact that Γ is a transformation from F_T to itself will be given in Lemma 2.8. To show that there exist solutions to the equations on X^b and on Y^b , we use the following result (see [SV79, Theorem 10.2.2] at page 255):

Proposition 2.5. Let $b : \mathbb{R}_+ \times \mathbb{R} \longrightarrow \mathbb{R}$ be a function satisfying the three following properties:

1. $\max_{s \geq 0} |b(s, 0)| < \infty$.
2. For any $n \in \mathbb{N}$, there exists a constant $c_n > 0$ such that $|b(s, x) - b(s, y)| \leq c_n |x - y|$ for any reals x and y satisfying $|x| < n$ and $|y| < n$.
3. There exists a constant $r > 0$ such that for any $|x| > r$, $\text{sign}(x)b(s, x) \geq 0$.

Then, for any random variable X_0 , the equation $E^{(b, X_0)}$ admits a unique strong solution where $E^{(b, X_0)}$ is defined by

$$X_t = X_0 - \int_0^t b(s, X_s) ds + \sigma B_t.$$

To show that there is a unique strong solution to the initial system, we search a fixed point to the transformation Γ . To do so, it is easy to check that for any $b \in F_T$, the equations (10) and (11) admit a unique strong solution. Indeed, the convexity at infinity of the potentials V_1 and V_2 guarantees that the third point of Proposition 2.5 is satisfied.

The following definition of moments will play a crucial role in the analysis of this paper.

Definition 2.6. For any $b \in F_T$ and $p > 0$, we define

$$\begin{aligned} \eta_p^b(t) &:= \mathbb{E} \left[|X_t^b|^p \right], \quad \widehat{\eta}_p^b(t) := \sup_{0 \leq s \leq t} \eta_p^b(s), \\ \xi_p^b(t) &:= \mathbb{E} \left[|Y_t^b|^p \right] \quad \text{and} \quad \widehat{\xi}_p^b(t) := \sup_{0 \leq s \leq t} \xi_p^b(s). \end{aligned}$$

To prove Theorem 1.1, we need several lemmas.

Lemma 2.7. Set $b \in F_T$, $n \geq 1$, $\rho := (\rho_0, \rho_0, \rho_0, \rho_0)$ with $\rho_0(x) := \beta_0 x$, then $\widehat{\eta}_{2n}^\rho(T) + \widehat{\xi}_{2n}^\rho < \infty$, for $n \geq 0$ such that $\mathbb{E} [|X_0^{2n}|] < \infty$ and $\mathbb{E} [|Y_0^{2n}|] < \infty$. Moreover:

$$\widehat{\eta}_{2n}^b(T) \leq k_1(n) \left[T^{2n} + (\|b_1 - \rho_0\|_T^{2n} + \|b_2 - \rho_0\|_T^{2n}) \left(T^{2n} + \widehat{\eta}_{4qn}^\rho(T) \right) \right],$$

and

$$\widehat{\xi}_{2n}^b(T) \leq k_2(n) \left[T^{2n} + (\|b_3 - \rho_0\|_T^{2n} + \|b_4 - \rho_0\|_T^{2n}) \left(T^{2n} + \widehat{\xi}_{4qn}^\rho(T) \right) \right],$$

where $k_1(n)$ and $k_2(n)$ are constants which do not depend on b , ρ or T .

Proof. Step 1. By considering $\rho := (\rho_0, \rho_0, \rho_0, \rho_0)$, we thus have the following equation:

$$X_t^\rho = X_0 + \sigma B_t - \int_0^t \nabla V_1(X_s^\rho) ds - 2\beta_0 \int_0^t X_s^\rho ds.$$

For any $n \geq 1$, Itô formula yields

$$\begin{aligned} \frac{d}{dt} \mathbb{E} \left[(X_t^\rho)^{2n} \right] &= n(2n-1) \sigma^2 \mathbb{E} \left[(X_t^\rho)^{2n-2} \right] - 4n\beta_0 \mathbb{E} \left[(X_t^\rho)^{2n} \right] - 2n \mathbb{E} \left[(X_t^\rho)^{2n-1} V_1'(X_t^\rho) \right] \\ &\leq n(2n-1) \sigma^2 \left(\mathbb{E} \left[(X_t^\rho)^{2n} \right] \right)^{1-\frac{1}{n}} - 4n\beta_0 \mathbb{E} \left[(X_t^\rho)^{2n} \right] \\ &\quad - 2nC_4 \mathbb{E} \left[(X_t^\rho)^{2n+2} \right] + 2nC_2 \mathbb{E} \left[(X_t^\rho)^{2n} \right] \\ &\leq n(2n-1) \sigma^2 \left(\mathbb{E} \left[(X_t^\rho)^{2n} \right] \right)^{1-\frac{1}{n}} + 2n(C_2 - 2\beta_0) \mathbb{E} \left[(X_t^\rho)^{2n} \right] \end{aligned}$$

$$-2nC_4 \left(\mathbb{E} \left[(X_t^\rho)^{2n} \right] \right)^{1+\frac{1}{n}}.$$

Note that we have used Assumption (H4) to obtain the first inequality in the above estimates. We deduce that, if $\left(\mathbb{E} \left[(X_t^\rho)^{2n} \right] \right)^{\frac{1}{n}} \geq \frac{C_2 - 2\beta_0 + \sqrt{(C_2 - 2\beta_0)^2 + 2C_4(2n-1)\sigma^2}}{2C_4}$, the right hand-side is nonpositive so that $\frac{d}{dt} \mathbb{E} \left[(X_t^\rho)^{2n} \right]$ is nonpositive. As a consequence,

$$\mathbb{E} \left[(X_t^\rho)^{2n} \right] \leq \max \left\{ \mathbb{E} \left[(X_0)^{2n} \right] ; \left(\frac{C_2 - 2\beta_0 + \sqrt{(C_2 - 2\beta_0)^2 + 2C_4(2n-1)\sigma^2}}{2C_4} \right)^n \right\} < \infty.$$

We deduce that

$$\sup_{t \geq 0} \mathbb{E} \left[(X_t^\rho)^{2n} \right] \leq c_n \left(1 + \mathbb{E} \left[(X_0)^{2n} \right] \right),$$

where c_n is constant. Similarly we obtain

$$\sup_{t \geq 0} \mathbb{E} \left[(Y_t^\rho)^{2n} \right] \leq c_n \left(1 + \mathbb{E} \left[(Y_0)^{2n} \right] \right).$$

As a consequence, if $\mathbb{E} \left[(X_0)^{2n} \right]$ and $\mathbb{E} \left[(Y_0)^{2n} \right]$ are finite, we have that $\widehat{\eta_{2n}^\rho}(T) < \infty$ and $\widehat{\xi_{2n}^\rho}(T) < \infty$.

Step 2. We have

$$\begin{aligned} X_t^b - X_t^\rho &= - \int_0^t [V_1'(X_s^b) - V_1'(X_s^\rho)] ds \\ &\quad - \int_0^t [b_1(s, X_s^b) - p_1 \circ \rho(s, X_s^\rho)] ds - \int_0^t [b_2(s, X_s^b) - p_2 \circ \rho(s, X_s^\rho)] ds. \end{aligned}$$

Consequently, for any $\alpha > 1$, we obtain that $|X_t^b - X_t^\rho|^\alpha$ is equal to

$$\begin{aligned} &- \alpha \int_0^t \text{sign}(X_s^b - X_s^\rho) |X_s^b - X_s^\rho|^{\alpha-1} \mathbb{1}_{X_s^b \neq X_s^\rho} [V_1'(X_s^b) - V_1'(X_s^\rho)] ds \\ &- \alpha \int_0^t \text{sign}(X_s^b - X_s^\rho) |X_s^b - X_s^\rho|^{\alpha-1} \mathbb{1}_{X_s^b \neq X_s^\rho} [b_1(s, X_s^b) - p_1 \circ \rho(s, X_s^\rho)] ds \\ &- \alpha \int_0^t \text{sign}(X_s^b - X_s^\rho) |X_s^b - X_s^\rho|^{\alpha-1} \mathbb{1}_{X_s^b \neq X_s^\rho} [b_2(s, X_s^b) - p_2 \circ \rho(s, X_s^\rho)] ds. \end{aligned}$$

Taking the limit as α goes to 1^+ we get

$$\begin{aligned} |X_t^b - X_t^\rho| &= - \int_0^t \text{sign}(X_s^b - X_s^\rho) [V_1'(X_s^b) - V_1'(X_s^\rho)] ds \\ &\quad - \int_0^t \text{sign}(X_s^b - X_s^\rho) [b_1(s, X_s^b) - p_1 \circ \rho(s, X_s^\rho)] ds \\ &\quad - \int_0^t \text{sign}(X_s^b - X_s^\rho) [b_2(s, X_s^b) - p_2 \circ \rho(s, X_s^\rho)] ds. \end{aligned} \tag{12}$$

We will control each term on the right-hand side of (12). The first one can be controlled by, see [Tug10, Proof of Lemma 2.7]

$$- \int_0^t \text{sign}(X_s^b - X_s^\rho) [V_1'(X_s^b) - V_1'(X_s^\rho)] ds \leq -\gamma \int_0^t |X_s^b - X_s^\rho| ds + T\tilde{\gamma},$$

for any $t \leq T$. In the last formula, γ and $\tilde{\gamma}$ are constants which depend on V_1 and are defined such that $\text{sign}(x - y) (V_1'(x) - V_1'(y)) \geq \gamma(x - y) - \tilde{\gamma}$. The existence of these constants is ensured

by the fact that the lack of convexity of V_1 is only on a compact set. For the second term: since b_1 is increasing

$$\text{sign}(X_s^b - X_s^\rho) [b_1(s, X_s^b) - b_1(s, X_s^\rho)] \geq 0,$$

which implies that

$$\begin{aligned} & -\text{sign}(X_s^b - X_s^\rho) [b_1(s, X_s^b) - p_1 \circ \rho(s, X_s^\rho)] \\ & \leq -\text{sign}(X_s^b - X_s^\rho) [b_1(s, X_s^b) - p_1 \circ \rho(s, X_s^\rho)] + \text{sign}(X_s^b - X_s^\rho) [b_1(s, X_s^b) - b_1(s, X_s^\rho)] \\ & = -\text{sign}(X_s^b - X_s^\rho) [b_1(s, X_s^\rho) - p_1 \circ \rho(s, X_s^\rho)] \\ & \leq |b_1(s, X_s^\rho) - p_1 \circ \rho(s, X_s^\rho)|. \end{aligned}$$

As a consequence, the second term of (12) is bounded above by

$$\int_0^t |b_1(s, X_s^\rho) - p_1 \circ \rho(s, X_s^\rho)| ds.$$

Similarly the third term is bounded above by

$$\int_0^t |b_2(s, X_s^\rho) - p_2 \circ \rho(s, X_s^\rho)| ds.$$

Substituting these estimates back into (12) we get

$$\begin{aligned} |X_t^b - X_t^\rho| & \leq T\tilde{\gamma} + \int_0^t |b_1(s, X_s^\rho) - p_1 \circ \rho(s, X_s^\rho)| ds \\ & \quad + \int_0^t |b_2(s, X_s^\rho) - p_2 \circ \rho(s, X_s^\rho)| ds. \end{aligned}$$

As b_1, b_2 and ρ are in the space Λ_T , we know that $\|b_1\|_T + \|b_2\|_T + \|\rho\|_T < \infty$ so that $\|b_1 - \rho\|_T < \infty$ and $\|b_2 - \rho\|_T < \infty$. We directly deduce:

$$|X_t^b - X_t^\rho| \leq T\tilde{\gamma} + (\|b_1 - \rho\|_T + \|b_2 - \rho\|_T) \int_0^t (1 + (X_s^\rho)^{2q}) ds.$$

By using triangular inequality, we obtain:

$$|X_t^b|^{2n} \leq (|X_t^\rho| + |X_t^b - X_t^\rho|)^{2n} \leq 2^{2n} \left\{ |X_t^\rho|^{2n} + |X_t^b - X_t^\rho|^{2n} \right\}.$$

Consequently, we have:

$$\widehat{\eta}_{2n}^b(T) = \sup_{0 \leq t \leq T} \mathbb{E} \left[|X_t^b|^{2n} \right] \leq 2^{2n} \left(\widehat{\eta}_{2n}^\rho(T) + \sup_{0 \leq t \leq T} \mathbb{E} \left[|X_t^b - X_t^\rho|^{2n} \right] \right).$$

But, we can write

$$\begin{aligned} |X_t^b - X_t^\rho|^{2n} & \leq 2^{2n} \left\{ T^{2n} \tilde{\gamma}^{2n} + 2^{2n} (\|b_1 - \rho\|_T^{2n} + \|b_2 - \rho\|_T^{2n}) \left[\int_0^T (1 + |X_t^\rho|^{2q}) dt \right]^{2n} \right\} \\ & \leq 2^{2n} \left\{ T^{2n} \tilde{\gamma}^{2n} + 2^{4n} (\|b_1 - \rho\|_T^{2n} + \|b_2 - \rho\|_T^{2n}) \left[T^{2n} + \left(\int_0^T |X_t^\rho|^{2q} dt \right)^{2n} \right] \right\} \\ & \leq 2^{2n} \left\{ T^{2n} \tilde{\gamma}^{2n} + 2^{4n} (\|b_1 - \rho\|_T^{2n} + \|b_2 - \rho\|_T^{2n}) \left[T^{2n} + \int_0^T |X_t^\rho|^{4qn} dt \right] \right\}. \end{aligned}$$

By taking the expectation then the supremum over $[0, T]$, we find the formula for $\widehat{\eta}_{2n}^b(T)$. The same computations hold for the second diffusion. \square

Lemma 2.8. Γ is an application from F_T to F_T and

$$\|\Gamma b\|_T^F \leq C_0 \left(1 + \widehat{\eta_{2q}^b}(T) + \widehat{\xi_{2q}^b}(T)\right), \quad (13)$$

where C_0 is a positive constant.

Proof. Step 1. We first need to prove that $p_i \circ \Gamma(b)$ lives in $\Lambda_T^1 \cap \Lambda_T^2$ for any $1 \leq i \leq 4$. We will do so only for $i = 1$. As ∇F_{11} is increasing and continuous, we deduce that $p_1 \circ \Gamma(b)$ is continuous and increasing in x . It is also locally Lipschitz (uniformly in the time variable). Due to the assumptions on the potential F_{11} , we have for any $x \geq y$

$$\begin{aligned} p_1 \circ \Gamma(b)(t, x) - p_1 \circ \Gamma(b)(t, y) &= a \mathbb{E} [\nabla F_{11}(x - X_t^b) - \nabla F_{11}(y - X_t^b)] \\ &\geq a \mathbb{E} [\beta_{11}^1((x - X_t^b) - (y - X_t^b)) + \beta_{11}^0] , \\ &\geq a \beta_{11}^1(x - y) + a \beta_{11}^0 , \end{aligned}$$

for some real numbers $\beta_{11}^1, \beta_{11}^0$. The existence of these constants is ensured by Assumption (H7). Similarly we obtain the following estimates

$$\begin{aligned} p_2 \circ \Gamma(b)(t, x) - p_2 \circ \Gamma(b)(t, y) &\geq (1 - a) \beta_{12}^1(x - y) + (1 - a) \beta_{12}^0, \\ p_3 \circ \Gamma(b)(t, x) - p_3 \circ \Gamma(b)(t, y) &\geq a \beta_{21}^1(x - y) + a \beta_{21}^0, \\ p_4 \circ \Gamma(b)(t, x) - p_4 \circ \Gamma(b)(t, y) &\geq (1 - a) \beta_{22}^1(x - y) + (1 - a) \beta_{22}^0, \end{aligned}$$

for some real numbers $\beta_{12}^1, \beta_{21}^1, \beta_{22}^1$ and $\beta_{12}^0, \beta_{21}^0, \beta_{22}^0$. The existence of these constants is ensured by Assumptions (H7) and (H8). By taking $\xi_1 := \inf \{a \beta_{11}^1; a \beta_{21}^1; (1 - a) \beta_{12}^1; (1 - a) \beta_{22}^1\}$ and $\xi_0 := \inf \{a \beta_{11}^0; a \beta_{21}^0; (1 - a) \beta_{12}^0; (1 - a) \beta_{22}^0\}$ we obtain that $p_i \circ \Gamma(b)$ is in $\Lambda_T^1 \cap \Lambda_T^2$.

Step 2. We will now prove Inequality (13) (which, by the way, proves that $p_i \circ \Gamma(b)$ lives in Λ_T^3). By definition, we have:

$$\begin{aligned} \|p_1 \circ \Gamma(b)\|_T &:= a \sup_{x \in \mathbb{R}} \frac{|\mathbb{E} [\nabla F_{11}(x - X_t^b)]|}{1 + x^{2q}} \\ &\leq a \sup_{x \in \mathbb{R}} \frac{\mathbb{E} [|\nabla F_{11}(x - X_t^b)|]}{1 + x^{2q}} \\ &\leq a \sup_{x \in \mathbb{R}} \frac{C \left(1 + |x|^{2q} + \mathbb{E} [|X_t^b|^{2q}]\right)}{1 + x^{2q}} \\ &\leq a C \left(1 + \widehat{\eta_{2q}^b}(T)\right). \end{aligned} \quad (14)$$

Note that we have used Assumption (H7) to obtain the second inequality in the above estimates. By proceeding similarly, we obtain

$$\|p_2 \circ \Gamma(b)\|_T \leq (1 - a) C \left(1 + \widehat{\xi_{2q}^b}(T)\right), \quad (15)$$

$$\|p_3 \circ \Gamma(b)\|_T \leq a C \left(1 + \widehat{\eta_{2q}^b}(T)\right), \quad (16)$$

$$\text{and } \|p_4 \circ \Gamma(b)\|_T \leq (1 - a) C \left(1 + \widehat{\xi_{2q}^b}(T)\right). \quad (17)$$

As a consequence, we have

$$\|\Gamma b\|_T^F \leq C_0 \left(1 + \widehat{\eta_{2q}^b}(T) + \widehat{\xi_{2q}^b}(T)\right).$$

□

Lemma 2.9. Γ is continuous and satisfies

$$\|p_1 \circ \Gamma(b) - p_1 \circ \Gamma(c)\|_T \leq (\|b_1 - c_1\|_T + \|b_2 - c_2\|_T) \sqrt{T} C'_0 \left(\widehat{\eta_{4q}^b}(T), \widehat{\eta_{4q}^c}(T) \right), \quad (18)$$

$$\|p_2 \circ \Gamma(b) - p_2 \circ \Gamma(c)\|_T \leq (\|b_1 - c_1\|_T + \|b_2 - c_2\|_T) \sqrt{T} C'_0 \left(\widehat{\eta_{4q}^b}(T), \widehat{\eta_{4q}^c}(T) \right), \quad (19)$$

$$\|p_3 \circ \Gamma(b) - p_3 \circ \Gamma(c)\|_T \leq (\|b_3 - c_3\|_T + \|b_4 - c_4\|_T) \sqrt{T} C'_0 \left(\widehat{\xi_{4q}^b}(T), \widehat{\xi_{4q}^c}(T) \right), \quad (20)$$

$$\text{and } \|p_4 \circ \Gamma(b) - p_4 \circ \Gamma(c)\|_T \leq (\|b_3 - c_3\|_T + \|b_4 - c_4\|_T) \sqrt{T} C'_0 \left(\widehat{\xi_{4q}^b}(T), \widehat{\xi_{4q}^c}(T) \right), \quad (21)$$

where C'_0 is an increasing function for both variables.

Proof. Set $s \in [0; T]$ and $x \in \mathbb{R}$. By triangular inequality, we have

$$|p_1 \circ \Gamma(b)(s, x) - p_1 \circ \Gamma(c)(s, x)| \leq \mathbb{E} \left[|F'_{11}(x - X_s^b) - F'_{11}(x - X_s^c)| \right].$$

By the assumptions on F_{11} , we get:

$$|p_1 \circ \Gamma(b)(s, x) - p_1 \circ \Gamma(c)(s, x)| \leq C_q \mathbb{E} \left[\Delta_s(b, c) \left(1 + (\Delta_s^b(x))^{2q-2} + (\Delta_s^c(x))^{2q-2} \right) \right],$$

where $\Delta_s(b, c) := |X_s^b - X_s^c|$,

and $\Delta_s^b(x) := |x - X_s^b|$ for any $b \in F_T$.

As $(a + b)^{2q-2} \leq 2^{2q-2} (a^{2q-2} + b^{2q-2})$, we deduce:

$$|p_1 \circ \Gamma(b)(s, x) - p_1 \circ \Gamma(c)(s, x)| \leq 2^{2q} C_q (1 + x^{2q-2}) \mathbb{E} \left[\Delta_s(b, c) \left(1 + (X_s^b)^{2q-2} + (X_s^c)^{2q-2} \right) \right]$$

We remind that $(a + b + c)^2 \leq 3(a^2 + b^2 + c^2)$ and $1 + x^{2q-2} \leq 2(1 + x^{2q})$. Then, Cauchy-Schwarz inequality yields

$$\begin{aligned} |p_1 \circ \Gamma(b)(s, x) - p_1 \circ \Gamma(c)(s, x)| &\leq 3 \times 2^{2q+1} C_q (1 + |x|^{2q}) \sqrt{\mathbb{E} \left[(\Delta_s(b, c))^2 \right]} \\ &\quad \times \sqrt{1 + \widehat{\eta_{4q-4}^b}(T) + \widehat{\eta_{4q-4}^c}(T)}. \end{aligned} \quad (22)$$

By using Itô formula with the function $x \mapsto |x|^2$, we can write

$$\begin{aligned} \Delta_t(b, c)^2 &= -2 \int_0^t (X_s^b - X_s^c) (V'_1(X_s^b) - V'_1(X_s^c)) ds \\ &\quad - 2 \int_0^t (X_s^b - X_s^c) (b_1(s, X_s^b) - c_1(s, X_s^c)) ds \\ &\quad - 2 \int_0^t (X_s^b - X_s^c) (b_2(s, X_s^b) - c_2(s, X_s^c)) ds. \end{aligned}$$

The first term is less than $2\theta_1 \int_0^t \Delta_s(b, c)^2 ds$. Since the functions b_1 and b_2 are increasing, we deduce that the quantities $(X_s^b - X_s^c) (b_1(s, X_s^b) - b_1(s, X_s^c))$ and $(X_s^b - X_s^c) (b_2(s, X_s^b) - b_2(s, X_s^c))$ are nonnegative. This implies

$$\begin{aligned} &- \int_0^t (X_s^b - X_s^c) (b_1(s, X_s^b) - c_1(s, X_s^c)) ds \\ &\leq \int_0^t |X_s^b - X_s^c| |b_1(s, X_s^c) - c_1(s, X_s^c)| ds \\ &\leq \frac{1}{2} \int_0^t \Delta_s(b, c)^2 ds + \frac{1}{2} \|b_1 - c_1\|_T^2 \int_0^t (1 + |X_s^c|^{2q})^2 ds \end{aligned}$$

$$\leq \frac{1}{2} \int_0^t \Delta_s(b, c)^2 ds + \|b_1 - c_1\|_T^2 \int_0^t (1 + |X_s^c|^{4q}) ds.$$

In the same way, we have

$$\begin{aligned} & - \int_0^t (X_s^b - X_s^c) (b_2(s, X_s^b) - c_2(s, X_s^c)) ds \\ & \leq \int_0^t |X_s^b - X_s^c| |b_2(s, X_s^c) - c_2(s, X_s^c)| ds \\ & \leq \frac{1}{2} \int_0^t \Delta_s(b, c)^2 ds + \frac{1}{2} \|b_2 - c_2\|_T^2 \int_0^t (1 + |X_s^c|^{2q})^2 ds \\ & \leq \frac{1}{2} \int_0^t \Delta_s(b, c)^2 ds + \|b_2 - c_2\|_T^2 \int_0^t (1 + |X_s^c|^{4q}) ds. \end{aligned}$$

We thus obtain:

$$\mathbb{E} [\Delta_t(b, c)^2] \leq 2(\theta + 1) \int_0^t \mathbb{E} [\Delta_s(b, c)^2] ds + 2 (\|b_1 - c_1\|_T^2 + \|b_2 - c_2\|_T^2) T \left(1 + \widehat{\eta_{4q}^c}(T)\right).$$

We apply Grönwall lemma and we get:

$$\mathbb{E} [\Delta_t(b, c)^2] \leq 2 (\|b_1 - c_1\|_T^2 + \|b_2 - c_2\|_T^2) T \left(1 + \widehat{\eta_{4q}^c}(T)\right) e^{2(\theta+1)t}.$$

As the role of b and c can be inverted, we obtain:

$$\mathbb{E} [\Delta_t(b, c)^2] \leq 2 (\|b_1 - c_1\|_T^2 + \|b_2 - c_2\|_T^2) T \left(1 + \frac{1}{2} \widehat{\eta_{4q}^b}(T) + \frac{1}{2} \widehat{\eta_{4q}^c}(T)\right) e^{2(\theta+1)t}.$$

We combine this with Inequality (22) and we finally have (18) with the function

$$C'_0(x, y) := 3 \times 2^{2q+\frac{3}{2}} C_q \sqrt{1 + \frac{|x| + |y|}{2}} \sqrt{1 + |x|^{\frac{4q-4}{4q}} + |y|^{\frac{4q-4}{4q}}}.$$

We obtain Inequalities (19), (20) and (21) by proceeding similarly. \square

As mentioned previously, we will use a fixed point theorem. We already have a continuous map. We will now restrict the space so that the map is a contraction.

Definition 2.10. Set $K > 0$ and $T > 0$. We consider

$$\Lambda_T^K := \{b \in \Lambda_T : \|b\|_T \leq K\}.$$

We also define $F_T^K := \Lambda_T^K \times \Lambda_T^K \times \Lambda_T^K \times \Lambda_T^K$.

Lemma 2.11. Let X_0 and Y_0 be two random variables such that $\mathbb{E} [X_0^{8q^2}] < \infty$ and $\mathbb{E} [Y_0^{8q^2}] < \infty$. Then, there exist two positive parameters K and T_0 such that for any $T < T_0$, we have the two following properties:

1. F_T^K is stable by Γ : $\Gamma F_T^K \subset F_T^K$.
2. The Lipschitz norm of the restriction of Γ on F_T^K is less than $\frac{1}{2}$.

Proof. Step 1. From (14), we have

$$\|p_1 \circ \Gamma(b)\|_T \leq aC \left(1 + \widehat{\eta_{2q}^b}(T)\right).$$

So, from Lemma 2.7, we have

$$\begin{aligned}
\|p_1 \circ \Gamma(b)\|_T &\leq aC \left(1 + k_1(q) \left[T^{2q} + \left(\|b_1 - \rho_0\|_T^{2q} + \|b_2 - \rho_0\|_T^{2q} \right) \left(T^{2q} + \widehat{\eta_{4q^2}^\rho(T)} \right) \right] \right) \\
&\leq aC \left(1 + k_1(q) \left[T^{2q} + 2^{2q} \left(\|b_1\|_T^{2q} + \|b_2\|_T^{2q} + 2\|\rho_0\|_T^{2q} \right) \left(T^{2q} + \widehat{\eta_{4q^2}^\rho(T)} \right) \right] \right) \\
&\leq aC \left(1 + k_1(q) \left[T^{2q} + 2^{2q+1} \left(K^{2q} + \|\rho_0\|_T^{2q} \right) \left(T^{2q} + \widehat{\eta_{4q^2}^\rho(T)} \right) \right] \right) \\
&\leq C \left(1 + k_1(q) \left[T^{2q} + 2^{2q+1} \left(K^{2q} + \|\rho_0\|_T^{2q} \right) \left(T^{2q} + \widehat{\eta_{4q^2}^\rho(T)} \right) \right] \right).
\end{aligned}$$

This can be rewritten as

$$\|p_1 \circ \Gamma(b)\|_T \leq C_1 + C_2 T^{2q} (1 + K^{2q}),$$

where C_1 and C_2 do not depend on T nor on K . We take $K \geq 2C_1$ and $T_0 \leq \left(\frac{C_1}{C_2(1+K^{2q})} \right)^{\frac{1}{2q}}$. As a consequence, for any $T < T_0$, we have

$$\|p_1 \circ \Gamma(b)\|_T \leq 2C_1 \leq K,$$

which proves that $p_1 \circ \Gamma(b) \in \Lambda_T^K$. We proceed similarly and we obtain that $p_2 \circ \Gamma(b) \in \Lambda_T^K$, $p_3 \circ \Gamma(b) \in \Lambda_T^K$ and $p_4 \circ \Gamma(b) \in \Lambda_T^K$. Consequently, $\Gamma(b) \in F_T^K$ if $b \in F_T^K$.

Step 2. We will now examine the Lipschitz constant. By making the sum of the inequalities in Lemma 2.9, we obtain

$$\|\Gamma(b) - \Gamma(c)\|_T^F \leq \alpha(T) \|b - c\|_T^F,$$

with $\alpha(T) := \max \left\{ 2\sqrt{T}C'_0 \left(\widehat{\eta_{4q}^b(T)}, \widehat{\eta_{4q}^c(T)} \right); 2\sqrt{T}C'_0 \left(\widehat{\xi_{4q}^b(T)}, \widehat{\xi_{4q}^c(T)} \right) \right\}$. We choose T_2 sufficiently small such that $\alpha(T_2) \leq \frac{1}{2}$. By taking $T := \min\{T_1; T_2\}$, the Lipschitz norm is less than $\frac{1}{2}$. \square

We point out that T depends on X_0 and Y_0 . This is why we will only be able to construct, in a first time, a solution on a finite time interval.

Proposition 2.12. Let X_0 and Y_0 be two random variables such that $\mathbb{E} \left[X_0^{8q^2} \right] < \infty$ and $\mathbb{E} \left[Y_0^{8q^2} \right] < \infty$. Then there exists $T_0 > 0$ such that for any $T < T_0$, the system of equations (1) admits a strong solution on the interval $[0; T]$. Moreover, we have

$$\sup_{0 \leq t \leq T} \mathbb{E} \left\{ |X_t|^{4q} \right\} + \sup_{0 \leq t \leq T} \mathbb{E} \left\{ |Y_t|^{4q} \right\} < \infty.$$

Proof. Step 1. We take K and T_0 as defined in Lemma 2.11. Thus, the Lipschitz norm of the restriction of Γ on F_T^K is smaller than $\frac{1}{2}$ for any $T < T_0$.

We take $b \in F_T^K$. We consider the sequence $(b_p)_{p \in \mathbb{N}}$ by $b_0 := b$ and $b_{p+1} := \Gamma(b_p)$ for any $p \in \mathbb{N}$.

We know that $b_p \in F_T^K$ for any $p \in \mathbb{N}$. Γ being a contraction, the sequence $(b_p)_p$ converges to an element $b_\infty \in F_T^K$. This element does not depend on b . Moreover, we have $\Gamma(b_\infty) = b_\infty$.

Consequently, $(X_t^{b_\infty}, Y_t^{b_\infty})_{t \in [0; T]}$ is a strong solution of the system (1) providing that $b_{\infty,1}, b_{\infty,2},$

$b_{\infty,3}$ and $b_{\infty,4}$ are locally Lipschitz (with $(b_{\infty,1}, b_{\infty,2}, b_{\infty,3}, b_{\infty,4}) =: b_\infty$).

As $b_{n+1} = \Gamma(b_n)$, then for $|x| \leq N$ and $|y| \leq N$, we have:

$$\begin{aligned}
|b_{n+1,1}(t, x) - b_{n+1,1}(t, y)| &= a \left| \mathbb{E} \left[F'_{11} \left(x - X_t^{b_n} \right) - F'_{11} \left(y - X_t^{b_n} \right) \right] \right| \\
&\leq a \mathbb{E} \left[\left| F'_{11} \left(x - X_t^{b_n} \right) - F'_{11} \left(y - X_t^{b_n} \right) \right| \right] \\
&\leq 2^{2q-1} a |x - y| \mathbb{E} \left[c + |x|^{2q-1} + |y|^{2q-1} + 2 \left| X_t^{b_n} \right| \right]
\end{aligned}$$

$$\leq k(N)|x - y| \left(1 + \widehat{\eta_{2q-1}^{b_n}}(T) \right).$$

Since $\|b_{n,1}\|_T \leq K$ and $\|b_{n,2}\|_T \leq K$, from Lemma 2.7, we deduce:

$$|b_{n+1,1}(t, x) - b_{n+1,1}(t, y)| \leq \psi(N, K, T, \rho) |x - y|.$$

By taking the limit as n goes to infinity, we deduce that

$$|b_{\infty,1}(t, x) - b_{\infty,1}(t, y)| \leq \psi(N, K, T, \rho) |x - y|.$$

So $b_{\infty,1}$ is locally Lipschitz. We can do the same reasoning for $b_{\infty,2}$, $b_{\infty,3}$ and $b_{\infty,4}$. Therefore, $(X_t^{b_\infty}, Y_t^{b_\infty})_{t \in [0; T]}$ is a strong solution of (1).

Step 2. According to the assumptions, $\mathbb{E}[X_0^{8q^2}]$ and $\mathbb{E}[Y_0^{8q^2}]$ are finite. We thus deduce

$$\sup_{s \in [0; T]} \mathbb{E}[|X_s^\rho|^{8q^2}] + \sup_{s \in [0; T]} \mathbb{E}[|Y_s^\rho|^{8q^2}] < \infty.$$

From Lemma 2.7, we have:

$$\widehat{\eta_{2n}^b}(T) \leq k_1(n) \left[T^{2n} + (\|b_1 - \rho_0\|_T^{2n} + \|b_2 - \rho_0\|_T^{2n}) \left(T^{2n} + \widehat{\eta_{4qn}^\rho}(T) \right) \right],$$

and

$$\widehat{\xi_{2n}^b}(T) \leq k_2(n) \left[T^{2n} + (\|b_3 - \rho_0\|_T^{2n} + \|b_4 - \rho_0\|_T^{2n}) \left(T^{2n} + \widehat{\xi_{4qn}^\rho}(T) \right) \right].$$

As $\widehat{\eta_{8q^2}^\rho}(T)$ and $\widehat{\xi_{4qn}^\rho}(T)$ are finite, we deduce the finiteness of $\widehat{\eta_{2n}^{b_\infty}}(T)$ and of $\widehat{\xi_{2n}^{b_\infty}}(T)$ for any n such that $4qn \leq 8q^2$. Consequently, we have $\widehat{\eta_{4q}^{b_\infty}}(T) + \widehat{\xi_{4q}^{b_\infty}}(T) < \infty$. \square

Let us point out that the uniqueness of the solution has not been proved for the moment. It will be proved subsequently.

We just obtained the result in finite time. We aim to establish it on the whole set \mathbb{R}_+ . To this end, we will assume that there exists a maximal time such that after this time, there is explosion. We will give a good control of the moments and then extend the solution after the maximal time. Thus we will obtain a contradiction proving that there is no such maximal time.

Lemma 2.13. Let X_0 and Y_0 be two random variables such that $\mathbb{E}[X_0^{2k}] < \infty$ and $\mathbb{E}[Y_0^{2k}] < \infty$ with $k > q$. Let T be a positive real and b an element of F_T . We assume that the function $\Gamma(b)$ is defined (which is not obvious since we did not assume the finiteness of the $8q^2$ -th moment) and that it satisfies $\Gamma(b) = b$. We put $X_t := X_t^b$ and $Y_t := Y_t^b$ the strong solution of the system starting from X_0 with the drift defined by b . Then, there exists a function C'' such that

$$\widehat{x_T} := \sup_{t \in [0; T]} \mathbb{E}\{|X_t|^{2k}\} \leq C'' \left(\mathbb{E}[|X_0|^{2k}]; \mathbb{E}[|Y_0|^{2k}] \right),$$

and

$$\widehat{y_T} := \sup_{t \in [0; T]} \mathbb{E}\{|Y_t|^{2k}\} \leq C'' \left(\mathbb{E}[|X_0|^{2k}]; \mathbb{E}[|Y_0|^{2k}] \right).$$

Proof. Step 1. We put $x_t := \mathbb{E}\{|X_t|^{2k}\}$ and $y_t := \mathbb{E}\{|Y_t|^{2k}\}$. We apply Itô formula, we take the integration, the expectation then we derive:

$$\begin{aligned} \frac{d}{dt} x_t &= -2k \mathbb{E} \left[\text{sign}(X_t) |X_t|^{2k-1} (V_1'(X_t) + F_{11}' * \mu_t(X_t) + F_{12}' * \nu_t(X_t)) \right] \\ &\quad + k(2k-1) \sigma^2 \mathbb{E} [|X_t|^{2k-2}]. \end{aligned}$$

Since F_{11} is convex (indeed, F'_{11} is odd and increasing due to (H7)), it is easy to prove that $\mathbb{E} \left[\text{sign}(X_t) |X_t|^{2k-1} F'_{11} * \mu_t(X_t) \right] \geq 0$. We deduce

$$\begin{aligned} \frac{d}{dt} x_t &\leq -2k \mathbb{E} \left[\text{sign}(X_t) |X_t|^{2k-1} (V'_1(X_t) + F'_{12} * \nu_t(X_t)) \right] \\ &\quad + k(2k-1) \sigma^2 x_t^{1-\frac{1}{k}}. \end{aligned}$$

Step 2. We now prove that the right hand side of the inequality is negative if x_t and y_t are too large. If V_1 was convex together with $V'_1(0) = 0$, the integral term with V'_1 would be easy to control since it would be nonnegative. Indeed, in this setting, we would have $xV'_1(x) = (x-0)(V'_1(x) - V'_1(0)) \geq 0$. However, V_1 is not convex. We take $\tau > 0$ arbitrarily large and we have:

$$\begin{aligned} \mathbb{E} \left[\text{sign}(X_t) |X_t|^{2k-1} V'_1(X_t) \right] &= \int_{-\tau}^{\tau} \text{sign}(x) |x|^{2k-1} V'_1(x) \mu_t(dx) \\ &\quad + \int_{[-\tau; \tau]^c} \text{sign}(x) |x|^{2k-1} V'_1(x) \mu_t(dx). \end{aligned}$$

If τ is large enough, the second integral is positive (due to Assumption (H5)) whilst the first may be negative (since the integrand is not uniformly nonnegative with respect to $x \in [-\tau; \tau]$).

Step 2.1. We begin by the first integral:

$$\begin{aligned} \int_{-\tau}^{\tau} \text{sign}(x) |x|^{2k-1} V'_1(x) \mu_t(dx) &\geq - \int_{-\tau}^{\tau} |x|^{2k-1} |V'_1(x)| \mu_t(dx) \\ &\geq -|\tau|^{2k-1} \sup_{x \in [-\tau; \tau]} |V'_1(x)| =: -f(\tau). \end{aligned}$$

Step 2.2. We now look at the second integral. Since $V''_1(\pm\infty) = +\infty$, we know that $g(\tau) := \inf_{x \in [-\tau; \tau]^c} \text{sign}(x) \frac{V'_1(x)}{|x|} > 0$ if τ is large enough. Thus:

$$\begin{aligned} \int_{[-\tau; \tau]^c} \text{sign}(x) |x|^{2k-1} V'_1(x) \mu_t(dx) &\geq g(\tau) \int_{[-\tau; \tau]^c} |x|^{2k} \mu_t(dx) \\ &\geq g(\tau) \left(\int_{\mathbb{R}} |x|^{2k} \mu_t(dx) - \int_{[-\tau; \tau]} |x|^{2k} \mu_t(dx) \right) \\ &\geq g(\tau) (x_t - \tau^{2k}). \end{aligned}$$

Step 2.3. We now control the mixed term $-2k \mathbb{E} \left[\text{sign}(X_t) |X_t|^{2k-1} F'_{12} * \nu_t(X_t) \right]$. We take \tilde{Y}_t an independent copy of Y_t . Then, we have:

$$\begin{aligned} -2k \mathbb{E} \left[\text{sign}(X_t) |X_t|^{2k-1} F'_{12} * \nu_t(X_t) \right] &= -2k \mathbb{E} \left[\text{sign}(X_t) |X_t|^{2k-1} F'_{12} (X_t - \tilde{Y}_t) \right] \\ &\leq 2k \mathbb{E} \left[|X_t|^{2k-1} \left| F'_{12} (X_t - \tilde{Y}_t) \right| \right]. \end{aligned}$$

Since $F'_{12}(0) = 0$ and F'_{12} is Lipschitz, we obtain:

$$\begin{aligned} -2k \mathbb{E} \left[\text{sign}(X_t) |X_t|^{2k-1} F'_{12} * \nu_t(X_t) \right] &\leq 2kC \mathbb{E} \left[|X_t|^{2k-1} |X_t - \tilde{Y}_t| \right] \\ &\leq 2kC \left\{ \mathbb{E} \left[|X_t|^{2k} \right] + \mathbb{E} \left[|X_t|^{2k-1} \right] \mathbb{E} [|Y_t|] \right\} \\ &\leq 2kC x_t + 2kC x_t^{1-\frac{1}{2k}} y_t^{\frac{1}{2k}}. \end{aligned}$$

Step 3. We combine the inequalities and we get:

$$\frac{d}{dt} x_t \leq k(2k-1) \sigma^2 x_t^{1-\frac{1}{k}} + 2kf(\tau) - 2kg(\tau)x_t$$

$$+ 2kg(\tau)\tau^{2k} + 2kCx_t + 2kCx_t^{1-\frac{1}{2k}}y_t^{\frac{1}{2k}}.$$

In the same way, we have:

$$\begin{aligned} \frac{d}{dt}y_t &\leq k(2k-1)\sigma^2y_t^{1-\frac{1}{k}} + 2kf(\tau) - 2kg(\tau)y_t \\ &\quad + 2kg(\tau)\tau^{2k} + 2kCy_t + 2kCy_t^{1-\frac{1}{2k}}x_t^{\frac{1}{2k}}. \end{aligned}$$

If $y_t \leq x_t$, we thus have

$$\begin{aligned} \frac{d}{dt}x_t &\leq k(2k-1)\sigma^2x_t^{1-\frac{1}{k}} + 2kf(\tau) - 2kg(\tau)x_t \\ &\quad + 2kg(\tau)\tau^{2k} + 4kCx_t. \end{aligned}$$

By taking τ large enough, $g(\tau) > 4C$ so that if x_t is larger than a constant $\chi(\tau)$, $\frac{d}{dt}x_t \leq 0$. Conversely, if $x_t \leq y_t$, if y_t is larger than $\chi(\tau)$, $\frac{d}{dt}y_t \leq 0$. We immediately deduce:

$$\widehat{x}_T \leq \max \left\{ \chi(\tau); \mathbb{E} \left[|X_0|^{2k} \right]; \mathbb{E} \left[|Y_0|^{2k} \right] \right\}$$

and

$$\widehat{y}_T \leq \max \left\{ \chi(\tau); \mathbb{E} \left[|X_0|^{2k} \right]; \mathbb{E} \left[|Y_0|^{2k} \right] \right\}$$

□

We are now able to obtain the main theorem

Theorem 2.1. Set two random variables X_0 and Y_0 such that $\mathbb{E} \left[X_0^{2q} \right] < \infty$ and $\mathbb{E} \left[Y_0^{2q} \right] < \infty$. Then, the system admits a unique strong solution on \mathbb{R}_+ .

Proof. Step 1. We consider

$$U := \sup \left\{ T > 0 : \text{(E) admits a unique solution on } [0; T], \sup_{0 \leq t \leq T} \mathbb{E} \left[X_t^{8q^2} \right] + \sup_{0 \leq t \leq T} \mathbb{E} \left[Y_t^{8q^2} \right] < \infty \right\}$$

with the convention $\sup \emptyset = 0$. We begin to show that $U > 0$. By taking K large enough, there exists $T > 0$ and a unique $b \in F_T^K$ such that $\Gamma(b) = b$, by Lemma 2.11. Then (X^b, Y^b) is a strong solution of the system (E) on $[0; T]$. We now consider a solution $(\widetilde{X}_t, \widetilde{Y}_t)_{t \in [0; T]}$. To this solution, we associate the following drifts

$$\begin{aligned} c_1(t, x) &:= a\mathbb{E} \left[F'_{11}(x - \widetilde{X}_t) \right], \quad c_2(t, x) := (1-a)\mathbb{E} \left[F'_{12}(x - \widetilde{Y}_t) \right], \\ c_3(t, x) &:= a\mathbb{E} \left[F'_{21}(x - \widetilde{X}_t) \right] \quad \text{and} \quad c_4(t, x) := (1-a)\mathbb{E} \left[F'_{22}(x - \widetilde{Y}_t) \right]. \end{aligned}$$

We put $c := (c_1, c_2, c_3, c_4)$. By the assumptions on F_{11} , we obtain:

$$\frac{|c_1(t, x)|}{1 + x^{2q}} = a \frac{\left| F'_{11}(x - \widetilde{X}_t) \right|}{1 + x^{2q}} \leq C \left(1 + \sup_{0 \leq t \leq T} \mathbb{E} \left[|\widetilde{X}_t|^{2q} \right] \right).$$

In the same way, we have

$$\begin{aligned} \frac{|c_2(t, x)|}{1 + x^{2q}} &\leq C \left(1 + \sup_{0 \leq t \leq T} \mathbb{E} \left[|\widetilde{Y}_t|^{2q} \right] \right), \\ \frac{|c_3(t, x)|}{1 + x^{2q}} &\leq C \left(1 + \sup_{0 \leq t \leq T} \mathbb{E} \left[|\widetilde{X}_t|^{2q} \right] \right), \end{aligned}$$

$$\text{and } \frac{|c_4(t, x)|}{1+x^{2q}} \leq C \left(1 + \sup_{0 \leq t \leq T} \mathbb{E} \left[|\tilde{Y}_t|^{2q} \right] \right).$$

However, according to Lemma 2.13, the moment at time t of \widetilde{X}_t and the one of \tilde{Y}_t are bounded by a function which depends on the moments of X_0 and Y_0 .

If we have a function c such that $\Gamma(c) = c$ then the associated process verifies the equations of Lemma 2.13. Then, by taking K large enough, we deduce that $\frac{|c_1(t, x)|}{1+x^{2q}} \leq K$, $\frac{|c_2(t, x)|}{1+x^{2q}} \leq K$, $\frac{|c_3(t, x)|}{1+x^{2q}} \leq K$ and $\frac{|c_4(t, x)|}{1+x^{2q}} \leq K$. This means that $c \in F_T^K$. Consequently, for any random variables X_0 and Y_0 with $8q^2$ th moment finite, by taking K large enough (which depends on the initial moments), we know that a solution of the equation $\Gamma(c) = c$ is in F_T^K . However, the equation has a unique solution on F_T^K since the map Γ is a contraction on F_T^K .

We immediately deduce that there is a unique system of stochastic differential equations which corresponds to (E). And, there is a unique strong solution to this equation. We thus deduce that $U \geq T > 0$.

Step 2. We assume that $U < \infty$. This U does depend on the moments of X_0 and Y_0 . We know that the moments of order 1 to $8q^2$ of (X_t, Y_t) are bounded by a constant C''_0 which depends only on the initial moments. We thus consider the system of equations

$$\begin{aligned} X'_t &= X'_0 - \int_0^t \nabla V_1(X'_s) ds - a \int_0^t (\nabla F_{11} * \mu_s)(X'_s) ds - (1-a) \int_0^t (\nabla F_{12} * \nu_s)(X'_s) ds + \sigma B_t, \\ Y'_t &= Y'_0 - \int_0^t \nabla V_2(Y'_s) ds - a \int_0^t (\nabla F_{21} * \mu_s)(Y'_s) ds - (1-a) \int_0^t (\nabla F_{22} * \nu_s)(Y'_s) ds + \sigma \tilde{B}_t, \end{aligned}$$

with X'_0 and Y'_0 such that $\mathbb{E}[|X'_0|^r] \leq C''_0$ and $\mathbb{E}[|Y'_0|^r] \leq C''_0$. We can associate a time $T' > 0$ to this equation such that it admits a unique strong solution on $[0; T']$. To the new random variable X'_0 is associated a new constant K' . Without any change to the generality, we take $K' \geq K$.

We put $X'_0 := X_{U-\frac{T'}{2}}$ and $Y'_0 := Y_{U-\frac{T'}{2}}$. These new initial random variables satisfy the conditions so we can define a unique strong solution on $[0; T']$. This implies that we have extended $(X_t, Y_t)_{t \in [0; U]}$ to $(X_t, Y_t)_{t \in [0; U+\frac{T'}{2}]}$. Indeed, on $[U-\frac{T'}{2}; U]$, there is uniqueness. This contradicts the definition of U .

□

By using the proof of Theorem 2.1, we can directly obtain the following result:

Proposition 2.14. Let (X, Y) be a solution of the system (1). Assume that $\mathbb{E}(X_0^{2n}) + \mathbb{E}(Y_0^{2n}) < \infty$ for some $n \in \mathbb{N}^*$. Then, we have

$$\sup_{t \geq 0} \mathbb{E}(X_t^{2n}) + \sup_{t \geq 0} \mathbb{E}(Y_t^{2n}) < \infty.$$

3 Propagation of chaos

We recall the interacting particle system defined in (6):

$$\begin{aligned} dX_t^i &= -\nabla V_1(X_t^i) dt - \frac{1}{N_n + M_n} \sum_{j=1}^{N_n} \nabla F_{11}(X_t^i - X_t^j) dt \\ &\quad - \frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \nabla F_{12}(X_t^i - Y_t^k) dt + \sigma dW_t^i, \quad i = 1, \dots, N_n; \\ dY_t^i &= -\nabla V_2(Y_t^i) dt - \frac{1}{N_n + M_n} \sum_{j=1}^{N_n} \nabla F_{21}(Y_t^i - X_t^j) dt \end{aligned}$$

$$- \frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \nabla F_{22}(Y_t^i - Y_t^k) dt + \sigma d\widetilde{W}_t^i, \quad i = 1, \dots, M_n;$$

and its identically independent copies given in (8)

$$\begin{cases} d\widehat{X}_t^i = -\nabla V_1(\widehat{X}_t^i) dt - a(\nabla F_{11} * \mu_t)(\widehat{X}_t^i) dt - (1-a)(\nabla F_{12} * \nu_t)(\widehat{X}_t^i) dt + \sigma dW_t^i \\ \quad i = 1, \dots, N_n; \\ d\widehat{Y}_t^i = -\nabla V_2(\widehat{Y}_t^i) dt - a(\nabla F_{21} * \mu_t)(\widehat{Y}_t^i) dt - (1-a)(\nabla F_{22} * \nu_t)(\widehat{Y}_t^i) dt + \sigma d\widetilde{W}_t^i, \\ \quad i = 1, \dots, M_n, \end{cases}$$

where $a = \lim_{n \rightarrow \infty} \frac{N_n}{N_n + M_n}$ and $\mu_t = \text{Law}(\widehat{X}_t^i)$, $\nu_t = \text{Law}(\widehat{Y}_t^i)$.

The rest of this section is devoted to prove Theorem 1.2 that is to show that the interacting particle system satisfies propagation of chaos. We adapt the proof of [BRTV98, Her03]. In Proposition 3.2 we prove a weaker statement than (7) where the expectation and the supremum are interchanged. We then strengthens Proposition 3.2 to the fourth power in Proposition 3.4. Finally, Theorem 1.2 will be derived from these propositions.

We will need the following lemma on a nonlinear generalisation of Grönwall's inequality.

Lemma 3.1. *Let ϕ be a positive function such that $\phi(0) = 0$. Suppose that there exist constants $A > 0, B \geq 0$ and $0 \leq \alpha < 1$ such that*

$$\phi(t) \leq A \int_0^t \phi(s) ds + B \int_0^t \phi(s)^\alpha ds,$$

then

$$\phi(t) \leq \left(\frac{B}{A} (e^{(1-\alpha)At} - 1) \right)^{\frac{1}{1-\alpha}}.$$

Proof. We note that a special case of this lemma for $\alpha = \frac{1}{2}$ has appeared in [Her03, Lemma 2] while a more general version where A and B are functions of s can be found in [Dra03, Theorem 21]. For the convenience of the reader we provide a simplified proof for the case of constant coefficients and arbitrary α here. Suppose ψ solve the integral equation

$$\psi(t) = A \int_0^t \psi(s) ds + B \int_0^t \psi(s)^\alpha ds \quad \psi(0) = 0.$$

We take the derivative with respect to t both sides to obtain

$$\frac{d\psi(t)}{A\psi(t) + B\psi(t)^\alpha} = dt, \quad \psi(0) = 0.$$

Taking the anti-derivative of this ODE gives

$$\frac{1}{A(\alpha-1)} \log \left[\frac{\psi(t)^\alpha}{A\psi(t) + B\psi(t)^\alpha} \right] = t + C.$$

Solving this equation with the initial data $\psi(0) = 0$ we obtain

$$\psi(t) = \left(\frac{B}{A} (e^{A(1-\alpha)t} - 1) \right)^{\frac{1}{1-\alpha}}.$$

A comparison principle gives $\phi(t) \leq \psi(t)$, which is the assertion of the lemma. \square

Proposition 3.2. *We have*

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \mathbb{E} \left[(X_t^i - \widehat{X}_t^i)^2 \right] = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \mathbb{E} \left[(Y_t^i - \widehat{Y}_t^i)^2 \right] = 0. \quad (24)$$

Proof. We define

$$\omega(t) := \mathbb{E} \left[(X_t^1 - \widehat{X}_t^1)^2 \right] \quad \text{and} \quad \widehat{\omega}(t) := \mathbb{E} \left[(Y_t^1 - \widehat{Y}_t^1)^2 \right]. \quad (25)$$

We have

$$\omega(t) = \mathbb{E} \left[(X_t^i - \widehat{X}_t^i)^2 \right] \quad \forall i = 1, \dots, N_n \quad \text{and} \quad \widehat{\omega}(t) = \mathbb{E} \left[(Y_t^i - \widehat{Y}_t^i)^2 \right] \quad \forall i = 1, \dots, M_n. \quad (26)$$

Using the Itô formula, we compute

$$\begin{aligned} \omega(t) &= \mathbb{E} \left[(X_t^i - \widehat{X}_t^i)^2 \right] \\ &= -2\mathbb{E} \int_0^t (\nabla V_1(X_s^i) - \nabla V_1(\widehat{X}_s^i)) \cdot (X_s^i - \widehat{X}_s^i) ds \\ &\quad - 2\mathbb{E} \int_0^t \left[\frac{1}{N_n + M_n} \sum_{j=1}^{N_n} \nabla F_{11}(X_s^i - X_s^j) - a(\nabla F_{11} * \mu_s)(\widehat{X}_s^i) \right] \cdot (X_s^i - \widehat{X}_s^i) ds \\ &\quad - 2\mathbb{E} \int_0^t \left[\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \nabla F_{12}(X_s^i - Y_s^k) - (1-a)(\nabla F_{12} * \mu_s)(\widehat{X}_s^i) \right] \cdot (X_s^i - \widehat{X}_s^i) ds \\ &= 2 \int_0^t \mathbb{E} (A_i(s) + B_i(s) + C_i(s) + D_i(s) + E_i(s)) ds, \end{aligned} \quad (27)$$

where

$$\begin{aligned} A_i(t) &= -(\nabla V_1(X_t^i) - \nabla V_1(\widehat{X}_t^i)) \cdot (X_t^i - \widehat{X}_t^i), \\ B_i(t) &= \left[-\frac{1}{N_n + M_n} \sum_{j=1}^{N_n} (\nabla F_{11}(X_t^i - X_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i)) \right] \cdot (X_t^i - \widehat{X}_t^i), \\ C_i(t) &= \left(a - \frac{N_n}{N_n + M_n} \right) (\nabla F_{11} * \mu_t)(\widehat{X}_t^i) \cdot (X_t^i - \widehat{X}_t^i), \\ D_i(t) &= \left[-\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} (\nabla F_{12}(X_t^i - Y_t^k) - (\nabla F_{12} * \nu_t)(\widehat{X}_t^i)) \right] \cdot (X_t^i - \widehat{X}_t^i), \\ E_i(t) &= \left((1-a) - \frac{M_n}{N_n + M_n} \right) (\nabla F_{12} * \nu_t)(\widehat{X}_t^i) \cdot (X_t^i - \widehat{X}_t^i). \end{aligned}$$

Next we estimate each term in (27). We start with A_i :

$$\begin{aligned} A_i(t) &= -(\nabla V_1(X_t^i) - \nabla V_1(\widehat{X}_t^i)) \cdot (X_t^i - \widehat{X}_t^i) \\ &\leq \theta_1 (X_t^i - \widehat{X}_t^i)^2, \end{aligned}$$

where we have used Assumption (H3) to obtain the last inequality. This implies that

$$\sum_{i=1}^{N_n} \mathbb{E} [A_i(t)] \leq N_n \theta_1 \omega(t). \quad (28)$$

Next we estimate B_i :

$$\begin{aligned} B_i(t) &= \left[-\frac{1}{N_n + M_n} \sum_{j=1}^{N_n} (\nabla F_{11}(X_t^i - X_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i)) \right] \cdot (X_t^i - \widehat{X}_t^i) \\ &= \left[-\frac{1}{N_n + M_n} \sum_{j=1}^{N_n} (\nabla F_{11}(X_t^i - X_t^j) - \nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j)) \right] \cdot (X_t^i - \widehat{X}_t^i) \end{aligned}$$

$$\begin{aligned}
& - \left[\frac{1}{N_n + M_n} \sum_{j=1}^{N_n} \left(\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i) \right) \right] \cdot (X_t^i - \widehat{X}_t^i) \\
& = - \frac{1}{N_n + M_n} \sum_{j=1}^{N_n} \varrho_{ij}^1(t) - \frac{1}{N_n + M_n} \sum_{j=1}^{N_n} \varrho_{ij}^2(t),
\end{aligned} \tag{29}$$

where

$$\varrho_{ij}^1(t) := \left(\nabla F_{11}(X_t^i - X_t^j) - \nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) \right) \cdot (X_t^i - \widehat{X}_t^i), \tag{30a}$$

$$\varrho_{ij}^2(t) := \left(\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i) \right) \cdot (X_t^i - \widehat{X}_t^i). \tag{30b}$$

We have

$$\sum_{i=1}^{N_n} \sum_{j=1}^{N_n} \varrho_{ij}^1(t) = \sum_{1 \leq i < j \leq N_n} \varrho_{ij}^3(t),$$

where $\varrho_{ij}^3(t) = \varrho_{ij}^1(t) + \varrho_{ji}^1(t)$. Since ∇F_{11} is an odd function, we have

$$\begin{aligned}
\varrho_{ij}^3(t) &= \left(\nabla F_{11}(X_t^i - X_t^j) - \nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) \right) \cdot (X_t^i - \widehat{X}_t^i) \\
&\quad + \left(\nabla F_{11}(X_t^j - X_t^i) - \nabla F_{11}(\widehat{X}_t^j - \widehat{X}_t^i) \right) \cdot (X_t^j - \widehat{X}_t^j) \\
&= \left(\nabla F_{11}(X_t^i - X_t^j) - \nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) \right) \cdot \left((X_t^i - \widehat{X}_t^i) - (X_t^j - \widehat{X}_t^j) \right).
\end{aligned}$$

If $X_t^i - X_t^j \geq \widehat{X}_t^i - \widehat{X}_t^j$ (resp. $X_t^i - X_t^j \leq \widehat{X}_t^i - \widehat{X}_t^j$) then $X_t^i - \widehat{X}_t^i \geq X_t^j - \widehat{X}_t^j$ (resp. $X_t^i - \widehat{X}_t^i \leq X_t^j - \widehat{X}_t^j$) and $\nabla F_{11}(X_t^i - X_t^j) \geq \nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j)$ (resp. $\nabla F_{11}(X_t^i - X_t^j) \leq \nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j)$) as ∇F_{11} is increasing. Thus we always have $\varrho_{ij}^3(t) \geq 0$. Therefore,

$$\sum_{i,j=1}^{N_n} \varrho_{ij}^1(t) \geq 0. \tag{31}$$

On the other hand, using Cauchy-Schwarz inequality, we get

$$\mathbb{E} \left(\sum_{j=1}^{N_n} \varrho_{ij}^2(t) \right) \leq \left(\mathbb{E}((X_t^i - \widehat{X}_t^i)^2 \kappa_i(t)) \right)^{\frac{1}{2}}, \tag{32}$$

where

$$\kappa_i(t) = \mathbb{E} \left(\left[\sum_{j=1}^{N_n} (\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i)) \right]^2 \right).$$

We rewrite κ_i as

$$\begin{aligned}
\kappa_i(t) &= \sum_{j=1}^{N_n} \xi_{j,j}(t) + \sum_{1 \leq j < k \leq N_n} \xi_{j,k}(t) \quad \text{with} \\
\xi_{j,k}(t) &= \mathbb{E} \left([\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i)] [\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^k) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i)] \right).
\end{aligned}$$

If $j \neq k$, $\widehat{X}^i, \widehat{X}^j$ and \widehat{X}^k are three independent copies of \widehat{X}^1 . This implies that

$$\xi_{j,k} = \mathbb{E}_{\widehat{X}^i} \left(\mathbb{E}_{\widehat{X}^j} [\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i)] \mathbb{E}_{\widehat{X}^k} [\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^k) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i)] \right)$$

$$= \mathbb{E}_{\widehat{X}^i}[0] = 0,$$

where we have used the fact that $\widehat{X}^i, \widehat{X}^j$ and \widehat{X}^k have the same law μ_t . For $j = k$, we get

$$\begin{aligned} \xi_{j,j}(t) &= \mathbb{E} \left(\left[\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i) \right]^2 \right) \\ &\leq 2\mathbb{E} \left(|\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j)|^2 + |(\nabla F_{11} * \mu_t)(\widehat{X}_t^i)|^2 \right). \end{aligned} \quad (33)$$

Since $|\nabla F_{11}(x)| \leq C(1 + |x|^{2q})$, we have $|\nabla F_{11}(x)|^2 \leq C(1 + |x|^{4q})$. Applying this inequality we obtain

$$\begin{aligned} \mathbb{E}|\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j)|^2 &\leq C\mathbb{E}(1 + |\widehat{X}_t^i - \widehat{X}_t^j|^{4q}) \leq C\mathbb{E}(1 + |\widehat{X}_t^i|^{4q} + |\widehat{X}_t^j|^{4q}) \leq C, \\ \mathbb{E}(|(\nabla F_{11} * \mu_t)(\widehat{X}_t^i)|^2) &= \mathbb{E} \left(\left| \int \nabla F_{11}(\widehat{X}_t^i - y) \mu_t(y) dy \right|^2 \right) \leq \mathbb{E} \left(\int |\nabla F_{11}(\widehat{X}_t^i - y)|^2 \mu_t(dy) \right) \\ &\leq C\mathbb{E} \left(\int (1 + |\widehat{X}_t^i|^{4q} + |y|^{4q}) \mu_t(y) dy \right) \leq C\mathbb{E}(1 + |\widehat{X}_t^i|^{4q}) \leq C. \end{aligned} \quad (34)$$

Therefore, we obtain

$$\kappa_i(t) = \sum_{j=1}^{N_n} \xi_{j,j}(t) \leq CN_n$$

Substituting this estimate back into (32) gives

$$\mathbb{E} \left(\sum_{j=1}^{N_n} \rho_{ij}^2(t) \right) \leq CN_n^{\frac{1}{2}} \left(\mathbb{E}(X_t^i - \widehat{X}_t^i)^2 \right)^{\frac{1}{2}}. \quad (35)$$

Substituting (35) and (31) back into (29), we achieve

$$\sum_{i=1}^{N_n} \mathbb{E}[B_i(t)] \leq \frac{CN_n^{3/2}}{N_n + M_n} \left(\mathbb{E}(X_t^i - \widehat{X}_t^i)^2 \right)^{\frac{1}{2}} = \frac{CN_n^{3/2}}{N_n + M_n} \sqrt{\omega(t)}. \quad (36)$$

We proceed with estimating C_i . Using Cauchy-Schwarz inequality we get

$$\begin{aligned} |\mathbb{E}[C_i(t)]| &= \left| a - \frac{N_n}{N_n + M_n} \right| \mathbb{E}(\nabla F_{11} * \mu_t)(\widehat{X}_t^i)(X_t^i - \widehat{X}_t^i) \\ &\leq \left| a - \frac{N_n}{N_n + M_n} \right| \left(\mathbb{E}[X_t^i - \widehat{X}_t^i]^2 \right)^{\frac{1}{2}} \left(\mathbb{E}(|(\nabla F_{11} * \mu_t)(\widehat{X}_t^i)|^2) \right)^{\frac{1}{2}} \\ &\leq C \left| a - \frac{N_n}{N_n + M_n} \right| \left(\mathbb{E}[X_t^i - \widehat{X}_t^i]^2 \right)^{\frac{1}{2}}, \end{aligned}$$

where we have used $\mathbb{E}(|(\nabla F_{11} * \mu_t)(\widehat{X}_t^i)|^2) \leq C$ which was proved in (34). Therefore,

$$\sum_{i=1}^{N_n} \mathbb{E}C_i \leq CN_n \left| a - \frac{N_n}{N_n + M_n} \right| \left(\mathbb{E}[X_t^i - \widehat{X}_t^i]^2 \right)^{\frac{1}{2}} = CN_n \left| a - \frac{N_n}{N_n + M_n} \right| \sqrt{\omega(t)}. \quad (37)$$

Now we estimate D_i . This is a cross term that involves both species and we will need to use assumptions on the Lipschitz property of F_{12} . We first add and subtract appropriate terms similarly as in B_i .

$$D_i(t) = \left[-\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \left(\nabla F_{12}(X_t^i - Y_t^k) - (\nabla F_{12} * \nu_t)(\widehat{X}_t^i) \right) \right] \cdot (X_t^i - \widehat{X}_t^i)$$

$$\begin{aligned}
&= \left[-\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \left(\nabla F_{12}(X_t^i - Y_t^k) - \nabla F_{12}(\widehat{X}_t^i - Y_t^k) \right) \right] \cdot (X_t^i - \widehat{X}_t^i) \\
&\quad - \left[\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \left(\nabla F_{12}(\widehat{X}_t^i - Y_t^k) - (\nabla F_{12} * \nu_t)(\widehat{X}_t^i) \right) \right] \cdot (X_t^i - \widehat{X}_t^i) \\
&=: D_i^1(t) + D_i^2(t).
\end{aligned} \tag{38}$$

Using Assumption (H8) on the Lipschitz property of ∇F_{12} we have

$$|\nabla F_{12}(X_t^i - Y_t^k) - \nabla F_{12}(\widehat{X}_t^i - Y_t^k)| \leq K |X_t^i - \widehat{X}_t^i|,$$

where K is the Lipschitz constant for ∇F_{12} , which implies that

$$\mathbb{E}[D_i^1(t)] \leq \frac{CM_n}{N_n + M_n} \mathbb{E}(X_t^i - \widehat{X}_t^i)^2. \tag{39}$$

Taking the expectation of $D_i^2(t)$, noting that $\nu_t = \text{Law}(\widehat{Y}_t^k)$, we obtain

$$\mathbb{E}[D_i^2(t)] = \mathbb{E} \left[-\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \left(\nabla F_{12}(\widehat{X}_t^i - Y_t^k) - (\nabla F_{12}(\widehat{X}_t^i - \widehat{Y}_t^k)) \right) \right] \cdot (X_t^i - \widehat{X}_t^i).$$

Then similarly as in $D_i^1(t)$, we have

$$\mathbb{E}[D_i^2(t)] \leq \frac{C}{N_n + M_n} \sum_{k=1}^{M_n} \mathbb{E}(|Y_t^k - \widehat{Y}_t^k| |X_t^i - \widehat{X}_t^i|),$$

which implies that

$$\begin{aligned}
\sum_{i=1}^{N_n} \mathbb{E}[D_i^2(t)] &\leq \frac{C}{N_n + M_n} \mathbb{E} \left(\sum_{i=1}^{N_n} |X_t^i - \widehat{X}_t^i| \sum_{k=1}^{M_n} |Y_t^k - \widehat{Y}_t^k| \right) \\
&\leq \frac{C\sqrt{N_n}\sqrt{M_n}}{N_n + M_n} \left(\sum_{i=1}^{N_n} \mathbb{E}(X_t^i - \widehat{X}_t^i)^2 \right)^{\frac{1}{2}} \left(\sum_{k=1}^{M_n} \mathbb{E}(Y_t^k - \widehat{Y}_t^k)^2 \right)^{\frac{1}{2}} \\
&= \frac{CM_n N_n}{N_n + M_n} \left(\mathbb{E}(X_t^i - \widehat{X}_t^i)^2 \right)^{\frac{1}{2}} \left(\mathbb{E}(Y_t^k - \widehat{Y}_t^k)^2 \right)^{\frac{1}{2}} \\
&= \frac{CM_n N_n}{N_n + M_n} \sqrt{\omega(t)\widehat{\omega}(t)}.
\end{aligned} \tag{40}$$

Substituting (39) and (40) into (38) we obtain

$$\sum_{i=1}^{N_n} \mathbb{E}[D_i(t)] \leq \frac{CM_n N_n}{N_n + M_n} \left(\omega(t) + \sqrt{\omega(t)\widehat{\omega}(t)} \right). \tag{41}$$

Finally we estimate E_i analogously as in C_i and get

$$\mathbb{E}[E_i(t)] \leq C \left| (1-a) - \frac{M_n}{N_n + M_n} \right| \left(\mathbb{E}(X_t^i - \widehat{X}_t^i)^2 \right)^{\frac{1}{2}} = C \left| (1-a) - \frac{M_n}{N_n + M_n} \right| \sqrt{\omega(t)}.$$

Taking the sum over i from 1 to N_n yields

$$\sum_{i=1}^{N_n} \mathbb{E}[E_i(t)] \leq CN_n \left| (1-a) - \frac{M_n}{N_n + M_n} \right| \sqrt{\omega(t)}. \tag{42}$$

Substituting (28), (36), (41) and (42) into (27) we obtain

$$\begin{aligned} N_n \omega(t) \leq & 2 \int_0^t \left(N_n \theta_1 \omega(s) + \frac{C N_n^{3/2}}{N_n + M_n} \sqrt{\omega(s)} + C N_n \left| a - \frac{N_n}{N_n + M_n} \right| \sqrt{\omega(t)} + \frac{C M_n N_n}{N_n + M_n} \omega(s) \right. \\ & \left. + \frac{C M_n N_n}{N_n + M_n} \sqrt{\omega(s) \widehat{\omega}(s)} + C N_n \left| (1-a) - \frac{M_n}{N_n + M_n} \right| \sqrt{\omega(t)} \right) ds. \end{aligned}$$

By dividing both sides by N_n we get

$$\begin{aligned} \omega(t) \leq & C \int_0^t \left(\omega(s) + \left(\frac{N_n^{1/2}}{N_n + M_n} + \left| a - \frac{N_n}{N_n + M_n} \right| + \left| (1-a) - \frac{M_n}{N_n + M_n} \right| \right) \sqrt{\omega(s)} \right. \\ & \left. + \sqrt{\omega(s) \widehat{\omega}(s)} \right) ds \end{aligned} \quad (43)$$

$$\leq C \int_0^t \left(\omega(s) + \widehat{\omega}(s) + f(n) \sqrt{\omega(s) + \widehat{\omega}(s)} \right) ds, \quad (44)$$

where $0 \leq f(n) \leq C \frac{N_n^{1/2}}{N_n + M_n} + \left| a - \frac{N_n}{N_n + M_n} \right| + \left| (1-a) - \frac{M_n}{N_n + M_n} \right|$ (hence $f(n) \rightarrow 0$ as $n \rightarrow 0$). Note that we have used $\widehat{\omega}(s) \geq 0$ and the elementary $\sqrt{xy} \leq \frac{1}{2}(x+y)$ to obtain the last estimate.

Analogously we obtain

$$\widehat{\omega}(t) \leq C \int_0^t \left(\omega(s) + \widehat{\omega}(s) + \widehat{f}(n) \sqrt{\omega(s) + \widehat{\omega}(s)} \right) ds \quad (45)$$

for some function $0 \leq \widehat{f}(n)$ that tends to 0 as n goes to infinity.

Taking the sum of (43) and (45) yields

$$\omega(t) + \widehat{\omega}(t) \leq C \int_0^t \left(\omega(s) + \widehat{\omega}(s) + (f(n) + \widehat{f}(n)) \sqrt{\omega(s) + \widehat{\omega}(s)} \right) ds. \quad (46)$$

Applying Lemma 3.1, we obtain

$$\omega(t) + \widehat{\omega}(t) \leq C \left((f(n) + \widehat{f}(n)) \left(e^{\frac{Ct}{2}} - 1 \right) \right)^2. \quad (47)$$

Since $f(n) + \widehat{f}(n) \rightarrow 0$ as $n \rightarrow 0$, the last estimate implies that

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \omega(t) = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \widehat{\omega}(t) = 0.$$

This completes the proof of Proposition 3.2. \square

Remark 3.3. If N_n and M_n tend to $+\infty$ simultaneously but $\frac{N_n}{M_n}$ is a constant, then from the computations in the proof of Proposition 3.2, we obtain explicit estimates

$$\sup_{t \in [0, T]} \mathbb{E} (X_t^i - \widehat{X}_t^i)^2 \leq \frac{C}{N_n} \quad \text{and} \quad \sup_{t \in [0, T]} \mathbb{E} (Y_t^i - \widehat{Y}_t^i)^2 \leq \frac{\hat{C}}{M_n}, \quad (48)$$

for some positive constants C and \hat{C} . \square

The following proposition strengthens Proposition 3.2.

Proposition 3.4. *We have*

$$\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \mathbb{E} \left[\left(X_t^i - \widehat{X}_t^i \right)^4 \right] = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \sup_{t \in [0, T]} \mathbb{E} \left[\left(Y_t^i - \widehat{Y}_t^i \right)^4 \right] = 0. \quad (49)$$

Proof. Let us define

$$\zeta(t) := \mathbb{E}[(X_t^i - \widehat{X}_t^i)^4] \quad \text{and} \quad \widehat{\zeta}(t) := \mathbb{E}[(Y_t^i - \widehat{Y}_t^i)^4]. \quad (50)$$

The strategy of the proof will be similar to that of Proposition 3.2 that consists of three steps: (1) using Itô's lemma to obtain an expression for $\zeta(t)$, (2) estimating each term that appears in the expression to derive a Grönwall type inequality for $\zeta(t)$ and (3) applying Lemma 3.1 to deduce the assertion.

We now carry out this procedure and will refer to the proof of Proposition 3.2 when similar arguments apply. We first use Itô formula to obtain

$$\begin{aligned} \zeta(t) &= \mathbb{E}[(X_t^i - \widehat{X}_t^i)^4] \\ &= -4\mathbb{E} \int_0^t (\nabla V_1(X_s^i) - \nabla V_1(\widehat{X}_s^i)) \cdot (X_s^i - \widehat{X}_s^i)^3 ds \\ &\quad - 4\mathbb{E} \int_0^t \left[\frac{1}{N_n + M_n} \sum_{j=1}^{N_n} \nabla F_{11}(X_s^i - X_s^j) - a(\nabla F_{11} * \mu_s)(\widehat{X}_s^i) \right] \cdot (X_s^i - \widehat{X}_s^i)^3 ds \\ &\quad - 4\mathbb{E} \int_0^t \left[\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \nabla F_{12}(X_s^i - Y_s^k) - (1-a)(\nabla F_{12} * \mu_s)(\widehat{X}_s^i) \right] \cdot (X_s^i - \widehat{X}_s^i)^3 ds \\ &=: 4 \int_0^t (F_i(s) + G_i(s) + H_i(s) + I_i(s)) ds, \end{aligned} \quad (52)$$

where

$$\begin{aligned} F_i(t) &:= -(\nabla V_1(X_t^i) - \nabla V_1(\widehat{X}_t^i)) \cdot (X_t^i - \widehat{X}_t^i)^3, \\ G_i(t) &:= \left[-\frac{1}{N_n + M_n} \sum_{j=1}^{N_n} (\nabla F_{11}(X_t^i - X_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i)) \right] \cdot (X_t^i - \widehat{X}_t^i)^3, \\ H_i(t) &:= \left(a - \frac{N_n}{N_n + M_n} \right) (\nabla F_{11} * \mu_t)(\widehat{X}_t^i) \cdot (X_t^i - \widehat{X}_t^i)^3, \\ I_i(t) &:= \left[-\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} (\nabla F_{12}(X_t^i - Y_t^k) - (\nabla F_{12} * \nu_t)(\widehat{X}_t^i)) \right] \cdot (X_t^i - \widehat{X}_t^i)^3, \\ J_i(t) &:= \left((1-a) - \frac{M_n}{N_n + M_n} \right) (\nabla F_{12} * \nu_t)(\widehat{X}_t^i) \cdot (X_t^i - \widehat{X}_t^i)^3. \end{aligned}$$

Next we estimate each term F_i, G_i, H_i and I_i . According to Assumption 2.1, we have

$$\mathbb{E}[F_i(s)] \leq \theta_1 \mathbb{E}[(X_s^i - \widehat{X}_s^i)^4]. \quad (53)$$

We write $G_i(t) = -\frac{1}{N_n + M_n}(G_i^1(t) + G_i^2(t))$, where

$$\begin{aligned} G_i^1(t) &:= \sum_{j=1}^{N_n} (\nabla F_{11}(X_t^i - X_t^j) - \nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j)) \cdot (X_t^i - \widehat{X}_t^i)^3, \\ G_i^2(t) &:= \sum_{j=1}^{N_n} (\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i)) \cdot (X_t^i - \widehat{X}_t^i)^3. \end{aligned}$$

Similarly as in the proof of Proposition 3.2, we have

$$\sum_{i=1}^{N_n} G_i^1(t) \geq 0.$$

Using Hölder's inequality

$$\left| \int f g d\gamma \right| \leq \left(\int |f|^{4/3} d\gamma \right)^{3/4} \left(\int |g|^4 d\gamma \right)^{1/4}, \quad (54)$$

we get

$$\begin{aligned} \mathbb{E} [G_i^2(t)] &= \mathbb{E} \left[\sum_{j=1}^{N_n} \left(\nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i) \right) \cdot (X_t^i - \widehat{X}_t^i)^3 \right] \\ &\leq \left(\mathbb{E} [(X_t^i - \widehat{X}_t^i)^4] \right)^{3/4} (\widehat{\kappa}_i(t))^{1/4}, \end{aligned}$$

where

$$\widehat{\kappa}_i(t) := \mathbb{E} \left[\left(\sum_{j=1}^{N_n} \nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i) \right)^4 \right] =: \mathbb{E} \left[\left(\sum_{j=1}^{N_n} a_j(t) \right)^4 \right]$$

where $a_j(t) := \nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j) - (\nabla F_{11} * \mu_t)(\widehat{X}_t^i)$. Using the multinomial theorem

$$\begin{aligned} \left(\sum_{j=1}^n a_j \right)^4 &= \sum_{j=1}^n a_j^4 + \sum_{1 \leq j \neq k \leq n} 4a_j^3 a_k + \sum_{1 \leq j < k \leq n} 6a_j^2 a_k^2 \\ &\quad + \sum_{1 \leq j \neq k \neq \ell \leq n} 12a_j^2 a_k a_\ell + \sum_{1 \leq j \neq k \neq \ell \neq m \leq n} 24a_j a_k a_\ell a_m, \end{aligned}$$

we decompose $\widehat{\kappa}_i(t)$ as follows

$$\widehat{\kappa}_i(t) = \widehat{\kappa}_i^{(1)}(t) + \widehat{\kappa}_i^{(2)}(t) + \widehat{\kappa}_i^{(3)}(t) + \widehat{\kappa}_i^{(4)}(t) + \widehat{\kappa}_i^{(5)}(t),$$

where

$$\begin{aligned} \widehat{\kappa}_i^{(1)}(t) &= \mathbb{E} \left(\sum_{j=1}^n a_j(t)^4 \right), \quad \widehat{\kappa}_i^{(2)}(t) = \mathbb{E} \left(\sum_{1 \leq j \neq k \leq n} 4a_j(t)^3 a_k(t) \right), \quad \widehat{\kappa}_i^{(3)}(t) = \mathbb{E} \left(\sum_{1 \leq j < k \leq n} 6a_j(t)^2 a_k(t)^2 \right), \\ \widehat{\kappa}_i^{(4)}(t) &= \mathbb{E} \left(\sum_{1 \leq j \neq k \neq \ell \leq n} 12a_j(t)^2 a_k(t) a_\ell(t) \right), \quad \widehat{\kappa}_i^{(5)}(t) = \mathbb{E} \left(\sum_{1 \leq j \neq k \neq \ell \neq m \leq n} 24a_j(t) a_k(t) a_\ell(t) a_m(t) \right). \end{aligned}$$

As in the proof of Proposition 3.2

$$\widehat{\kappa}_i^{(2)}(t) = \widehat{\kappa}_i^{(4)}(t) = \widehat{\kappa}_i^{(5)}(t) = 0 \quad \text{and} \quad \widehat{\kappa}_i^{(1)}(t) \leq C N_n^2, \quad \widehat{\kappa}_i^{(3)}(t) \leq C N_n^2.$$

Therefore, we obtain $\widehat{\kappa}_i(t) \leq C N_n^2$, thus

$$\mathbb{E} [G_i^2(t)] \leq C \sqrt{N_n} \left(\mathbb{E} [(X_t^i - \widehat{X}_t^i)^4] \right)^{3/4} \quad \text{and} \quad \mathbb{E} [G_i(t)] \leq \frac{C \sqrt{N_n}}{N_n + M_n} \left(\mathbb{E} [(X_t^i - \widehat{X}_t^i)^4] \right)^{3/4}. \quad (55)$$

Next we estimate H_i . Using Hölder's inequality (54) again we have

$$\mathbb{E} [H_i(t)] \leq \left| a - \frac{N_n}{N_n + M_n} \right| \left(\mathbb{E} [(X_t^i - \widehat{X}_t^i)^4] \right)^{3/4} \left(\mathbb{E} [(\nabla F_{11} * \mu_t)(\widehat{X}_t^i)^4] \right)^{1/4}.$$

As in the proof of (34), it holds that

$$\mathbb{E} [(\nabla F_{11} * \mu_t)(\widehat{X}_t^i)^4] \leq C,$$

which implies that

$$\mathbb{E} (H_i) \leq C \left| a - \frac{N_n}{N_n + M_n} \right| \left(\mathbb{E} [(X_t^i - \widehat{X}_t^i)^4] \right)^{3/4}. \quad (56)$$

We proceed with the term I_i

$$\begin{aligned}
I_i(t) &= \left[-\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \left(\nabla F_{12}(X_t^i - Y_t^k) - (\nabla F_{12} * \nu_t)(\widehat{X}_t^i) \right) \right] \cdot (X_t^i - \widehat{X}_t^i)^3 \\
&= \left[-\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \left(\nabla F_{12}(X_t^i - Y_t^k) - \nabla F_{12}(\widehat{X}_t^i - Y_t^k) \right) \right] \cdot (X_t^i - \widehat{X}_t^i)^3 \\
&\quad - \left[\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \left(\nabla F_{12}(\widehat{X}_t^i - Y_t^k) - (\nabla F_{12} * \nu_t)(\widehat{X}_t^i) \right) \right] \cdot (X_t^i - \widehat{X}_t^i)^3 \\
&=: I_i^{(1)}(t) + I_i^{(2)}(t).
\end{aligned} \tag{57}$$

Using the Lipschitz property of ∇F_{12} we get

$$|I_i^{(1)}(t)| \leq \frac{K}{N_n + M_n} \sum_{k=1}^{M_n} (X_t^i - \widehat{X}_t^i)^4,$$

which implies that

$$\mathbb{E} \left[I_i^{(1)}(t) \right] \leq \frac{K M_n}{N_n + M_n} \mathbb{E} \left[(X_t^i - \widehat{X}_t^i)^4 \right].$$

Summing this estimate over i yields

$$\sum_{i=1}^{N_n} \mathbb{E} \left[I_i^{(1)}(t) \right] \leq \frac{K M_n N_n}{N_n + M_n} \mathbb{E} \left[(X_t^i - \widehat{X}_t^i)^4 \right]. \tag{58}$$

For the term $I_i^{(2)}(t)$, we have

$$\begin{aligned}
\mathbb{E} \left[I_i^{(2)}(t) \right] &= -\frac{1}{N_n + M_n} \mathbb{E} \left[\sum_{k=1}^{M_n} \left(\nabla F_{12}(\widehat{X}_t^i - Y_t^k) - (\nabla F_{12} * \nu_t)(\widehat{X}_t^i) \right) \cdot (X_t^i - \widehat{X}_t^i)^3 \right] \\
&= -\frac{1}{N_n + M_n} \mathbb{E} \left[\sum_{k=1}^{M_n} \left(\nabla F_{12}(\widehat{X}_t^i - Y_t^k) - \nabla F_{12}(\widehat{X}_t^i - \widehat{Y}_t^k) \right) \cdot (X_t^i - \widehat{X}_t^i)^3 \right] \\
&\leq \frac{K}{M_n + N_n} \sum_{k=1}^{M_n} \mathbb{E} \left(|Y_t^k - \widehat{Y}_t^k| |X_t^{1,i} - \widehat{X}_t^i|^3 \right) \\
&\leq \frac{K}{M_n + N_n} \sum_{k=1}^{M_n} \left(\mathbb{E}(Y_t^k - \widehat{Y}_t^k)^4 \right)^{1/4} \left(\mathbb{E}(X_t^i - \widehat{X}_t^i)^4 \right)^{3/4}.
\end{aligned}$$

Hence

$$\begin{aligned}
\sum_{i=1}^{N_n} \mathbb{E} \left[I_i^{(2)}(t) \right] &\leq \frac{K}{M_n + N_n} \left(\sum_{k=1}^{M_n} \left(\mathbb{E}(Y_t^k - \widehat{Y}_t^k)^4 \right)^{1/4} \right) \left(\sum_{i=1}^{N_n} \left(\mathbb{E}(X_t^i - \widehat{X}_t^i)^4 \right)^{3/4} \right) \\
&= \frac{K M_n N_n}{M_n + N_n} \left(\mathbb{E}(Y_t^k - \widehat{Y}_t^k)^4 \right)^{1/4} \left(\mathbb{E}(X_t^i - \widehat{X}_t^i)^4 \right)^{3/4}.
\end{aligned} \tag{59}$$

Substituting (58) and (59) into (57), we obtain

$$\begin{aligned}
\sum_{i=1}^{N_n} \mathbb{E} [I_i(t)] &\leq \frac{K M_n N_n}{N_n + M_n} \mathbb{E} \left[(X_t^i - \widehat{X}_t^i)^4 \right] \\
&\quad + \frac{K M_n N_n}{M_n + N_n} \left(\mathbb{E}(Y_t^k - \widehat{Y}_t^k)^4 \right)^{1/4} \left(\mathbb{E}(X_t^i - \widehat{X}_t^i)^4 \right)^{3/4}.
\end{aligned} \tag{60}$$

Similarly as H_i we get

$$\mathbb{E}[J_i(t)] \leq C \left| (1-a) - \frac{M_n}{N_n + M_n} \right| \left(\mathbb{E}[(X_t^i - \widehat{X}_t^i)^4] \right)^{3/4}. \quad (61)$$

Taking the sum over i in (52) and from estimates (53), (55), (56) and (60) we get

$$\begin{aligned} N_n \zeta(t) \leq & 4 \int_0^t \left[N_n \theta_1 \zeta(s) + \frac{C N_n^{3/2}}{N_n + M_n} \zeta^{3/4}(s) + N_n \left| a - \frac{N_n}{N_n + M_n} \right| \zeta^{3/4}(s) \right. \\ & \left. + C N_n \left| (1-a) - \frac{M_n}{N_n + M_n} \right| \zeta^{3/4}(s) + \frac{K M_n N_n}{N_n + M_n} \zeta(s) + \frac{K M_n N_n}{M_n + N_n} \zeta^{3/4}(s) \widehat{\zeta}^{1/4}(s) \right] ds. \end{aligned}$$

By dividing both sides of the above estimate by N_n , noting that $\frac{M_n}{M_n + N_n} \leq 1$, we write the result in a compact form

$$\zeta(t) \leq C \int_0^t \left(\zeta(s) + g(n) \zeta^{3/4}(s) + \zeta^{3/4} \widehat{\zeta}^{1/4}(s) \right) ds \quad (62)$$

where

$$g(n) = \frac{\sqrt{N_n}}{N_n + M_n} + \left| a - \frac{N_n}{N_n + M_n} \right| + \left| (1-a) - \frac{M_n}{N_n + M_n} \right| \xrightarrow{n \rightarrow \infty} 0$$

Using the inequality of arithmetic and geometric means,

$$3a + b = a + a + a + b \geq 4a^{3/4}b^{1/4} \quad \text{for all } a, b \geq 0,$$

and the non-negativity of ζ and $\widehat{\zeta}$, we obtain the following estimate

$$\zeta(t) \leq C \int_0^t \left[\zeta(s) + \widehat{\zeta}(s) + g(n)(\zeta(s) + \widehat{\zeta}(s))^{3/4} \right] ds. \quad (63)$$

Analogously we obtain a similar estimate for $\widehat{\zeta}$

$$\widehat{\zeta}(t) \leq C \int_0^t \left[\zeta(s) + \widehat{\zeta}(s) + \widehat{g}(n)(\zeta(s) + \widehat{\zeta}(s))^{3/4} \right] ds, \quad (64)$$

where

$$\widehat{g}(n) = \frac{\sqrt{M_n}}{N_n + M_n} + \left| a - \frac{N_n}{N_n + M_n} \right| + \left| (1-a) - \frac{M_n}{N_n + M_n} \right| \xrightarrow{n \rightarrow \infty} 0.$$

Adding (63) and (64) gives

$$(\zeta + \widehat{\zeta})(t) \leq C \int_0^t \left[\zeta(s) + \widehat{\zeta}(s) + (g(n) + \widehat{g}(n))(\zeta(s) + \widehat{\zeta}(s))^{3/4} \right] ds. \quad (65)$$

Applying Lemma 3.1 for $\alpha = \frac{3}{4}$ we get

$$(\zeta + \widehat{\zeta})(t) \leq C \left((g(n) + \widehat{g}(n))(e^{Ct} - 1) \right)^4, \quad (66)$$

from which we deduce the statement of the proposition. \square

Remark 3.5. If N_n and M_n tend to $+\infty$ simultaneously but $\frac{N_n}{M_n}$ is a constant, then from the computations in the proof of Proposition 3.2, we obtain explicit estimates

$$\sup_{t \in [0, T]} \mathbb{E} \left[(X_t^i - \widehat{X}_t^i)^4 \right] \leq \frac{C}{N_n^2} \quad \text{and} \quad \sup_{t \in [0, T]} \mathbb{E} \left[(Y_t^i - \widehat{Y}_t^i)^4 \right] \leq \frac{\hat{C}}{M_n^2}, \quad (67)$$

for some positive constants C and \hat{C} . \square

We are now ready to prove Theorem 1.2.

Proof of Theorem 1.2. According to (27) we have

$$\begin{aligned}
& (X_t^i - \widehat{X}_t^i)^2 \\
&= -2 \int_0^t (\nabla V_1(X_s^i) - \nabla V_1(\widehat{X}_s^i)) \cdot (X_s^i - \widehat{X}_s^i) ds \\
&\quad - 2 \int_0^t \left[\frac{1}{N_n + M_n} \sum_{j=1}^{N_n} \nabla F_{11}(X_s^i - X_s^j) - a(\nabla F_{11} * \mu_s)(\widehat{X}_s^i) \right] \cdot (X_s^i - \widehat{X}_s^i) ds \\
&\quad - 2 \int_0^t \left[\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \nabla F_{12}(X_s^i - Y_s^k) - (1-a)(\nabla F_{12} * \mu_s)(\widehat{X}_s^i) \right] \cdot (X_s^i - \widehat{X}_s^i) ds. \tag{68}
\end{aligned}$$

We define

$$\begin{aligned}
L &:= 2 \int_0^T \left| \left[\frac{1}{N_n + M_n} \sum_{j=1}^{N_n} \nabla F_{11}(X_s^i - X_s^j) - a(\nabla F_{11} * \mu_s)(\widehat{X}_s^i) \right] \cdot (X_s^i - \widehat{X}_s^i) \right| ds \\
&\quad + 2 \int_0^T \left| \left[\frac{1}{N_n + M_n} \sum_{k=1}^{M_n} \nabla F_{12}(X_s^i - Y_s^k) - (1-a)(\nabla F_{12} * \mu_s)(\widehat{X}_s^i) \right] \cdot (X_s^i - \widehat{X}_s^i) \right| ds \tag{69}
\end{aligned}$$

$$=: L_1 + L_2. \tag{70}$$

From (68) and (9) we can estimate

$$(X_t^i - \widehat{X}_t^i)^2 \leq 2\theta_1 \int_0^t |X_s^i - \widehat{X}_s^i|^2 ds + L. \tag{71}$$

Setting $\varphi(t) := \int_0^t |X_s^i - \widehat{X}_s^i|^2 ds$, we get

$$\varphi'(t) \leq 2\theta_1 \varphi(t) + L.$$

Using Grönwall Lemma and the fact that $\varphi(0) = 0$, we deduce that, for any $t \in [0, T]$,

$$\varphi(t) \leq \frac{L}{2\theta_1} (e^{2\theta_1 t} - 1).$$

Substituting this back into (71), we obtain

$$(X_t^i - \widehat{X}_t^i)^2 = \varphi'(t) \leq M e^{2\theta_1 t} \leq e^{2\theta_1 T} L,$$

from which we deduce that

$$\mathbb{E} \left[\sup_{t \in [0, T]} (X_t^i - \widehat{X}_t^i)^2 \right] \leq e^{2\theta_1 T} \mathbb{E}[L]. \tag{72}$$

Next we find an upper bound for L . For the first term, L_1 , proceeding similarly as the terms B_i and C_i in the proof of Proposition 3.2, we have

$$\begin{aligned}
\mathbb{E}[L_1] &\leq \frac{2}{M_n + N_n} \sum_{j=1}^{N_n} \int_0^T (|\rho_{ij}^1(s)| + |\rho_{ij}^2(s)|) ds + 2C \left| a - \frac{N_n}{M_n + N_n} \right| \int_0^T \left(E[X_s^i - \widehat{X}_s^i]^2 \right)^{\frac{1}{2}} ds \\
&\leq \frac{2}{M_n + N_n} \sum_{j=1}^{N_n} \int_0^T (|\rho_{ij}^1(s)| + |\rho_{ij}^2(s)|) ds + 2CT \left| a - \frac{N_n}{M_n + N_n} \right| \left(\sup_{s \in [0, T]} E[X_s^i - \widehat{X}_s^i]^2 \right)^{\frac{1}{2}} \\
&=: K_1 + K_2, \tag{73}
\end{aligned}$$

where $\rho_{ij}^1(s)$ and $\rho_{ij}^2(s)$ are defined in (30).

By Proposition 3.2, K_2 tends to 0 as n goes to $+\infty$. If $\frac{M_n}{N_n}$ is constant, then $K_2 = 0$. We now estimate both terms in K_1 . According to (35), we have

$$E\left(\sum_{j=1}^{N_n} |\rho_{ij}^2(s)|\right) \leq C\sqrt{N_n} \left(\sup_{s \in [0, T]} E[X_s^i - \widehat{X}_s^i]^2\right)^{\frac{1}{2}}.$$

Thus,

$$\frac{2}{M_n + N_n} \int_0^T E\left(\sum_{j=1}^{N_n} |\rho_{ij}^2(s)|\right) ds \leq \frac{2CT\sqrt{N_n}}{M_n + N_n} \left(\sup_{s \in [0, T]} E[X_s^i - \widehat{X}_s^i]^2\right)^{\frac{1}{2}},$$

which converges to 0 as $n \rightarrow +\infty$. If $\frac{M_n}{N_n}$ is constant, then

$$\sup_{s \in [0, T]} E\left[(X_s^i - \widehat{X}_s^i)^2\right] \leq \frac{C}{N_n},$$

which implies that

$$\frac{2}{M_n + N_n} \int_0^T E\left(\sum_{j=1}^{N_n} |\rho_{ij}^2(s)|\right) ds \leq \frac{2CT}{N_n}. \quad (74)$$

For the ρ_{ij}^1 term, using (30), Cauchy-Schwarz inequality and Proposition 2.14, we get

$$\begin{aligned} \mathbb{E}[|\rho_{ij}^1(s)|] &\leq \left\{ \mathbb{E}[(X_s^i - \widehat{X}_s^i)^2] \mathbb{E}[(\nabla F_{11}(X_t^i - X_t^j) - \nabla F_{11}(\widehat{X}_t^i - \widehat{X}_t^j))^2] \right\}^{\frac{1}{2}} \\ &\leq \left\{ \mathbb{E}[(X_s^i - \widehat{X}_s^i)^2] \right\}^{\frac{1}{2}} \\ &\quad \times \left\{ \mathbb{E}[(X_s^i - \widehat{X}_s^i + \widehat{X}_s^j - X_s^j)^2 (c + |X_s^i - X_s^j|^{2q} + |\widehat{X}_s^i - \widehat{X}_s^j|^{2q})^2] \right\}^{\frac{1}{2}} \\ &\leq \left\{ \sup_{s \in [0, T]} \mathbb{E}[(X_s^i - \widehat{X}_s^i)^2] \right\}^{\frac{1}{2}} \\ &\quad \times \left\{ \mathbb{E}[(X_s^i - \widehat{X}_s^i + \widehat{X}_s^j - X_s^j)^4] \mathbb{E}[(c + |X_s^i - X_s^j|^{2q} + |\widehat{X}_s^i - \widehat{X}_s^j|^{2q})^4] \right\}^{\frac{1}{4}} \\ &\leq C \left\{ \sup_{s \in [0, T]} \mathbb{E}[(X_s^i - \widehat{X}_s^i)^2] \right\}^{\frac{1}{2}} \times \left\{ \sup_{s \in [0, T]} \mathbb{E}[(X_s^i - \widehat{X}_s^i)^4] \right\}^{\frac{1}{4}}, \end{aligned}$$

which tends to 0 as n tends to $+\infty$ according to Propositions 3.2 and 3.4. If $\frac{N_n}{M_n}$ is constant then,

$$\mathbb{E}[|\rho_{ij}^1(s)|] \leq C \left\{ \sup_{s \in [0, T]} \mathbb{E}[(X_s^i - \widehat{X}_s^i)^2] \right\}^{\frac{1}{2}} \times \left\{ \sup_{s \in [0, T]} \mathbb{E}[(X_s^i - \widehat{X}_s^i)^4] \right\}^{\frac{1}{4}} \leq \frac{C}{N_n}.$$

Thus

$$\frac{2}{M_n + N_n} \sum_{j=1}^{N_n} \int_0^T \mathbb{E}[|\rho_{ij}^1(s)|] ds \leq \frac{C}{N_n}. \quad (75)$$

From (73), (75) and (74), we get

$$\mathbb{E}[L_1] \leq \frac{C}{N_n}. \quad (76)$$

The second term L_2 can be estimated similarly as the terms D_i and E_i in the proof of Proposition 3.2. We have

$$\mathbb{E}L_2 \leq 2 \int_0^T \left\{ \frac{CM_n}{M_n + N_n} \mathbb{E}(X_s^i - \widehat{X}_s^i)^2 + \frac{C}{M_n + N_n} \sum_{k=1}^{M_n} \left(\mathbb{E}[(Y_s^k - \widehat{Y}_s^k)^2] \mathbb{E}[(X_s^i - \widehat{X}_s^i)^2] \right) \right\}^{\frac{1}{2}}$$

$$\begin{aligned}
& + C \left| (1-a) - \frac{M_n}{M_n + N_n} \right| \left[\mathbb{E}(X_s^i - \widehat{X}_s^i)^2 \right]^{\frac{1}{2}} \Big\} ds \\
& \leq 2 \int_0^T \left\{ \frac{CM_n}{M_n + N_n} \sup_{s \in [0, T]} \mathbb{E}(X_s^i - \widehat{X}_s^i)^2 + C \left| (1-a) - \frac{M_n}{M_n + N_n} \right| \left[\sup_{s \in [0, T]} \mathbb{E}(X_s^i - \widehat{X}_s^i)^2 \right]^{\frac{1}{2}} \right. \\
& \quad \left. + \frac{CM_n}{M_n + N_n} \left(\sup_{s \in [0, T]} \mathbb{E}[(Y_s^i - \widehat{Y}_s^i)^2] \sup_{s \in [0, T]} \mathbb{E}[(X_s^i - \widehat{X}_s^i)^2] \right)^{\frac{1}{2}} \right\} ds \\
& \leq 2T \left\{ \frac{CM_n}{M_n + N_n} \sup_{s \in [0, T]} \mathbb{E}(X_s^i - \widehat{X}_s^i)^2 + C \left| (1-a) - \frac{M_n}{M_n + N_n} \right| \left[\sup_{s \in [0, T]} \mathbb{E}(X_s^i - \widehat{X}_s^i)^2 \right]^{\frac{1}{2}} \right. \\
& \quad \left. + \frac{CM_n}{M_n + N_n} \left(\sup_{s \in [0, T]} \mathbb{E}[(Y_s^i - \widehat{Y}_s^i)^2] \sup_{s \in [0, T]} \mathbb{E}[(X_s^i - \widehat{X}_s^i)^2] \right)^{\frac{1}{2}} \right\}, \tag{77}
\end{aligned}$$

According to Proposition 3.2 the RHS of (77) tends to 0 as $n \rightarrow \infty$. If $\frac{N_n}{M_n}$ is constant then it is bounded by $\frac{C}{N_n}$. Substituting estimates of $\mathbb{E}[L_1]$ and $\mathbb{E}[L_2]$ back into (72), we obtain

$$\mathbb{E} \left[\sup_{t \in [0, T]} (X_t^i - \widehat{X}_t^i)^2 \right] \leq \frac{C e^{2\theta_1 T}}{N_n},$$

from which the first assertion of the theorem follows. The second assertion is obtained analogously. \square

4 Existence and non-uniqueness of invariant measures

In this section we prove Theorem 1.3 showing the existence and non-uniqueness of invariant measures of the PDE system (2) that is given below.

$$\partial_t \mu_t = \operatorname{div} \left((\nabla V_1 + a(\nabla F_{11} * \mu_t) + (1-a)(\nabla F_{12} * \nu_t)) \mu_t \right) + \frac{\sigma^2}{2} \Delta \mu_t, \tag{78a}$$

$$\partial_t \nu_t = \operatorname{div} \left((\nabla V_2 + a(\nabla F_{21} * \mu_t) + (1-a)(\nabla F_{22} * \nu_t)) \nu_t \right) + \frac{\sigma^2}{2} \Delta \nu_t. \tag{78b}$$

We only consider the quadratic interaction potentials

$$F_{ij}(x) = \frac{\alpha_{ij} x^2}{2}.$$

We expect that extensions to polynomial potentials could be possible but the analysis will be much more intricate. We leave this for future investigation. Stationary solutions $(\mu(x) dx, \nu(x) dx)$ of the above system is determined by

$$\mu(x) = \frac{\exp \left(-\frac{2}{\sigma^2} (V_1(x) + aF_{11} * \mu(x) + (1-a)F_{12} * \nu(x)) \right)}{\int \exp \left(-\frac{2}{\sigma^2} (V_1(x) + aF_{11} * \mu(x) + (1-a)F_{12} * \nu(x)) \right) dx}, \tag{79a}$$

$$\nu(x) = \frac{\exp \left(-\frac{2}{\sigma^2} (V_2(x) + aF_{21} * \mu(x) + (1-a)F_{22} * \nu(x)) \right)}{\int \exp \left(-\frac{2}{\sigma^2} (V_2(x) + aF_{21} * \mu(x) + (1-a)F_{22} * \nu(x)) \right) dx}. \tag{79b}$$

We define

$$m_1 := \int x \mu(x) dx \quad \text{and} \quad m_2 = \int x \nu(x) dx.$$

Using explicit formulas $F_{ij}(x) = \frac{\alpha_{ij}}{2} x^2$ for $i, j = 1, 2$, we obtain

$$\mu(x) = \frac{\exp \left(-\frac{2}{\sigma^2} (V_1(x) + a\frac{\alpha_{11}}{2} x^2 - a\alpha_{11} m_1 x + (1-a)\frac{\alpha_{12}}{2} x^2 - (1-a)\alpha_{12} m_2 x) \right)}{\int \exp \left(-\frac{2}{\sigma^2} (V_1(x) + a\frac{\alpha_{11}}{2} x^2 - a\alpha_{11} m_1 x + (1-a)\frac{\alpha_{12}}{2} x^2 - (1-a)\alpha_{12} m_2 x) \right) dx}, \tag{80a}$$

$$\nu(x) = \frac{\exp\left(-\frac{2}{\sigma^2}\left(V_2(x) + a\frac{\alpha_{21}}{2}x^2 - a\alpha_{21}m_1x + (1-a)\frac{\alpha_{22}}{2}x^2 - (1-a)\alpha_{22}m_2x\right)\right)}{\int \exp\left(-\frac{2}{\sigma^2}\left(V_2(x) + a\frac{\alpha_{21}}{2}x^2 - a\alpha_{21}m_1x + (1-a)\frac{\alpha_{22}}{2}x^2 - (1-a)\alpha_{22}m_2x\right)\right) dx}. \quad (80b)$$

Therefore, (m_1, m_2) satisfies the following system

$$m_1 = \frac{\int x \exp\left(-\frac{2}{\sigma^2}\left(V_1(x) + a\frac{\alpha_{11}}{2}x^2 - a\alpha_{11}m_1x + (1-a)\frac{\alpha_{12}}{2}x^2 - (1-a)\alpha_{12}m_2x\right)\right) dx}{\int \exp\left(-\frac{2}{\sigma^2}\left(V_1(x) + a\frac{\alpha_{11}}{2}x^2 - a\alpha_{11}m_1x + (1-a)\frac{\alpha_{12}}{2}x^2 - (1-a)\alpha_{12}m_2x\right)\right) dx}, \quad (81a)$$

$$m_2 = \frac{\int x \exp\left(-\frac{2}{\sigma^2}\left(V_2(x) + a\frac{\alpha_{21}}{2}x^2 - a\alpha_{21}m_1x + (1-a)\frac{\alpha_{22}}{2}x^2 - (1-a)\alpha_{22}m_2x\right)\right) dx}{\int \exp\left(-\frac{2}{\sigma^2}\left(V_2(x) + a\frac{\alpha_{21}}{2}x^2 - a\alpha_{21}m_1x + (1-a)\frac{\alpha_{22}}{2}x^2 - (1-a)\alpha_{22}m_2x\right)\right) dx}. \quad (81b)$$

We define $\Phi_1(m_1, m_2)$ and $\Phi_2(m_1, m_2)$ to be the right-hand sides of (81a) and (81b), respectively. Setting $\Phi(m_1, m_2) := (\Phi_1, \Phi_2)(m_1, m_2)$. We rewrite (81) as

$$(m_1, m_2) = \Phi(m_1, m_2), \quad (82)$$

where $\Phi(m_1, m_2)$ denotes its right-hand side.

4.1 Symmetrical invariant measure

We suppose that V_1 and V_2 are symmetrical.

Lemma 4.1. *There exists a unique pair of symmetric invariant measures that are given by*

$$\mu^0(x) = \frac{\exp\left(-\frac{2}{\sigma^2}\left(V_1(x) + a\frac{\alpha_{11}}{2}x^2 + (1-a)\frac{\alpha_{12}}{2}x^2\right)\right)}{\int \exp\left(-\frac{2}{\sigma^2}\left(V_1(x) + a\frac{\alpha_{11}}{2}x^2 + (1-a)\frac{\alpha_{12}}{2}x^2\right)\right) dx}, \quad (83a)$$

$$\nu^0(x) = \frac{\exp\left(-\frac{2}{\sigma^2}\left(V_2(x) + a\frac{\alpha_{21}}{2}x^2 + (1-a)\frac{\alpha_{22}}{2}x^2\right)\right)}{\int \exp\left(-\frac{2}{\sigma^2}\left(V_2(x) + a\frac{\alpha_{21}}{2}x^2 + (1-a)\frac{\alpha_{22}}{2}x^2\right)\right) dx}. \quad (83b)$$

Proof. Since (μ^0, ν^0) satisfy (79a) with their mean values $(0, 0)$ that fulfill (81b), they are invariant measures and are symmetric because V_1 and V_2 are symmetric. Now suppose that (μ^0, ν^0) is an arbitrary symmetric invariant measure. Then $(\hat{\mu}^0, \hat{\nu}^0)$ satisfies (79a) with (m_1, m_2) replaced by their mean values $(m_{\hat{\mu}^0}, m_{\hat{\nu}^0})$. Since $(\hat{\mu}^0$ and $\hat{\nu}^0)$ are symmetric, we have

$$m_{\hat{\mu}^0} = \int x \hat{\mu}^0(x) dx = 0 = \int x \hat{\nu}^0(x) dx = m_{\hat{\nu}^0}.$$

Substituting these values back into (79a) we obtain $(\hat{\mu}^0, \hat{\nu}^0) = (\mu^0, \nu^0)$. \square

4.2 Other invariant measures

We are now interested in non-symmetrical invariant measures.

Assumption 4.1. Suppose that V_1 and V_2 have a common unique minimizer m^*

$$V_1'(m^*) = V_2'(m^*) = 0, \quad V_1''(m^*) > 0 \quad \text{and} \quad V_2''(m^*) > 0.$$

We will make use of the following result.

Lemma 4.2. [HT10, Lemma A.3] Let U and G be two $C^\infty(\mathbb{R})$ -continuous functions. Let λ be a parameter that belongs to some compact interval \mathcal{I} of \mathbb{R} . We define $U_\lambda = U + \lambda G$. Suppose that $U_\lambda(z) \geq z^2$ for $|z|$ larger than some value R independent of λ and that U_λ has a unique global minimum at z_λ with $U''_\lambda(z_\lambda) > 0$. Let f_m be a C^3 -continuous function depending on some parameter m that belongs to a compact set \mathcal{M} . Furthermore, we also assume that there exists some constant $\theta > 0$ such that $|f_m(z)| \leq \exp[\theta|U_\lambda(z)|]$ for all $z \geq R, \lambda \in \mathcal{I}, m \in \mathcal{M}$ and $f_m^{(k)}$ is locally bounded uniformly with respect to the parameter $m \in \mathcal{M}$ for $0 \leq k \leq 3$. Let $a, b \in \mathbb{R}$ such that $a < z_\lambda < b$. Then the following asymptotic result holds as ε tends to 0:

$$\int_a^b f_m(z) \exp\left[-\frac{2U_\lambda(z)}{\varepsilon}\right] dz = \sqrt{\frac{\pi\varepsilon}{\mathcal{U}_2}} \exp\left[-\frac{2U_\lambda(z_\lambda)}{\varepsilon}\right] \left\{ f_m(z_\lambda) + \gamma_0(\lambda)\varepsilon + o_{\mathcal{I}\mathcal{M}}^{(1)}(\varepsilon) \right\}, \quad (84)$$

with

$$\gamma_0(\lambda) = f_m(z_\lambda) \left(\frac{5\mathcal{U}_3^2}{48\mathcal{U}_2^3} - \frac{\mathcal{U}_4}{16\mathcal{U}_2^2} \right) - f'_m(z_\lambda) \frac{\mathcal{U}_3}{4\mathcal{U}_2^2} + \frac{f''_m(z_\lambda)}{4\mathcal{U}_2}. \quad (85)$$

Here $\mathcal{U}_k = U_\lambda^{(k)}(z_\lambda)$ and $o_{\mathcal{I}\mathcal{M}}^{(1)}(\varepsilon)/\varepsilon$ converges to 0 as ε goes to 0 uniformly with respect to the parameters m and λ . Moreover, for any $n \geq 1$, we have

$$\frac{\int_{\mathbb{R}} z^n e^{f_m(z)} e^{-\frac{2U_\lambda(z)}{\varepsilon}} dz}{\int_{\mathbb{R}} e^{f_m(z)} e^{-\frac{2U_\lambda(z)}{\varepsilon}} dz} - z_\lambda^n \approx -\frac{n z_\lambda^{n-2}}{4\mathcal{U}_2} \left[z_\lambda \frac{\mathcal{U}_3}{\mathcal{U}_2} - n + 1 - 2z_\lambda f'_m(z_\lambda) \right] \varepsilon, \quad (86)$$

where the estimate is uniform with respect to the parameters m and λ as $\varepsilon \rightarrow 0$.

We are now ready to prove Theorem 1.3.

Proof of Theorem 1.3. We recall that $\rho > 0$ is defined such that

$$\rho \geq \max \left\{ \frac{|V_1^{(3)}(m^*)|}{4V_1''(m^*)(V_1''(m^*) + a\alpha_{11} + (1-a)\alpha_{12})}, \frac{|V_2^{(3)}(m^*)|}{4V_2''(m^*)(V_2''(m^*) + a\alpha_{21} + (1-a)\alpha_{22})} \right\}. \quad (87)$$

We define

$$D(\sigma) := [m^* - \rho\sigma^2, m^* + \rho\sigma^2] \times [m^* - \rho\sigma^2, m^* + \rho\sigma^2].$$

Let $(m_1, m_2) \in D(\sigma)$. Then there exist ρ_1, ρ_2 with $0 \leq |\rho_1|, |\rho_2| \leq \rho$ such that

$$m_i = m^* + \rho_i \sigma^2, \quad i = 1, 2.$$

We have

$$\begin{aligned} \Phi_1(m_1, m_2) &= \frac{\int x \exp\left(-\frac{2}{\sigma^2}(V_1(x) + a\frac{\alpha_{11}}{2}x^2 - a\alpha_{11}(m^* + \rho_1\sigma^2)x + (1-a)\frac{\alpha_{12}}{2}x^2 - (1-a)\alpha_{12}(m^* + \rho_2\sigma^2)x)\right) dx}{\int \exp\left(-\frac{2}{\sigma^2}(V_1(x) + a\frac{\alpha_{11}}{2}x^2 - a\alpha_{11}(m^* + \rho_1\sigma^2)x + (1-a)\frac{\alpha_{12}}{2}x^2 - (1-a)\alpha_{12}(m^* + \rho_2\sigma^2)x)\right) dx} \\ &= \frac{\int x e^{2a\alpha_{11}\rho_1 x + 2(1-a)\alpha_{12}\rho_2 x} \exp\left(-\frac{2}{\sigma^2}(V_1(x) + a\frac{\alpha_{11}}{2}x^2 - a\alpha_{11}m^*x + (1-a)\frac{\alpha_{12}}{2}x^2 - (1-a)\alpha_{12}m^*x)\right) dx}{\int e^{2a\alpha_{11}\rho_1 x + 2(1-a)\alpha_{12}\rho_2 x} \exp\left(-\frac{2}{\sigma^2}(V_1(x) + a\frac{\alpha_{11}}{2}x^2 - a\alpha_{11}m^*x + (1-a)\frac{\alpha_{12}}{2}x^2 - (1-a)\alpha_{12}m^*x)\right) dx}. \end{aligned}$$

Set $U(x) = V_1(x) + a\frac{\alpha_{11}}{2}x^2 - a\alpha_{11}m^*x + (1-a)\frac{\alpha_{12}}{2}x^2 - (1-a)\alpha_{12}m^*x$. We have

$$\begin{aligned} U'(x) &= V_1'(x) + (a\alpha_{11} + (1-a)\alpha_{12})(x - m^*), \\ U''(x) &= V_1''(x) + a\alpha_{11} + (1-a)\alpha_{12}, \\ U^{(3)}(x) &= V_1^{(3)}(x). \end{aligned}$$

Since $U'(m^*) = V_1'(m^*) = 0, U''(m^*) = V_1''(m^*) > 0$, m^* is the unique minimizer of U . Applying Lemma 4.2 for $f(x) = 2a\alpha_{11}\rho_1x + 2(1-a)\alpha_{12}\rho_2x, n = 1, U(x)$ and $\lambda = 0$, we get

$$\begin{aligned}\Phi_1(m^* + \rho_1\sigma^2, m^* + \rho_2\sigma^2) &= m^* - \frac{1}{4m^*\mathcal{U}_2} \left[m^*\frac{\mathcal{U}_3}{\mathcal{U}_2} - 2m^*f'(m^*) \right] \sigma^2 + o(\sigma^2) \\ &= m^* - \left[\frac{V_1^{(3)}(m^*)}{4(V_1''(m^*) + \tau)^2} + \frac{\zeta}{V_1''(m^*) + \tau} \right] \sigma^2 + o(\sigma^2) \\ &=: m^* - k_1\sigma^2 + o(\sigma^2),\end{aligned}$$

where

$$\tau := a\alpha_{11} + (1-a)\alpha_{12}, \quad \zeta := a\alpha_{11}\rho_1 + (1-a)\alpha_{12}\rho_2.$$

We have

$$\begin{aligned}|k_1| &\leq \frac{|V_1^{(3)}|}{4(V_1''(m^*) + \tau)^2} + \frac{a\alpha_{11}|\rho_1| + (1-a)\alpha_{12}|\rho_2|}{V_1''(m^*) + \tau} \\ &\leq \frac{|V_1^{(3)}|}{4(V_1''(m^*) + \tau)^2} + \frac{a\alpha_{11}\rho + (1-a)\alpha_{12}\rho}{V_1''(m^*) + \tau} \\ &= \frac{|V_1^{(3)}|}{4(V_1''(m^*) + \tau)^2} + \frac{\tau\rho}{V_1''(m^*) + \tau} \\ &\stackrel{(87)}{\leq} \rho.\end{aligned}$$

Similarly we have

$$\Phi_2(m^* + \rho_1\sigma^2, m^* + \rho_2\sigma^2) = m^* - k_2\sigma^2 + o(\sigma^2) \quad \text{where } |k_2| \leq \rho.$$

Thus for σ small enough, we have $\Phi(m^* + \rho_1\sigma^2, m^* + \rho_2\sigma^2) \in D(\sigma)$. By Brouwer's fixed-point theorem, there exist $(m_1, m_2) \in D(\sigma)$ that satisfy (81), thus the measures μ and ν defined in (80) are invariant measures for the coupled MV-equations. \square

Remark 4.3. Assumption 4.1 has been used to obtain that the two functions

$$\begin{aligned}U(x) &= V_1(x) + a\frac{\alpha_{11}}{2}x^2 - a\alpha_{11}m^*x + (1-a)\frac{\alpha_{12}}{2}x^2 - (1-a)\alpha_{12}m^*x, \\ \hat{U}(x) &= V_2(x) + a\frac{\alpha_{21}}{2}x^2 - a\alpha_{21}m^*x + (1-a)\frac{\alpha_{22}}{2}x^2 - (1-a)\alpha_{22}m^*x,\end{aligned}$$

have the common unique minimizer m^* which is also the minimizer of V_1 and V_2 . We expect that this assumption can be removed. To this end, one would need to find a solution (m_1^*, m_2^*) to the following system

$$\begin{aligned}V_1'(m_1) + (1-a)\alpha_{12}(m_1 - m_2) &= 0, \\ V_2'(m_2) + a\alpha_{21}(m_1 - m_2) &= 0.\end{aligned}$$

Then one apply Brouwer's fixed-point theorem for $D(\sigma) = [m_1^* + \rho_1\sigma^2, m_2^* + \rho_2\sigma^2]$ where $0 \leq |\rho_1|, |\rho_2| \leq \rho$ with a suitable choice of ρ . \square

References

- [BGG13] F. Bolley, I. Gentil, and A. Guillin. Uniform convergence to equilibrium for granular media. *Archive for Rational Mechanics and Analysis*, 208(2):429–445, May 2013.
- [BGM10] F. Bolley, A. Guillin, and F. Malrieu. Trend to equilibrium and particle approximation for a weakly selfconsistent Vlasov-Fokker-Planck equation. *M2AN Math. Model. Numer. Anal.*, 44(5):867–884, 2010.

- [BnCD16] A. Barbaro, J. Ca nizo, J. Carrillo, and P. Degond. Phase transitions in a kinetic flocking model of Cucker–Smale type. *Multiscale Modeling & Simulation*, 14(3):1063–1088, 2016.
- [BRTV98] S. Benachour, B. Roynette, D. Talay, and P. Vallois. Nonlinear self-stabilizing processes. I. Existence, invariant probability, propagation of chaos. *Stochastic Process. Appl.*, 75(2):173–201, 1998.
- [BRV98] S. Benachour, B. Roynette, and P. Vallois. Nonlinear self-stabilizing processes. II. Convergence to invariant probability. *Stochastic Process. Appl.*, 75(2):203–224, 1998.
- [BT08] J. Binney and S. Tremaine. *Galactic Dynamics: Second Edition*. Princeton University Press, 2008.
- [CDJ18] X. Chen, Es. S. Daus, and A. Jüngel. Global existence analysis of cross-diffusion population systems for multiple species. *Archive for Rational Mechanics and Analysis*, 227(2):715–747, Feb 2018.
- [CEV11] C. Conca, E. Espejo, and K. Vilches. Remarks on the blowup and global existence for a two species chemotactic Keller–Segel system in \mathbb{R}^2 . *European Journal of Applied Mathematics*, 22(6):553–580, 2011.
- [CGM08] P. Cattiaux, A. Guillin, and F. Malrieu. Probabilistic approach for granular media equations in the non-uniformly convex case. *Probability Theory and Related Fields*, 140(1):19–40, Jan 2008.
- [CGPS18] J. A. Carrillo, R. S. Gvalani, G. A. Pavliotis, and A. Schlichting. Long-time behaviour and phase transitions for the McKean–Vlasov equation on the torus, 2018.
- [CHS18] J. Carrillo, Y. Huang, and M. Schmidtchen. Zoology of a nonlocal cross-diffusion model for two species. *SIAM Journal on Applied Mathematics*, 78(2):1078–1104, 2018.
- [CJ17] X. Chen and A. Jüngel. A note on the uniqueness of weak solutions to a class of cross-diffusion systems. *Journal of Evolution Equations*, Nov 2017.
- [CL16] G. Carlier and M. Laborde. Remarks on continuity equations with nonlinear diffusion and nonlocal drifts. *Journal of Mathematical Analysis and Applications*, 444(2):1690 – 1702, 2016.
- [CLM12] R. M. Colombo and M. Lécureux-Mercier. Nonlocal crowd dynamics models for several populations. *Acta Mathematica Scientia*, 32(1):177 – 196, 2012.
- [CLM13] G. Crippa and M. Lécureux-Mercier. Existence and uniqueness of measure solutions for a system of continuity equations with non-local flow. *Nonlinear Differential Equations and Applications NoDEA*, 20(3):523–537, Jun 2013.
- [CMV03] J. A. Carrillo, R. J. McCann, and C. Villani. Kinetic equilibration rates for granular media and related equations: entropy dissipation and mass transportation estimates. *Rev. Mat. Iberoamericana*, 19(3):971–1018, 12 2003.
- [CMV06] J. A. Carrillo, R. J. McCann, and C. Villani. Contractions in the 2-Wasserstein length space and thermalization of granular media. *Archive for Rational Mechanics and Analysis*, 179(2):217–263, Feb 2006.
- [CP10] L. Chayes and V. Panferov. The McKean-Vlasov equation in finite volume. *J. Stat. Phys.*, 138(1-3):351–380, 2010.
- [Daw83] D. A. Dawson. Critical dynamics and fluctuations for a mean-field model of cooperative behavior. *J. Statist. Phys.*, 31(1):29–85, 1983.

- [DEGZ18] A. Durmus, A. Eberle, A. Guillin, and R. Zimmer. An elementary approach to uniform in time propagation of chaos, 2018.
- [DFF16] M. Di Francesco and S. Fagioli. A nonlocal swarm model for predators–prey interactions. *Mathematical Models and Methods in Applied Sciences*, 26(02):319–355, 2016.
- [DMR17] M. H. Duong, A. Muntean, and O. M. Richardson. Discrete and continuum links to a nonlinear coupled transport problem of interacting populations. *The European Physical Journal Special Topics*, 226(10):2345–2357, Jul 2017.
- [Dra03] S. S. Dragomir. *Some Gronwall type inequalities and applications*. Nova Science Publishers, Inc., Hauppauge, NY, 2003.
- [Duo15] M. H. Duong. Long time behaviour and particle approximation of a generalised Vlasov dynamic. *Nonlinear Anal.*, 127:1–16, 2015.
- [EFK17] J. Evers, R. Fetecau, and T. Kolokolnikov. Equilibria for an aggregation model with two species. *SIAM Journal on Applied Dynamical Systems*, 16(4):2287–2338, 2017.
- [EK16] J. Evers and T. Kolokolnikov. Metastable states for an aggregation model with noise. *SIAM Journal on Applied Dynamical Systems*, 15(4):2213–2226, 2016.
- [ESV10] E.E. Espejo, A. Stevens, and J.J.L. Velázquez. A note on non-simultaneous blow-up for a drift-diffusion model. *Differential Integral Equations*, 23(5/6):451–462, 05 2010.
- [FEF18] M. Di Francesco, A. Esposito, and S. Fagioli. Nonlinear degenerate cross-diffusion systems with nonlocal interaction. *Nonlinear Analysis*, 169:94 – 117, 2018.
- [FF13] M. Di Francesco and S. Fagioli. Measure solutions for non-local interaction PDEs with two species. *Nonlinearity*, 26(10):2777, 2013.
- [Fun84] T. Funaki. A certain class of diffusion processes associated with nonlinear parabolic equations. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 67(3):331–348, Oct 1984.
- [GP18] S. N. Gomes and G. A. Pavliotis. Mean Field Limits for Interacting Diffusions in a Two-Scale Potential. *J. Nonlinear Sci.*, 28(3):905–941, 2018.
- [GPY13] J. Garnier, G. Papanicolaou, and T.-W. Yang. Large deviations for a mean field model of systemic risk. *SIAM J. Financial Math.*, 4(1):151–184, 2013.
- [GPY17] J. Garnier, G. Papanicolaou, and T.-W. Yang. Consensus convergence with stochastic effects. *Vietnam J. Math.*, 45(1-2):51–75, 2017.
- [Her03] S. Herrmann. Système de processus auto-stabilisants. *Dissertationes Math. (Rozprawy Mat.)*, 414:49, 2003.
- [HIP08] S. Herrmann, P. Imkeller, and D. Peithmann. Large deviations and a Kramers’ type law for self-stabilizing diffusions. *Ann. Appl. Probab.*, 18(4):1379–1423, 08 2008.
- [HM14] M. Hauray and S. Mischler. On Kac’s chaos and related problems. *Journal of Functional Analysis*, 266(10):6055 – 6157, 2014.
- [HT10] S. Herrmann and J. Tugaut. Non-uniqueness of stationary measures for self-stabilizing processes. *Stochastic Process. Appl.*, 120(7):1215–1246, 2010.
- [JW16] P.-E. Jabin and Z. Wang. Mean field limit and propagation of chaos for Vlasov systems with bounded forces. *Journal of Functional Analysis*, 271(12):3588 – 3627, 2016.
- [JW17a] P.-E. Jabin and Z. Wang. *Mean Field Limit for Stochastic Particle Systems*, pages 379–402. Springer International Publishing, Cham, 2017.

- [JW17b] P.-E. Jabin and Z. Wang. Quantitative estimate of propagation of chaos for stochastic systems with $w^{-1,\infty}$ kernels, 2017.
- [Kac56] M. Kac. Foundations of kinetic theory. In *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, Volume 3: Contributions to Astronomy and Physics*, pages 171–197, Berkeley, Calif., 1956. University of California Press.
- [KO03] M. Kurokiba and T. Ogawa. Finite time blow-up of the solution for a nonlinear parabolic equation of drift-diffusion type. *Differential Integral Equations*, 16(4):427–452, 2003.
- [KRZ18] N. I. Kavallaris, T. Ricciardi, and G. Zecca. A multi-species chemotaxis system: Lyapunov functionals, duality, critical mass. *European Journal of Applied Mathematics*, 29(3):515–542, 2018.
- [KS71] E. F. Keller and L. A. Segel. Model for chemotaxis. *Journal of Theoretical Biology*, 30(2):225 – 234, 1971.
- [Lab17] M. Laborde. On some nonlinear evolution systems which are perturbations of Wasserstein gradient flows. In M. Bergounioux, É. Oudet, M. Rumpf, G. Carlier, T. Champion, and F. Santambrogio, editors, *Topological Optimization and Optimal Transport: In the Applied Sciences*, chapter 12, pages 304–332. Berlin, Boston: De Gruyter, 2017.
- [LM17] T. Lepoutre and A. Moussa. Entropic structure and duality for multiple species cross-diffusion systems. *Nonlinear Analysis*, 159:298 – 315, 2017. Advances in Reaction-Cross-Diffusion Systems.
- [Mal03] F. Malrieu. Convergence to equilibrium for granular media equations and their Euler schemes. *Ann. Appl. Probab.*, 13(2):540–560, 05 2003.
- [McK66] H. P. McKean, Jr. A class of Markov processes associated with nonlinear parabolic equations. *Proc. Nat. Acad. Sci. U.S.A.*, 56:1907–1911, 1966.
- [Mél96] S. Méléard. *Asymptotic behaviour of some interacting particle systems; McKean-Vlasov and Boltzmann models*, pages 42–95. Springer Berlin Heidelberg, Berlin, Heidelberg, 1996.
- [MM13] S. Mischler and C. Mouhot. Kac’s program in kinetic theory. *Inventiones mathematicae*, 193(1):1–147, Jul 2013.
- [MMW15] S. Mischler, C. Mouhot, and B. Wennberg. A new approach to quantitative propagation of chaos for drift, diffusion and jump processes. *Probability Theory and Related Fields*, 161(1):1–59, Feb 2015.
- [Mon17] P. Monmarché. Long-time behaviour and propagation of chaos for mean field kinetic particles. *Stochastic Process. Appl.*, 127(6):1721–1737, 2017.
- [Shi87] M. Shiino. Dynamical behavior of stochastic systems of infinitely many coupled nonlinear oscillators exhibiting phase transitions of mean-field type: H theorem on asymptotic approach to equilibrium and critical slowing down of order-parameter fluctuations. *Phys. Rev. A*, 36:2393–2412, Sep 1987.
- [SV79] Daniel W. Stroock and S. R. Srinivasa Varadhan. *Multidimensional diffusion processes*, volume 233 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin-New York, 1979.
- [Szn91] A.-S. Sznitman. Topics in propagation of chaos. In P.-L. Hennequin, editor, *Ecole d’Eté de Probabilités de Saint-Flour XIX — 1989*, pages 165–251, Berlin, Heidelberg, 1991. Springer Berlin Heidelberg.

- [Tam84] Y. Tamura. On asymptotic behaviors of the solution of a nonlinear diffusion equation. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 31(1):195–221, 1984.
- [Tug10] J. Tugaut. *Self-stabilizing processes in a multi-wells landscape*. Theses, Université Henri Poincaré - Nancy I, July 2010.
- [Tug14a] J. Tugaut. Phase transitions of McKean-Vlasov processes in double-wells landscape. *Stochastics*, 86(2):257–284, 2014.
- [Tug14b] J. Tugaut. Self-stabilizing processes in multi-wells landscape in \mathbb{R}^d -invariant probabilities. *J. Theoret. Probab.*, 27(1):57–79, 2014.
- [WLEC17] C. Wang, Q. Li, W. E, and B. Chazelle. Noisy Hegselmann-Krause systems: phase transition and the $2R$ -conjecture. *J. Stat. Phys.*, 166(5):1209–1225, 2017.