

Lattice self-similar sets on the real line are not Minkowski measurable

Kombrink, Sabrina; Winter, Steffen

DOI:

[10.1017/etds.2018.26](https://doi.org/10.1017/etds.2018.26)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Kombrink, S & Winter, S 2020, 'Lattice self-similar sets on the real line are not Minkowski measurable', *Ergodic Theory and Dynamical Systems*, vol. 40, no. 1, pp. 221-232. <https://doi.org/10.1017/etds.2018.26>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This article has been published in a revised form in *Ergodic Theory and Dynamical Systems*, DOI: <https://doi.org/10.1017/etds.2018.26>. This version is free to view and download for private research and study only. Not for re-distribution or re-use. © Cambridge University Press, 2018.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Lattice self-similar sets on the real line are not Minkowski measurable

Sabrina Kombrink[†] and Steffen Winter[‡] [†]

[†] *Universität zu Lübeck, Institut für Mathematik, Ratzeburger Allee 160, 23562
Lübeck, Germany*

(e-mail: kombrink@math.uni-luebeck.de)

[‡] *Karlsruhe Institute of Technology, Department of Mathematics, Englerstr. 2,
76131 Karlsruhe, Germany*

(e-mail: steffen.winter@kit.edu)

(Received 02/03/2018)

Abstract. We show that any nontrivial self-similar subset of the real line that is invariant under a lattice iterated function system (IFS) satisfying the open set condition (OSC) is not Minkowski measurable. So far, this was only known for special classes of such sets. Thereby, we provide the last puzzle-piece in proving that under OSC a nontrivial self-similar subset of the real line is Minkowski measurable iff it is invariant under a nonlattice IFS, a 25-year-old conjecture.

1. Introduction

The Minkowski content was proposed by B. B. Mandelbrot [[Man95](#)] as texture parameter for irregular sets (a measure of “lacunarity”). Indeed, the Minkowski content can be used to understand the geometry of a fractal set beyond its (Hausdorff or Minkowski) dimension and in particular is a tool to distinguish between sets of the same dimension. Besides its geometric relevance, the Minkowski content has attracted attention in connection with the Weyl-Berry conjecture concerning the distribution of the eigenvalues of the Laplacian on bounded domains $\Omega \subseteq \mathbb{R}^d$ with fractal boundaries. More precisely, M. L. Lapidus and C. Pomerance showed in [[LP93](#)] that if $\Omega \subseteq \mathbb{R}$, then the second asymptotic term of the eigenvalue counting function can be expressed in terms of the Minkowski dimension and the Minkowski content of the boundary of Ω , whenever these quantities exist. However, although

[†] This research was initiated and carried out while the authors were staying at the Institut Mittag-Leffler. The authors would like to thank the staff of the institute as well as the participants and organisers of the 2017 research programme *Fractal Geometry and Dynamics* for the stimulating atmosphere, the excellent working environment, warm hospitality and financial support.

much progress has been made in recent years, in general it is not easy to decide, whether the Minkowski content of a given fractal set exists or not.

Assuming the open set condition (OSC) it was conjectured in [Lap93, Conjecture 3] (see also [Gat00, Section 5.2]) that a nontrivial self-similar set is Minkowski measurable (i. e. its Minkowski content exists, and is positive and finite) iff it arises from a nonlattice iterated function system (IFS). The progress in resolving this conjecture is as follows: Self-similar subsets of \mathbb{R} generated from nonlattice IFS satisfying OSC were shown to be Minkowski measurable in [Lap93, Fal95, Gat00] (the results of [Gat00] hold for self-similar subsets of \mathbb{R}^d , too). For nontrivial self-similar subsets of \mathbb{R} the converse, i. e. lattice sets are not Minkowski measurable, was shown in [LvF00] under additional assumptions. These assumptions address the geometric structure of the underlying feasible open set for the OSC and have been weakened in [KK15, KPW16], see Section 2.3 for more details. However, up to now the conjecture remained unresolved for large classes of self-similar sets, see Section 2.4 for examples.

In the present article we fully remove the assumptions of [LvF00, KK15, KPW16] and in this way provide the last puzzle-piece in proving that under OSC a nontrivial self-similar subset of \mathbb{R} is Minkowski measurable iff it arises from a nonlattice IFS. This resolves the conjecture stated in [Lap93, Conjecture 3] and [Gat00, Section 5.2] for self-similar sets in \mathbb{R} .

The article is organised as follows. After some preliminaries in Sections 2.1 and 2.2 we give a brief exposition of the key results from the literature in Section 2.3. A class of self-similar sets for which Minkowski measurability had previously not been understood is discussed in Section 2.4. Our main results are stated in Section 3 and proved in Section 4. We conclude by showing in Section 5 that for sets in \mathbb{R} the above-mentioned results from [KK15, KPW16] are equivalent.

2. Preliminaries

2.1. *Minkowski measurability* Let A denote a compact subset of the one-dimensional Euclidean space $(\mathbb{R}, |\cdot|)$ and let $\varepsilon > 0$. Define the ε -parallel set of A to be $A_\varepsilon := \{x \in \mathbb{R} \mid \inf_{a \in A} |x - a| \leq \varepsilon\}$. If the *Minkowski dimension* $\dim_M(A) := 1 - \lim_{\varepsilon \searrow 0} \log(\lambda(A_\varepsilon)) / \log(\varepsilon)$ exists, then we consider the *rescaled volume function* $\varepsilon \mapsto \varepsilon^{\dim_M(A)-1} \lambda(A_\varepsilon)$ defined on $(0, \infty)$, where λ denotes Lebesgue measure in \mathbb{R} . If its limit as $\varepsilon \searrow 0$ exists, then we write

$$\mathcal{M}(A) := \lim_{\varepsilon \searrow 0} \varepsilon^{\dim_M(A)-1} \lambda(A_\varepsilon)$$

and call this value the *Minkowski content* of A . If $\mathcal{M}(A)$ exists, and is positive and finite then we say that A is *Minkowski measurable*.

2.2. *Self-similar sets, open set condition, (non-)lattice and nontrivial* We let $\Phi := \{\phi_1, \dots, \phi_N\}$ with $N \in \mathbb{N}$, $N \geq 2$ denote an iterated function system (IFS) consisting of similarities ϕ_j acting on \mathbb{R} . Note that the ϕ_j are not required to be orientation preserving. Suppose that the IFS Φ satisfies the *open set condition*

(OSC), that is, there exists a nonempty open set O such that

$$\phi_i(O) \subseteq O \quad \text{and} \quad \phi_i(O) \cap \phi_j(O) = \emptyset \quad \text{for } i, j \in \Sigma, i \neq j \quad (2.1)$$

where $\Sigma := \{1, \dots, N\}$. Any nonempty open set O satisfying (2.1) shall be called a *feasible open set* for the IFS Φ . Let r_i denote the similarity ratio of ϕ_i . We say that Φ is *lattice*, if the set $\{\log(r_i) \mid i \in \Sigma\}$ generates a discrete subgroup of $(\mathbb{R}, +)$. Otherwise, Φ is said to be *nonlattice*. If Φ is lattice, then there exists a maximal $a > 0$ such that $\{\log(r_i) \mid i \in \Sigma\} \subseteq a\mathbb{Z}$ and we call $r := e^a$ the *base* of Φ .

The natural action of Φ on the class of subsets of \mathbb{R} is defined via $\Phi A := \bigcup_{i \in \Sigma} \phi_i A$ for $A \subseteq \mathbb{R}$. By Hutchinson's theorem, there exists a unique nonempty compact set F satisfying the invariance relation $\Phi F = F$. This set F is called the *self-similar set* associated with Φ . It is well-known that under OSC $\dim_M(F)$ exists and that $\#F > 1$, where $\#$ denotes the cardinality.

F is called *nontrivial* if $\dim_M(F) < 1$. Nontriviality of F is equivalent to the assertion that any feasible open set O satisfies $O \setminus \overline{\Phi O} \neq \emptyset$, see [PW12, Corollary 5.6]. Here, \overline{B} and ∂B denote the topological closure and boundary of a set B respectively. A feasible open set O for Φ is called *strong*, if it has nonempty intersection with F , i. e. $O \cap F \neq \emptyset$. Moreover, following [KPW16, Win15] O is called *compatible*, if $\partial O \subseteq F$. (Notice, in [PW12] O is called *compatible* if $\partial \overline{O} \subseteq F$, which is a weaker condition on O .) Let π_F denote the *metric projection* onto F , which is defined on the set of points $x \in \mathbb{R}$ with a unique nearest neighbour y in F by $\pi_F(x) := y$. The set O is said to satisfy the *projection condition* if $\phi_i O \subseteq \pi_F^{-1}(\phi_i F)$ for $i \in \Sigma$.

2.3. *Known results on Minkowski measurability of self-similar sets in \mathbb{R}* Let $F \subseteq \mathbb{R}$ be the self-similar set of an IFS Φ as defined in Section 2.2 and let I denote the interior of the convex hull of F , that is, \overline{I} is the smallest closed interval containing F . Note that since F is not a singleton, I is nonempty.

THEOREM 2.1. *Suppose that Φ satisfies OSC.*

- (i) [Lap93, Fal95] *If Φ is nonlattice and $\phi_i(\overline{I}) \cap \phi_j(\overline{I}) = \emptyset$ for $i \neq j$ (i. e. the strong separation condition is satisfied), then F is Minkowski measurable.*
- (ii) [Gat00] *If Φ is nonlattice, then F is Minkowski measurable.*
- (iii) [LvF00] *If Φ is lattice, F is nontrivial and I is a feasible open set for Φ , then F is not Minkowski measurable.*
- (iv) [KK15] *If Φ is lattice, F is nontrivial and $\Phi^m I$ is a feasible open set for Φ for some $m \in \mathbb{N}_0$, then F is not Minkowski measurable.*
- (v) [KPW16] *Assume existence of a strong feasible open set O for Φ that satisfies the projection condition and for which there exists a finite partition of $(0, \infty)$ so that $\varepsilon \mapsto \lambda(F_\varepsilon \cap (O \setminus \Phi O))$ is polynomial on each partition interval. If Φ is lattice and F is nontrivial, then F is not Minkowski measurable.*

We point out that the results of [Gat00, KPW16] stated above in (ii) and (v) hold in arbitrary dimension. For self-similar subsets of \mathbb{R} the assumptions in (iv) and (v) are equivalent, which we prove below in Theorem 5.1. To clarify that there exist lattice self-similar sets which are not covered by the results (iii)–(v), we now discuss some examples with more complicated feasible open sets.

2.4. *Self-similar sets with complicated feasible open sets* Let $A > 1$ and let $\mathcal{D} := \{d_1, \dots, d_N\} \subseteq \mathbb{R}$ be a digit set. Define similarities ϕ_j acting on \mathbb{R} by $\phi_j(x) = (x + d_j)/A$ for $j \in \{1, \dots, N\}$. Further, let

$$\mathcal{D}_1 := \mathcal{D}, \quad \mathcal{D}_n := \mathcal{D} + A\mathcal{D}_{n-1}, \quad n \geq 2 \quad \text{and} \quad \mathcal{D}_\infty := \bigcup_{n=1}^\infty \mathcal{D}_n.$$

By [HL08, Theorem 4.4] the IFS $\Phi := \{\phi_1, \dots, \phi_N\}$ satisfies OSC iff \mathcal{D}_∞ is uniformly discrete and $\#\mathcal{D}_k = N^k$ for all $k \geq 1$. (\mathcal{D}_∞ is *uniformly discrete* if there exists $r > 0$ so that $|x - y| \geq r$ for all $x \neq y \in \mathcal{D}_\infty$.) Thus, if one chooses A, d_1, \dots, d_N to be nonnegative integers, then OSC is satisfied iff $d_i \not\equiv d_j \pmod{A}$ for $i \neq j$. Depending on the choice of A and \mathcal{D} feasible open sets can be rather complicated. E. g. for the IFS Φ given by $N = 3$, $A = 4$, $d_1 = 0$, $d_2 = 1$ and $d_3 = 6$, displayed in Figure 1, OSC is satisfied but the assumptions of Theorem 2.1(iii)–(v) are violated, which can be seen as follows. For (iii) and (iv) we provide a proof in the next paragraph. The statement for (v) then directly follows from the equivalence of (iv) and (v) which we prove in Theorem 5.1 below.

Fix $m \in \mathbb{N}_0$ and let $U := \Phi^m I$, where $I = (0, 2)$ in this example. We claim that U is not feasible for Φ . Without loss of generality we can assume that m is odd, i. e. $m = 2k + 1$ for some $k \in \mathbb{N}_0$, since feasibility of $\Phi^m I$ would imply feasibility of $\Phi^{m+1} I$. Writing $\phi_\omega := \phi_{\omega_1} \circ \dots \circ \phi_{\omega_n}$ for $\omega \in \Sigma^n$ the claim directly follows from

$$\phi_1(U) \cap \phi_2(U) \supset \phi_1(\phi_3 \phi_{23}^k(I)) \cap \phi_2(\phi_{23}^k \phi_2(I)) \neq \emptyset, \quad (2.2)$$

which we now prove: First observe that $\phi_1(0) = 0$, $\phi_3(2) = 2$ and $\phi_{23}(2/3) = 2/3$. Second, note that for the left endpoints of the intervals $\phi_{13} \phi_{23}^k(I)$ and $\phi_2 \phi_{23}^k \phi_2(I)$ we have

$$\begin{aligned} \phi_{13} \phi_{23}^k(0) - \phi_2 \phi_{23}^k \phi_2(0) &= \phi_{13} \left(\phi_{23}^k \left(\frac{2}{3} \right) - \frac{2}{3} \left(\frac{1}{4} \right)^{2k} \right) - \phi_2 \left(\phi_{23}^k \left(\frac{2}{3} \right) - \frac{5}{12} \left(\frac{1}{4} \right)^{2k} \right) \\ &= \left(\frac{1}{4} \right)^{2k+2} > 0. \end{aligned}$$




FIGURE 1. The interval $[0, 2]$ and its images under the first two iterations of the IFS $\{\frac{x}{4}, \frac{x+1}{4}, \frac{x+6}{4}\}$. The numbers above the intervals indicate their coding, e. g. ‘12’ encodes $\phi_1 \phi_2([0, 2])$.

Third, the intervals $\phi_{13}\phi_{23}^k(I)$ and $\phi_2\phi_{23}^k\phi_2(I)$ both have length $2 \cdot (1/4)^{2k+2}$. Therefore, they must overlap in an interval of length $(1/4)^{2k+2}$, showing (2.2).

Remark 2.2. Indeed, in the above example, any feasible open set necessarily has an infinite number of connected components, disqualifying, in particular, the sets $\Phi^m I$. This was pointed out to us by Christoph Bandt, whom we wish to thank for sharing the following arguments with us:

The *dynamical boundary* of F associated with Φ is the set $\text{db}(F) := \bigcup_h F \cap hF$, where the union is taken over all *neighbour maps* h , i. e. maps of the form $h = \phi_u^{-1}\phi_\omega$, where $u, \omega \in \Sigma^n$ for some $n \in \mathbb{N}$ are so that $\phi_u(F) \cap \phi_\omega(F) \neq \emptyset$ and $u_1 \neq \omega_1$. When $x \in \text{db}(F)$ then $\phi_u(x) \in \phi_u(F) \cap \phi_\omega(F)$. Thus, any feasible open set O for Φ may not intersect $\text{db}(F)$. On the other hand, $\text{db}(F) \subseteq F$ and whence $\text{db}(F) \subseteq \overline{O}$. Therefore, $\text{db}(F) \subseteq \partial O$. Now, if the dynamical boundary has infinite cardinality (which is the case here, see below), then O necessarily has infinitely many connected components.

In [BM09] general statements were obtained to determine the cardinality of the dynamical boundary of a limit set of a graph-directed system via neighbour graphs. The neighbour graph associated to the present example is depicted in Figure 2. Its root is the identity and its vertices are the neighbour maps. “An arrow with label i, j is drawn from vertex h to vertex \bar{h} if $\bar{h} = \phi_i^{-1}h\phi_j$ for two marks $i, j \in \Sigma$. We keep only those arrows which correspond to proper neighbors, that is $\phi_i(F) \cap h\phi_j(F) \neq \emptyset$.” [BMT18]. In our example $\phi_i(F) \cap \phi_j(F) \neq \emptyset$ iff $(i, j) \in \{(1, 2), (2, 1)\}$. Therefore, there are precisely two arrows leaving the root vertex id . The first arrow, labeled with ‘1,2’, terminates in the vertex $\phi_1^{-1}\phi_2$ with $\phi_1^{-1}\phi_2(x) = x + 1$. The second arrow, labeled with ‘2,1’, terminates in the vertex $\phi_2^{-1}\phi_1$ with $\phi_2^{-1}\phi_1(x) = x - 1$. The other vertices and arrows are constructed accordingly. Using the terminology from [BM09] the light shaded vertices are intermediate and the dark shaded ones are terminal. According to [BM09, Theorem 7] the terminal and intermediate vertices correspond to subsets of the dynamical boundary with cardinality one and countably infinite respectively. Thus, $\text{db}(F)$ is countably infinite here.




FIGURE 2. Neighbour graph for the IFS $\{\frac{x}{4}, \frac{x+1}{4}, \frac{x+6}{4}\}$.

3. Main results

THEOREM 3.1. *If F is a nontrivial self-similar set in \mathbb{R} generated by a lattice IFS Φ satisfying OSC, then F is not Minkowski measurable.*

Together with Theorem 2.1 (ii) we thus verify the conjecture of [Lap93, Conjecture 3] and [Gat00, Section 5.2] for self-similar sets in \mathbb{R} :

COROLLARY 3.2. *Suppose that F is a nontrivial self-similar set in \mathbb{R} generated by an IFS Φ satisfying OSC. Then F is Minkowski measurable iff Φ is nonlattice.*

Remark 3.3. The nontriviality condition, $\dim_M(F) < 1$, is necessary in the statements of Theorem 3.1, Corollary 3.2 and Theorem 2.1(iii)–(v) and cannot be removed: The unit interval $X := [0, 1]$ has Minkowski dimension $\dim_M(X) = 1$. It is the self-similar set associated with the lattice IFS $\{x \mapsto x/2, x \mapsto (x+1)/2\}$ acting on \mathbb{R} . However, its Minkowski content $\mathcal{M}(X) = \lim_{\varepsilon \rightarrow 0} (1 + 2\varepsilon) = 1$ exists as a positive and finite value. Hence X is Minkowski measurable. In fact, any self-similar set F in \mathbb{R} with $\dim_M F = 1$ is Minkowski measurable, see e. g. [KPW16, Theorem 1.1(i)].

A key ingredient in the proof of Theorem 3.1 is the construction of a relatively simple strong feasible open set, see Theorem 3.4 below and its proof. With this set at hand we can deduce Minkowski non-measurability from [KPW16, Theorem 3.1], see Theorem 4.1 below. Define $\Sigma^* := \bigcup_{n=0}^{\infty} \Sigma^n$ with $\Sigma^0 := \{\emptyset\}$, where \emptyset denotes the empty word. Moreover, for $\omega = (\omega_1, \dots, \omega_n) \in \Sigma^*$ write $\phi_\omega := \phi_{\omega_1} \circ \dots \circ \phi_{\omega_n}$ and let ϕ_\emptyset be the identity map.

THEOREM 3.4. *Let $F \subseteq \mathbb{R}$ be the self-similar set generated by an IFS Φ satisfying OSC. Then there exists a strong and compatible feasible open set U for Φ , i. e. one which satisfies $U \cap F \neq \emptyset$ and $\partial U \subseteq F$.*

What is more, there always exists such a set U that can be generated from a finite union of elementary intervals $\phi_u(I)$: there exist $m \in \mathbb{N}_0$ and $\Lambda \subseteq \Sigma^m$ such that

$$U_\Lambda := \bigcup_{\omega \in \Sigma^*} \phi_\omega \bigcup_{u \in \Lambda} \phi_u(I) \quad (3.1)$$

defines a strong and compatible feasible open set for F .

Remark 3.5. For any $m \in \mathbb{N}_0$ and any nonempty $\Lambda \subseteq \Sigma^m$, the set U_Λ in (3.1) has nonempty intersection with F , since $I \cap F \neq \emptyset$. Moreover, U_Λ is compatible, because $\partial U_\Lambda \subseteq \overline{\bigcup_{\omega \in \Sigma^*} \phi_\omega \bigcup_{u \in \Lambda} \phi_u(\partial I)} \subseteq F$, where the last inclusion follows since $\partial I \subseteq F$ and $\phi_\omega F \subseteq F$ for any $\omega \in \Sigma^*$. However, it is not obvious that U_Λ is a feasible open set and this is indeed only true for particular choices of Λ .

4. Proofs

4.1. *Construction of a feasible open set U_Λ – Proof of Theorem 3.4* Obviously, the first statement of the theorem follows from the second. In view of Remark 3.5, it therefore suffices to show that at least one of the sets U_Λ (defined by (3.1)) is feasible. First observe that for any $m \in \mathbb{N}$ and any nonempty $\Lambda \subseteq \Sigma^m$ the set U_Λ

is nonempty and open, since I has these properties, and that $\phi_i U_\Lambda \subseteq U_\Lambda$ for any $i \in \Sigma$. Therefore, all that remains to be shown is existence of a set $\Lambda \subseteq \Sigma^m$ for some $m \in \mathbb{N}_0$ such that $\phi_i(U_\Lambda) \cap \phi_j(U_\Lambda) = \emptyset$ for any $i \neq j \in \Sigma$. For this we adapt Schief's construction in [Sch94] of a strong feasible open set:

Let r_ω denote the similarity ratio of ϕ_ω for $\omega = (\omega_1, \dots, \omega_n) \in \Sigma^*$. Note that $r_\omega = r_{\omega_1} \cdots r_{\omega_n}$. Fix $\varepsilon \in (0, 1/6)$. Schief showed [Sch94, proof of Theorem 2.1] that there exists $\kappa \in \Sigma^*$ so that

$$O_\kappa := \bigcup_{u \in \Sigma^*} \phi_u \kappa (F_\varepsilon)$$

is a feasible open set for Φ . As $r_i < 1$ for all i , there is a minimal $k \in \mathbb{N}_0$ such that

$$\bigcup_{\omega \in \Sigma^k} \phi_\omega I \subseteq F_\varepsilon.$$

Set $m := k + |\kappa|$, where $|\kappa|$ denotes the length of κ , i. e. $\kappa \in \Sigma^{|\kappa|}$. Further, set $\Lambda := \{\kappa\omega \mid \omega \in \Sigma^k\}$. Then $\emptyset \neq \Lambda \subseteq \Sigma^m$ and $U_\Lambda \subseteq O_\kappa$, whence $\phi_i(U_\Lambda) \cap \phi_j(U_\Lambda) = \emptyset$ for any $i \neq j \in \Sigma$. This completes the proof of Theorem 3.4.

4.2. *A criterion for Minkowski measurability.* In the proof of Theorem 3.1 we will make use of a general Minkowski measurability criterion for self-similar sets in \mathbb{R}^d (satisfying OSC) derived in [KPW16]. It is based on feasible open sets satisfying the projection condition and was obtained via classical renewal theory. We briefly restate a version of this criterion here, boiled down to our present one-dimensional setting. Given Φ , O and F as in Section 2 we set

$$\Gamma := O \setminus \Phi O \quad \text{and} \quad g := \sup \left\{ \inf_{y \in F} |x - y| \mid x \in \Gamma \right\}. \quad (4.1)$$

THEOREM 4.1. [KPW16, Theorem 3.1 and Corollary 3.2] *Let $F \subseteq \mathbb{R}$ be a nontrivial self-similar set generated by a lattice IFS Φ with base r . Suppose that Φ satisfies OSC with a strong feasible set O satisfying the projection condition. Let $D := \dim_M(F)$ and let Γ and g be defined as in (4.1). Define the function $p : (rg, g] \rightarrow \mathbb{R}$ by*

$$p(\varepsilon) := \varepsilon^{D-1} \left[\frac{\lambda(\Gamma)}{r^{D-1} - 1} + \sum_{\ell=0}^{\infty} r^{\ell(D-1)} \lambda(F_{r^\ell \varepsilon} \cap \Gamma) \right]. \quad (4.2)$$

Then F is Minkowski measurable iff p is constant on $(rg, g]$.

Note that the series in the definition of p is uniformly convergent in ε , see [KPW16, proof of Theorem 3.1].

Remark 4.2. It is easily seen that a feasible open set of the form U_Λ given in (3.1) satisfies the projection condition. In fact, any strong and compatible feasible open set O satisfies the projection condition, see e. g. [Win15, Remark 3.20].

4.3. *Minkowski non-measurability – Proof of Theorem 3.1* Let r denote the lattice base of Φ and let U be a strong and compatible feasible open set for Φ . Such a set U exists due to Theorem 3.4. We want to apply Theorem 4.1 and note that all its assumptions are satisfied; in particular, the projection condition, see Remark 4.2. We infer that the set F is Minkowski measurable iff the function p defined in (4.2) is constant.

In the following we will demonstrate that the properties of U imply that p cannot be constant. We pursue a proof by contradiction, whence assume that there exists $C > 0$ so that $p(\varepsilon) = C$ or, equivalently,

$$L(\varepsilon) := C\varepsilon^{1-D} - \frac{\lambda(\Gamma)}{r^{D-1} - 1} = \sum_{\ell=0}^{\infty} r^{\ell(D-1)} \lambda(F_{r^\ell \varepsilon} \cap \Gamma) =: R(\varepsilon) \quad (4.3)$$

for $\varepsilon \in (rg, g]$. Define $G := U \setminus \overline{\Phi U}$. Clearly, G is open and $G \subseteq \Gamma$. Moreover, $\lambda(\Gamma \setminus G) = 0$, since $\Gamma \setminus G = U \cap \partial \Phi U \subseteq U \cap \Phi \partial U \subseteq F$ and $\dim_M(F) < 1$. Therefore, $\lambda(F_{r^\ell \varepsilon} \cap \Gamma) = \lambda(F_{r^\ell \varepsilon} \cap G)$. As stated in Section 2.2, nontriviality implies $G \neq \emptyset$. Hence G has countably many connected components G_j , $j \in J$, each of which is an open interval. Without loss of generality suppose that either $J = \mathbb{N}_0$ or $J = \{0, \dots, n\}$ for some $n \in \mathbb{N}_0$. Write $\text{diam}(G_j)$ for the diameter of G_j and assume that the G_j are ordered so that $\text{diam}(G_{j-1}) \geq \text{diam}(G_j)$ for all $j \in J \setminus \{0\}$. Since $\partial G_j \subseteq F$ and $G_j \cap F = \emptyset$, we have

$$\lambda(F_t \cap G_j) = \begin{cases} 2t & : 0 < 2t \leq \text{diam}(G_j), \\ \text{diam}(G_j) & : 2t > \text{diam}(G_j). \end{cases} \quad (4.4)$$

For $\ell \in \mathbb{N}_0$, $j \in J$ define $f_{\ell,j} : (rg, g] \rightarrow \mathbb{R}$ by

$$f_{\ell,j}(\varepsilon) := r^{\ell(D-1)} \lambda(F_{r^\ell \varepsilon} \cap G_j).$$

Then $R(\varepsilon) = \sum_{\ell=0}^{\infty} \sum_{j \in J} f_{\ell,j}(\varepsilon)$. Let $f_{\ell,j}^{(-)}(\varepsilon)$ and $f_{\ell,j}^{(+)}(\varepsilon)$ denote the left and right derivatives of $f_{\ell,j}$ at ε respectively. By (4.4), we have that

$$f_{\ell,j}^{(-)}(\varepsilon) \geq f_{\ell,j}^{(+)}(\varepsilon) \geq 0 \text{ for } \ell \in \mathbb{N}_0, j \in J \text{ and } \varepsilon \in (rg, g). \quad (4.5)$$

In fact, since $\lambda(F_\varepsilon \cap G_j)$ is piecewise linear with at most two different slopes, the derivative $f'_{\ell,j}(\varepsilon)$ of $f_{\ell,j}$ exists at all $\varepsilon \in (rg, g)$ except for at most one point.

LEMMA 4.3. *The series $\sum_{\ell=0}^{\infty} \sum_{j \in J} f_{\ell,j}^{(+)}$ and $\sum_{\ell=0}^{\infty} \sum_{j \in J} f_{\ell,j}^{(-)}$ converge uniformly on (rg, g) .*

Proof. Let $N_\varepsilon(\ell) := \#\{j \in J \mid 2r^\ell \varepsilon \leq \text{diam}(G_j)\}$. As remarked in Section 4.2, the series $\sum_{\ell=0}^{\infty} r^{\ell(D-1)} \lambda(F_{r^\ell \varepsilon} \cap \Gamma)$ from (4.2) is uniformly convergent in ε . Thus, there exists a sequence $(c_n)_n$ so that $\lim_{n \rightarrow \infty} c_n = 0$ and so that for $\varepsilon \in (rg, g]$, $n \in \mathbb{N}$

$$c_n \geq \sum_{\ell=n}^{\infty} r^{\ell(D-1)} \sum_{j \in J} \lambda(F_{r^\ell \varepsilon} \cap G_j) \geq \sum_{\ell=n}^{\infty} r^{\ell(D-1)} \sum_{j=0}^{N_\varepsilon(\ell)-1} 2r^\ell \varepsilon = \varepsilon \sum_{\ell=n}^{\infty} 2r^{\ell D} N_\varepsilon(\ell).$$

Since $f_{\ell,j}^{(-)}(\varepsilon) = 0$ if $2r^\ell\varepsilon > \text{diam}(G_j)$, this yields

$$\sum_{\ell=n}^{\infty} \sum_{j \in J} f_{\ell,j}^{(-)}(\varepsilon) = \sum_{\ell=n}^{\infty} \sum_{j=0}^{N_\varepsilon(\ell)-1} 2r^{\ell D} = \sum_{\ell=n}^{\infty} 2r^{\ell D} N_\varepsilon(\ell) \leq \frac{c_n}{\varepsilon} \leq \frac{c_n}{rg}$$

which proves uniform convergence of $\sum_{\ell=0}^{\infty} \sum_{j \in J} f_{\ell,j}^{(-)}$ and by (4.5) also of the series $\sum_{\ell=0}^{\infty} \sum_{j \in J} f_{\ell,j}^{(+)}$. \square

Remark 4.4. Observe that $f_{\ell,j}$ are Kneser functions of order 1, i. e. they satisfy

$$f_{\ell,j}(\mu b) - f_{\ell,j}(\mu a) \leq \mu(f_{\ell,j}(b) - f_{\ell,j}(a)),$$

for all $a, b \in (rg, g)$ with $a \leq b$ and any $\mu \geq 1$ such that $\mu b < g$. This can be checked directly, but it also follows from [Sta76, Lemma 5], since the intervals G_j are metrically associated with F (meaning that for each point $x \in G_j$ there is a point $y \in F$ with $|x - y| = \inf_{a \in F} |x - a|$ such that the whole segment between x and y is contained in G_j) and therefore $\lambda(F_t \cap G_j)$ is a Kneser function of order 1 on $(0, \infty)$. Hence, the assertion of Lemma 4.3 is a special case of [Sta76, Lemma 4].

In order to obtain a contradiction, we consider two cases:

CASE 1: There exist $\ell^* \in \mathbb{N}_0$, $j^* \in J$ and $x \in (rg, g)$ so that $f_{\ell^*,j^*}^{(-)}(x) \neq f_{\ell^*,j^*}^{(+)}(x)$. Equation (4.4) implies

$$f_{\ell^*,j^*}^{(-)}(x) = 2r^{\ell^* D} > 0 = f_{\ell^*,j^*}^{(+)}(x) \quad (4.6)$$

Lemma 4.3 shows that the right and left derivatives of R at x exist and are given by $R^{(+)}(x) = \sum_{\ell=0}^{\infty} \sum_{j \in J} f_{\ell,j}^{(+)}(x)$ and $R^{(-)}(x) = \sum_{\ell=0}^{\infty} \sum_{j \in J} f_{\ell,j}^{(-)}(x)$. With (4.5) and (4.6) we thus obtain

$$R^{(-)}(x) - R^{(+)}(x) \geq f_{\ell^*,j^*}^{(-)}(x) - f_{\ell^*,j^*}^{(+)}(x) = 2r^{\ell^* D} > 0.$$

Hence, unlike the function L , the function R is not differentiable at x , contradicting (4.3).

CASE 2: The derivative $f'_{\ell,j}$ exists on (rg, g) for all $\ell \in \mathbb{N}_0$, $j \in J$.

In this case, for any $j \in J$ there exists $k = k(j) \in \mathbb{N}_0$ so that $\text{diam}(G_j) = 2r^k g$, yielding

$$f'_{\ell,j} \equiv 2r^{\ell D} \text{ on } (rg, g) \text{ for all } \ell \geq k(j), \text{ and } f'_{\ell,j} \equiv 0 \text{ otherwise.} \quad (4.7)$$

By Lemma 4.3, R' exists and coincides with $\sum_{\ell=0}^{\infty} \sum_{j \in J} f'_{\ell,j}$ which by (4.7) is constant on (rg, g) . However, $L'(\varepsilon) = C(1 - D)\varepsilon^{-D}$ which, due to the nontriviality of F (and since $C > 0$), is clearly not constant. Therefore, we obtain a contradiction to (4.3) also in the second case. This completes the proof of Theorem 3.1.

5. Equivalence of (iv) and (v) of Theorem 2.1

During our discussions the question arose whether the classes of self-similar subsets of \mathbb{R} covered by the assertions (iv) and (v) of Theorem 2.1 are equivalent. The following statement gives an affirmative answer (irrespective of the IFS being lattice or nonlattice).

THEOREM 5.1. *Let Φ be an IFS in \mathbb{R} satisfying OSC such that the associated invariant set F is nontrivial. Then the following assertions are equivalent.*

- (i) $\Phi^m I$ is a feasible open set for Φ for some $m \in \mathbb{N}_0$.
- (ii) There exists a strong feasible open set O for Φ , satisfying the projection condition, for which there exists a finite partition of $(0, \infty)$ so that $\varepsilon \mapsto \lambda(F_\varepsilon \cap (O \setminus \Phi O))$ is polynomial on each partition interval.

Proof. To begin with, note that for any feasible open set of the form $\Phi^m I$ there exists a finite partition of $(0, \infty)$ so that $\varepsilon \mapsto \lambda(F_\varepsilon \cap (\Phi^m I \setminus \Phi^{m+1} I))$ is polynomial on each partition interval. Therefore, (i) implies (ii).

For the converse, suppose that O is as in (ii). Consider $U := \text{int}(\overline{O})$, where int denotes the topological interior. Then $U = \bigcup_{i \in E} U_i$ is a union of open intervals U_i with the property that the distance between any two U_i is strictly positive. Let $\tilde{E} := \{i \in E : U_i \cap I \neq \emptyset\}$. The key part of the proof is to show that

$$\#\tilde{E} < \infty. \quad (5.1)$$

Before proving (5.1) we demonstrate that (5.1) implies assertion (i). Since O is a strong feasible open set for Φ , so is U , which can be seen by contradiction. Therefore, $F \subseteq \overline{U}$ and so, by (5.1), $F \subseteq \overline{U} \cap \overline{I} \subseteq \bigcup_{i \in \tilde{E}} \overline{U}_i$, which implies that there exists $m \in \mathbb{N}$ so that $\Phi^m \overline{I} \subseteq \bigcup_{i \in \tilde{E}} \overline{U}_i$ (simply choose m large enough that, for any $w \in \Sigma^m$, $\text{diam}(\phi_w \overline{I})$ is smaller than the minimal distance between the finitely many U_i). The property that the U_i have positive distance to one another implies that $\Phi^m I \subseteq \bigcup_{i \in E} U_i$. From this inclusion it is easy to see that $\Phi^m I$ is feasible for Φ , whence assertion (i) holds.

To verify (5.1) let $c_1, \dots, c_k \in (0, \infty)$ denote the partition points of the partition of $(0, \infty)$ associated with O . Let $\{H_j\}_{j \in J}$ denote the collection of connected components of $\text{int}(O \setminus \Phi O)$. Clearly, each H_j is an open interval and it is easy to see that $H_j \cap F = \emptyset$. We show (5.1) in four steps. Our first one is to prove

$$(I) \quad \#J < \infty.$$

For this, set $h_j := \inf\{\varepsilon > 0 : H_j \subseteq F_\varepsilon\}$. Observe that $\varepsilon \mapsto \lambda(F_\varepsilon \cap H_j)$ is constant (and equal to $\lambda(H_j)$) for $\varepsilon > h_j$, and linear (and nonconstant) in a left neighborhood of h_j . In particular, the function $\varepsilon \mapsto \lambda(F_\varepsilon \cap H_j)$ is not differentiable at h_j and so h_j must be one of the partition points c_ℓ . Next we show that for each of the finitely many c_ℓ the associated set $J_\ell := \{j \in J \mid h_j = c_\ell\}$ is of finite cardinality: For $j \in J_\ell$ let \widetilde{H}_j be the largest open interval (or one of the two in case of non-uniqueness) satisfying $H_j \cap \widetilde{H}_j \neq \emptyset$ and $\widetilde{H}_j \subseteq F_{<c_\ell} \setminus F$, where $F_{<c_\ell} := \{x \in \mathbb{R} \mid \inf_{a \in F} |x - a| < c_\ell\}$. By construction, $\{\widetilde{H}_j\}_{j \in J_\ell}$ is a pairwise disjoint family (here it is important to restrict to $j \in J_\ell$). Furthermore, $\lambda(\widetilde{H}_j) = c_\ell$. Therefore,

$$\#J_\ell = \frac{1}{c_\ell} \sum_{j \in J_\ell} \lambda(\widetilde{H}_j) = \frac{1}{c_\ell} \lambda\left(\bigcup_{j \in J_\ell} \widetilde{H}_j\right) \leq \frac{1}{c_\ell} \lambda(F_{c_\ell}) < \infty,$$

whence $\#J = \sum_{\ell=1}^k \#J_\ell < \infty$, showing (I).

Our second step is to verify that

(II) the number of connected components of $U \setminus \Phi U$ is finite.

For this, note that the family of connected components of $U \setminus \Phi U$ essentially coincides with $\{H_j\}_{j \in J}$, with the only possible differences occurring at the boundary points $\bigcup_{j \in J} \partial H_j$. More precisely, each H_j is a subset of $U \setminus \Phi U$ but not necessarily a connected component of this set, and $U \setminus \Phi U \subseteq \bigcup_{j \in J} \overline{H_j}$. Therefore, $U \setminus \Phi U$ has at most $\#J$ connected components and (I) implies (II).

For the third step let U_{i_1}, \dots, U_{i_n} denote those connected components of U which intersect $U \setminus \Phi U$ and let $E^* := \{i_1, \dots, i_n\}$ denote the respective index set. (The finiteness of E^* is clear from assertion (II).) We prove

(III) $\bigcup_{j \in E^*} U_j \cap F \neq \emptyset$.

If $i \in E \setminus E^*$ then $U_i \subseteq \Phi U = \bigcup_{j,k} \phi_k(U_j)$, which is a disjoint union by OSC and definition of the U_j . As furthermore U_i and each $\phi_k(U_j)$ is open and connected there exist $k \in \Sigma$, $j \in E$ so that $U_i \subseteq \phi_k(U_j)$. On the other hand, since $\phi_k(U_j)$ is a connected subset of U and intersects the connected component U_i , it must be contained in U_i . Thus $U_i = \phi_k(U_j)$, i.e. U_i is the precise image of U_j under ϕ_k . Amongst $\{U_i \mid i \in E \setminus E^*\}$ there is at least one largest bounded one, U_{i^*} , which needs to be an image of some U_j with $j \in E^*$ by the contraction property of the ϕ_k . Amongst $\{U_i \mid i \in E \setminus E^*\} \setminus \{U_{i^*}\}$ there again is a largest bounded one and inductively we see that each bounded U_i with $i \in E \setminus E^*$ is the image of one of the sets U_j , $j \in E^*$ and that also the possible unbounded components need to be amongst U_{i_1}, \dots, U_{i_n} . Since U is strong, we conclude $\bigcup_{j \in E^*} U_j \cap F \neq \emptyset$, showing (III).

In the final step we deduce (5.1) from the above: By (III), we can find a minimal $m \in \mathbb{N}$ and $\omega \in \Sigma^m$ so that $\phi_\omega I \subseteq \bigcup_{j \in E^*} U_j$. Let $\omega = \omega_1 \dots \omega_m$. If $\phi_{\omega_2 \dots \omega_m} I$ intersected infinitely many of the positively separated U_i , then $U \setminus \Phi U$ would have infinitely many connected components. (Assume that $\phi_{\omega_2 \dots \omega_m} I$ intersects infinitely many U_i , say U_{j_1}, U_{j_2}, \dots . Then on the one hand, $\phi_\omega I$ intersects all the sets $\phi_{\omega_1} U_{j_k}$, i.e. infinitely many. On the other hand, $\phi_\omega I$ is an interval and thus contained in a connected component V of U . Thus, as each $\phi_{\omega_1} U_{j_k}$ is connected, $\phi_{\omega_1} U_{j_k} \subseteq V$. Further, the family $\{\phi_{\omega_1} U_{j_k}\}_k$ is disjoint, which implies that the set $V \setminus \phi_{\omega_1} U$ and thus $V \setminus \Phi U$ and $U \setminus \Phi U$ must have infinitely many connected components.) This contradicts assertion (II). Thus, $\phi_{\omega_2 \dots \omega_m} I$ intersects only finitely many U_i . Inductively, one can now show that I can only intersect finitely many U_i . (Suppose that $\phi_{\omega_k \dots \omega_m} I$ intersects infinitely many U_i . Then with a similar argument as above, the same must hold for $\phi_{\omega_{k-1} \dots \omega_m} I$, which provides a contradiction to the previous induction step.) Hence $\#\tilde{E} < \infty$, proving (5.1). \square

REFERENCES

- [BM09] C. Bandt and M. Mesing. Self-affine fractals of finite type. *Banach Center Publications*, 84(1):131–148, 2009.

- [BMT18] C. Bandt, D. Mekhontsev, and A. Tetenov. A single fractal pinwheel tile. *Proc. Amer. Math. Soc.*, 146(3):1271–1285, 2018.
- [Fal95] K. J. Falconer. On the Minkowski measurability of fractals. *Proc. Amer. Math. Soc.*, 123(4):1115–1124, 1995.
- [Gat00] D. Gatzouras. Lacunarity of self-similar and stochastically self-similar sets. *Trans. Amer. Math. Soc.*, 352(5):1953–1983, 2000.
- [HL08] X.-G. He and K.-S. Lau. On a generalized dimension of self-affine fractals. *Math. Nachr.*, 281(8):1142–1158, 2008.
- [KK15] M. Kesseböhmer and S. Kombrink. Minkowski content and fractal Euler characteristic for conformal graph directed systems. *J. Fractal Geom.*, 2(2):171–227, 2015.
- [KPW16] S. Kombrink, E. P. J. Pearse, and S. Winter. Lattice-type self-similar sets with pluriphase generators fail to be Minkowski measurable. *Math. Z.*, 283(3-4):1049–1070, 2016.
- [Lap93] M. L. Lapidus. Vibrations of fractal drums, the Riemann hypothesis, waves in fractal media and the Weyl-Berry conjecture. In *Ordinary and partial differential equations, Vol. IV (Dundee, 1992)*, volume 289 of *Pitman Res. Notes Math. Ser.*, pages 126–209. Longman Sci. Tech., Harlow, 1993.
- [LP93] M. L. Lapidus and C. Pomerance. The Riemann zeta-function and the one-dimensional Weyl-Berry conjecture for fractal drums. *Proc. London Math. Soc. (3)*, 66(1):41–69, 1993.
- [LvF00] M. L. Lapidus and M. van Frankenhuysen. *Fractal geometry and number theory*. Birkhäuser Boston, Inc., Boston, MA, 2000. Complex dimensions of fractal strings and zeros of zeta functions.
- [Man95] B. B. Mandelbrot. Measures of fractal lacunarity: Minkowski content and alternatives. In *Fractal geometry and stochastics (Finsterbergen, 1994)*, volume 37 of *Progr. Probab.*, pages 15–42. Birkhäuser, Basel, 1995.
- [PW12] E. P. J. Pearse and S. Winter. Geometry of canonical self-similar tilings. *Rocky Mountain J. Math.*, 42(4):1327–1357, 2012.
- [Sch94] A. Schief. Separation properties for self-similar sets. *Proc. Amer. Math. Soc.*, 122(1):111–115, 1994.
- [Sta76] L. L. Stachó. On the volume function of parallel sets. *Acta Sci. Math. (Szeged)*, 38(3–4):365–374, 1976.
- [Win15] S. Winter. Minkowski content and fractal curvatures of self-similar tilings and generator formulas for self-similar sets. *Adv. Math.*, 274:285–322, 2015.