

On inverse shadowing

Good, Chris; Mitchell, Joel; Thomas, Joe

DOI:

[10.1080/14689367.2020.1739228](https://doi.org/10.1080/14689367.2020.1739228)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Good, C, Mitchell, J & Thomas, J 2020, 'On inverse shadowing', *Dynamical Systems*, vol. 35, no. 3, pp. 539-547.
<https://doi.org/10.1080/14689367.2020.1739228>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is an Accepted Manuscript of an article published by Taylor & Francis in *Dynamical Systems* on 08/03/2020, available online:
<https://doi.org/10.1080/14689367.2020.1739228>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

On inverse shadowing

Chris Good^a, Joel Mitchell^a and Joe Thomas^b

^aSchool of Mathematics, University of Birmingham, Birmingham, B15 2TT, UK;

^bSchool of Mathematics, The University of Manchester, Manchester, M13 9PL, UK
and Joe Thomas^b

ARTICLE HISTORY

Compiled Wednesday 4th March, 2020

ABSTRACT

We give a reformulation of the inverse shadowing property with respect to the class of all pseudo-orbits. This reformulation bears witness to the fact that the property is far stronger than might initially seem. We give some implications of this reformulation, in particular showing that systems with inverse shadowing are not sensitive. Finally we show that, on compact spaces, inverse shadowing is equivalent to a finite version of it.

KEYWORDS

inverse shadowing; pseudo-orbit; minimality; sensitivity; equicontinuity

AMS CLASSIFICATION

37B05, 37C50, 54H20

1. Introduction

Let $f: X \rightarrow X$ be a continuous function on a compact metric space X . A δ -pseudo-orbit is a sequence $(x_i)_{i \in \omega} \subseteq X$ such that $d(f(x_i), x_{i+1}) < \delta$ for all $i \in \omega$. Such sequences arise naturally during the computation of orbits of points in dynamical systems, and so their study is of great importance, for example, in computational models of dynamical systems. Indeed, through calculation one often encounters rounding errors and so the generated sequence of points is not actually a true orbit of the system, but instead a δ -pseudo-orbit with δ dependent upon the degree of accuracy to which one can compute. One question that may then be asked is, to what extent does this sequence reflect any of the original dynamics in the system? Ultimately, this is a question regarding the stability of a dynamical system and one line of enquiry is to determine if such sequences are closely followed by true orbits of the system; thus leading directly to the notion of shadowing. The sequence $(y_i)_{i \in \omega} \subseteq X$ is said to ε -shadow the sequence $(x_i)_{i \in \omega}$ provided that $d(y_i, x_i) < \varepsilon$ for all $i \in \omega$. If $(y_i)_{i \in \omega} = (f^i(y))_{i \in \omega}$ for some $y \in X$, then this shadowing means that to some degree of accuracy $\varepsilon > 0$, the pseudo-orbit is followed by a true orbit. If this happens for all pseudo-orbits of some

Email: c.good@bham.ac.uk (C. Good)

Email: JSM140@bham.ac.uk (J. Mitchell)

Email: joe.thomas-3@postgrad.manchester.ac.uk (J. Thomas)

given accuracy, then the system (X, f) is said to have the shadowing property.

This concept of shadowing has a natural interpretation when modelling a system numerically and this has been studied in detail in the works of Corless [9], Palmer [28] and Pearson [30]. It is also an important theoretical concept. For example, Bowen [6] used shadowing implicitly as a key step in his proof that the nonwandering set of an Axiom A diffeomorphism is a factor of a shift of finite type. Since then it has been studied extensively as a key factor in stability theory [32, 34, 37], in understanding the structure of ω -limit sets and Julia sets [2, 3, 4, 5, 7, 17, 25], and as a property in and of itself [11, 13, 15, 16, 23, 26, 27, 31, 32, 36].

In this paper, we look at the related concept, inverse shadowing. First introduced by Corless and Pilyugin [10] as something akin to the “dual” of shadowing, and as part of the concept of bishadowing by Diamond *et al* [12], inverse shadowing in a system informally means that true trajectories may be recovered from computed orbits (within some given accuracy). Kloeden, Ombach and Pokorskii [20] later defined inverse shadowing using the notion of a δ -method which are functions mapping points to δ -pseudo-orbits originating from the point. This allows one to consider certain classes of maps from the space to the space of pseudo-orbits through the imposition of extra structure on such mappings such as continuity.

Such classes have been studied in a variety of different settings for example [18, 19, 20, 22, 29, 33]. Of particular interest has been its relationship to structural stability. In [33], Pilyugin showed that if an Axiom A diffeomorphism on a closed C^∞ manifold is structurally stable, then it has the inverse shadowing property with respect to classes of continuous methods. Meanwhile in [21], Kloeden and Ombach prove that a structurally stable homeomorphism on a compact space has inverse shadowing with respect to the class of methods induced by homeomorphism. Further results in this direction can be found in [8, 19, 22]. We remark that inverse shadowing, particularly with regard to the class of methods induced by homeomorphism, is closely related to Lewowicz’s notion of persistency [24] (see also [35]).

Within this paper, we examine inverse shadowing with respect to the class of all pseudo-orbits. In Section 3 we prove that inverse shadowing is equivalent to a property which essentially involves a quantifier swap in the definition. We use this reformulation to give a number of implications in Section 4, in particular showing that a system exhibiting inverse shadowing is not *eventually sensitive* (defined below). We conclude by showing that when the phase space is compact, for the classes \mathcal{T}_0 , \mathcal{T}_c and \mathcal{T}_h , inverse shadowing is equivalent to a finite version.

Although many of our results hold for metric spaces in general¹, we restrict our attention to compact metric spaces since inverse shadowing is a *dynamical property*, i.e. it is invariant under conjugacy, in this case. Thus (X, d) is a compact metric space throughout this paper. We denote by \mathbb{Z} the set of all integers; the set of positive integers $1, 2, 3, 4, \dots$ is denoted by \mathbb{N} whilst $\omega := \mathbb{N} \cup \{0\}$.

¹Theorems 3.1, 3.3, 4.1 and 4.2 together with Corollary 4.3 and Lemma 4.5 do not require the assumption of compactness.

2. Preliminaries

We firstly outline the notions of inverse shadowing that will be used here. Let $f: X \rightarrow X$ be a homeomorphism (resp. a continuous function) on a compact metric space (X, d) . We call the pair (X, f) a *dynamical system*. The *orbit* of x under f is the set of points $\{f^i(x)\}_{i \in A}$ and is denoted by $\text{Orb}_f(x)$, with the understanding that $A = \mathbb{Z}$ if f is a homeomorphism and a full version of inverse shadowing is under consideration whilst $A = \omega$ when it is a positive version of inverse shadowing under consideration. For $\delta > 0$, we refer to a bi-infinite sequence $(x_k)_{k \in \mathbb{Z}}$ such that $d(f(x_k), x_{k+1}) < \delta$ for all $k \in \mathbb{Z}$ as a δ -pseudo-orbit; and a mono-infinite sequence $(x_k)_{k \in \omega}$ such that $d(f(x_k), x_{k+1}) < \delta$ for all $k \in \omega$ as a *positive* δ -pseudo-orbit.

Let $X^{\mathbb{Z}}$ (resp. X^{ω}) be the product space of all bi-infinite (resp. mono-infinite) sequences, with the product topology (note that compactness of X implies compactness of the product). Then for any given $\delta > 0$, let $\Phi_f(\delta) \subseteq X^{\mathbb{Z}}$ be the set of all δ -pseudo-orbits with respect to f (resp. $\Phi_f^+(\delta) \subseteq X^{\omega}$ the set of all positive δ -pseudo-orbits with respect to f). A mapping $\varphi: X \rightarrow \Phi_f(\delta)$ (resp. $\varphi: X \rightarrow \Phi_f^+(\delta)$), such that, for each $x \in X$, $\varphi(x)_0 = x$, is called a (resp. positive) δ -method for f , where $\varphi(x)_k$ is used to denote the k^{th} term in the sequence $\varphi(x)$. We denote by $\mathcal{T}_0(f, \delta)$ the set of all respective δ -methods with understanding from the context whether this refers to positive methods or not. Similarly we denote by $\mathcal{T}_c(f, \delta)$ the set of all continuous (positive) δ -methods, and by $\mathcal{T}_h(f, \delta)$ the set of all (positive) δ -methods induced by a homeomorphism, i.e., $\mathcal{T}_h(f, \delta)$ is the set of (resp. positive) δ -methods φ for which there exists a homeomorphism $h: X \rightarrow X$ such that $d(f(x), h(x)) < \delta$ for each $x \in X$ and $\varphi(x)_k = h^k(x)$, for all relevant $k \in \mathbb{Z}$ (resp. $k \in \omega$).

Definition 2.1. Let $f: X \rightarrow X$ be a homeomorphism (resp. continuous function). We say that f experiences (*resp. positive*) \mathcal{T}_α -inverse shadowing if, for any $\varepsilon > 0$ there exists $\delta > 0$ such that for any $x \in X$ and any $\varphi \in \mathcal{T}_\alpha(f, \delta)$ there exists $y \in X$ such that $\varphi(y)$ ε -shadows x ; i.e.

$$\forall k \in \mathbb{Z} \text{ (resp. } k \in \omega), d(\varphi(y)_k, f^k(x)) < \varepsilon.$$

Definition 2.2. Let $f: X \rightarrow X$ be a homeomorphism (resp. continuous function). We say that f experiences (*resp. positive*) weak inverse shadowing with respect to the class \mathcal{T}_α if, for any $\varepsilon > 0$ there exists $\delta > 0$ such that for any $x \in X$ and any $\varphi \in \mathcal{T}_\alpha(f, \delta)$ there exists $y \in X$ such that

$$\varphi(y) \subseteq B_\varepsilon(\text{Orb}_f(x)).$$

(NB. As stated in the preliminaries, $\text{Orb}_f(x)$ is only the positive trajectory of x when considering the ‘positive’ version of the above statement).

Remark 1. Clearly if a system has \mathcal{T}_α -inverse shadowing then it has weak inverse shadowing with respect to the class \mathcal{T}_α .

3. An equivalent reformulation of inverse shadowing

Within the literature surrounding inverse shadowing, several authors [10, 22] have commented on the importance of restricting one's attention to certain admissible classes of pseudo-orbits. Theorem 3.1 brings to light exactly why such a restriction may be important; in particular, it demonstrates that \mathcal{T}_0 -inverse shadowing is a much stronger property than it may appear *prima facie*.

Theorem 3.1. *A system (X, f) with f a homeomorphism (resp. continuous map) has (resp. positive) \mathcal{T}_0 -inverse shadowing if and only if for any $\varepsilon > 0$ there exists $\delta > 0$ such that for any $x \in X$ there exists $y \in X$ such that for any $\varphi \in \mathcal{T}_0(f, \delta)$ and any $k \in \mathbb{Z}$ (resp. $k \in \omega$) $d(f^k(x), \varphi(y)_k) < \varepsilon$.*

Proof. Clearly the latter implies the former. Thus, suppose that f has (resp. positive) \mathcal{T}_0 -inverse shadowing and assume further that the latter is false, that is, there exists an $\varepsilon > 0$ such that

$$\forall \delta > 0 \exists x \in X : \forall y \in X \exists \varphi \in \mathcal{T}_0(f, \delta) \exists k \in \mathbb{Z} \text{ (resp. } k \in \omega) : d(f^k(x), \varphi(y)_k) \geq \varepsilon. \quad (1)$$

Take such an ε and let $\delta > 0$ correspond to this ε in the definition of inverse shadowing. Now, fix $x \in X$ corresponding to this ε and δ as in (1). Then by (1), for each $y \in X$, there will exist a $\varphi_y \in \mathcal{T}_0(f, \delta)$ and a $k \in \mathbb{Z}$ (resp. $k \in \omega$) for which $d(f^k(x), \varphi_y(y)_k) \geq \varepsilon$. Now, define a map $\varphi : X \rightarrow X^{\mathbb{Z}} : y \mapsto \varphi_y(y)$. By construction, $\varphi \in \mathcal{T}_0(f, \delta)$ and by (resp. positive) \mathcal{T}_0 -inverse shadowing, there will exist $y \in X$ such that $d(f^k(x), \varphi(y)_k) < \varepsilon$ for all $k \in \mathbb{Z}$ (resp. $k \in \omega$). But by the construction of φ , this can not be the case for all such k and thus one obtains a contradiction. \square

The compactness of X ensures the uniform continuity of f which we use to obtain the following corollary.

Corollary 3.2. *A system (X, f) with f a homeomorphism (resp. continuous map) has (resp. positive) \mathcal{T}_0 -inverse shadowing if and only if for any $\varepsilon > 0$ there exists $\delta > 0$ such that for any $x \in X$ there exists $y \in X$ such that for any $\varphi \in \mathcal{T}_0(f, \delta)$ and any $k \in \mathbb{Z}$ (resp. $k \in \omega$) and any $z \in B_\delta(y)$, $d(f^k(x), \varphi(z)_k) < \varepsilon$.*

Proof. That the latter entails the former is trivial. Therefore, suppose the former. Let $\varepsilon > 0$ be given. Then, by Theorem 3.1 there exists $\eta > 0$ (without loss of generality $\eta < \frac{\varepsilon}{2}$) such that for any $x \in X$, there exists $y \in X$ such that for any $\varphi \in \mathcal{T}_0(f, \eta)$ and any $k \in \mathbb{Z}$ (resp. $k \in \omega$) $d(f^k(x), \varphi(y)_k) < \frac{\varepsilon}{2}$.

First consider the case when f is a homeomorphism. Using uniform continuity, take $\delta > 0$ such that, for any $a, b \in X$, if $d(a, b) < \delta$ then $d(f(a), f(b)) < \frac{\eta}{2}$ and $d(f^{-1}(a), f^{-1}(b)) < \frac{\eta}{2}$; without loss of generality $\delta < \frac{\eta}{2}$. Now let $(z_k)_{k \in \mathbb{Z}}$ be a δ -pseudo-orbit where $z_0 \in B_\delta(y)$. Then, by the triangle inequality,

$$(\dots, z_{-k}, \dots, z_{-1}, y, z_1, \dots, z_k, \dots)$$

is a η -pseudo-orbit through y . It follows by the choice of y that $d(f^k(x), z_k) < \frac{\varepsilon}{2}$ for all $k \in \mathbb{Z} \setminus \{0\}$. Since $d(x, z_0) \leq d(x, y) + d(y, z_0) = \frac{\varepsilon}{2} + \eta < \varepsilon$, the sequence $(z_k)_{k \in \mathbb{Z}}$ ε -shadows x and we are done.

Now suppose that f is a continuous map; we are now considering positive \mathcal{T}_0 -inverse shadowing. We argue similarly to before. Using uniform continuity, take $\delta > 0$ such that, for any $a, b \in X$, if $d(a, b) < \delta$ then $d(f(a), f(b)) < \frac{\eta}{2}$. Without loss of generality $\delta < \frac{\eta}{2}$. Now let $(z_k)_{k \in \omega}$ be a δ -pseudo-orbit where $z_0 \in B_\delta(y)$. Then, by the triangle inequality, $(y, z_1, \dots, z_k, \dots)$ is a η -pseudo-orbit through y . It follows by the choice of y that $d(f^k(x), z_k) < \frac{\varepsilon}{2}$ for all $k \in \omega \setminus \{0\}$. Since

$$d(x, z_0) \leq d(x, y) + d(y, z_0) < \frac{\varepsilon}{2} + \delta < \frac{\varepsilon}{2} + \eta < \varepsilon,$$

the sequence $(z_k)_{k \in \omega}$ ε -shadows x and we are done. \square

Using the same technique as in the proof of Theorem 3.1 we come by the following reformulation of weak inverse shadowing. We omit the proof.

Theorem 3.3. *A map $f: X \rightarrow X$ has (resp. positive) weak inverse shadowing with respect to the class \mathcal{T}_0 if and only if for any $\varepsilon > 0$ there exists $\delta > 0$ such that for any $x \in X$ there exists $y \in X$ such that for any $\varphi \in \mathcal{T}_0(f, \delta)$,*

$$\varphi(y) \subseteq B_\varepsilon(\text{Orb}_f(x)).$$

4. Implications

Before giving some implications of Theorem 3.1, we will need to recall some standard definitions in dynamical systems. A dynamical system (X, f) is said to be *equicontinuous* at a point $x \in X$ if for any $\varepsilon > 0$ there exists $\delta > 0$ such that, for any $y \in X$, if $d(x, y) < \delta$ then, for any $n \in \omega$, $d(f^n(x), f^n(y)) < \varepsilon$. The system itself is said to be *equicontinuous* if it is equicontinuous at every point. We observe that, since X is compact, equicontinuity is equivalent to *uniform equicontinuity* (δ is chosen independently of the point $x \in X$). A dynamical system exhibits *sensitive dependence on initial conditions* (or is *sensitive*) if there exists $\delta > 0$ such that for any nonempty open set U there exist $x, y \in U$ and $k \in \mathbb{N}$ such that $d(f^k(x), f^k(y)) \geq \delta$. Such a δ is referred to as a *sensitivity constant* for (X, f) . A weakening of sensitivity was introduced in [14]; a dynamical system (X, f) is *eventually sensitive* if there exists $\delta > 0$ such that for any $x \in X$ and any $\varepsilon > 0$ there exist $n, k \in \omega$ and $y \in B_\varepsilon(f^n(x))$ such that $d(f^{n+k}(x), f^k(y)) \geq \delta$. We refer to such a δ as an *eventual-sensitivity constant*. Clearly sensitivity implies eventual sensitivity but, as demonstrated in [14], the converse is not true. It is also easy to see that neither sensitivity nor eventual sensitivity can be held in conjunction with equicontinuity. Finally a system is *expansive* if there exists $\delta > 0$ such that for any $x, y \in X$ there exists $k \in \omega$ with $d(f^k(x), f^k(y)) \geq \delta$. It is an easy exercise to show that if a system is *perfect*, that is the space has no isolated points, then an expansive system is sensitive.

In [19, Theorem 4] the authors show a homeomorphism with \mathcal{T}_0 -inverse shadowing is not expansive. Whilst not explicitly stated there, we remark that this result assumes that for any ε there exist distinct points $x, y \in X$ with $d(x, y) < \varepsilon$. Indeed, it is easy to see that a system consisting of a single periodic orbit has \mathcal{T}_0 -inverse shadowing but is also expansive. The reformulation given by Theorem 3.1 enables us to give the following much stronger result.

Theorem 4.1. *Let $f: X \rightarrow X$ be a continuous map. If f has the positive \mathcal{T}_0 -inverse shadowing property then the system (X, f) is not eventually sensitive. (If f is a homeomorphism with \mathcal{T}_0 -inverse shadowing, then neither (X, f) nor the inverse system (X, f^{-1}) are eventually sensitive.)*

Proof. Suppose that f is eventually sensitive and let $\delta_0 > 0$ be a constant of eventual sensitivity. Pick $\varepsilon > 0$ with $\varepsilon < \frac{\delta_0}{2}$. Now take a $\delta > 0$ corresponding to this ε as in the reformulation of positive inverse shadowing in Theorem 3.1. Pick $x \in X$. Then there exists $y \in X$ such that every δ -pseudo-orbit through y ε -shadows x . Consider $f(y)$: By eventual sensitivity there exist $n, k \in \mathbb{N}$ and $z \in B_\delta(f^{n+1}(y))$ such that $d(f^k(z), f^{n+k+1}(y)) \geq \delta_0$. But the sequence

$$(y, f(y), \dots, f^n(y), z, f(z), \dots, f^k(z), f^{k+1}(z), \dots),$$

is a δ -pseudo orbit from y . Similarly so is the orbit sequence of y . Therefore, by inverse shadowing, $f^{n+k+1}(y), f^k(z) \in B_\varepsilon(f^{n+k+1}(x))$. It follows by the triangle inequality that

$$d(f^{n+k+1}(y), f^k(z)) < 2\varepsilon < \delta_0,$$

which is a contradiction. Therefore f is not eventually sensitive. \square

Remark 2. Note that there is no need to make the assumption that for any $\varepsilon > 0$ there are points ε -close in Theorem 4.1.

In [19, Theorem 3] the authors show that a chain transitive homeomorphism on a compact metric space is minimal if and only if it has the weak inverse shadowing property with respect to \mathcal{T}_0 . With Theorem 3.3 this result, as well as the following analogous one, becomes much more elementary. (NB. In the statement of [19, Theorem 3] the authors assume the phase space is compact, this assumption does not appear necessary for their proof nor ours.) Recall first that a system (X, f) is said to be minimal if, for $A \subseteq X$ closed, $f(A) = A$ implies that $A = X$ or $A = \emptyset$. Equivalently it is minimal if $\overline{\text{Orb}_f(x)} = X$ for all $x \in X$.

Theorem 4.2 (see [19]). *Let $f: X \rightarrow X$ be a chain transitive continuous function. Then f is minimal if and only if f has positive weak inverse shadowing with respect to the class \mathcal{T}_0 .*

Proof. Since the orbit of every point is dense under a minimal map, it is trivial that such a map has positive weak inverse shadowing with respect to the class \mathcal{T}_0 .

Pick $z \in X$. Let $\varepsilon > 0$ and take $\delta > 0$ as in Theorem 3.3. Then there exists $y \in X$ such that for any $\varphi \in \mathcal{T}_0(f, \delta)$,

$$\varphi(y) \subseteq B_\varepsilon(\text{Orb}(z)).$$

Since f is chain transitive there is a δ -chain from y to every point in X . It follows then by the above that $x \in B_\varepsilon(\text{Orb}(z))$ for all $x \in X$. Since $\varepsilon > 0$ was arbitrary the result follows. \square

Corollary 4.3. *Let $f: X \rightarrow X$ be a chain transitive continuous function. If f has the positive \mathcal{T}_0 -inverse shadowing property then the system (X, f) is equicontinuous. (If f is a homeomorphism with \mathcal{T}_0 -inverse shadowing, then both (X, f) and the inverse system (X, f^{-1}) are equicontinuous.)*

Proof. By Theorem 4.2 the system is minimal. A minimal system is either equicontinuous or sensitive (see² [1, Corollary 2]). \square

We conclude this paper by showing that inverse shadowing with respect to any given class, is equivalent to what we call *finite inverse shadowing*, with respect to the same class. This is akin to the result that shadowing is equivalent to finite shadowing [32].

Definition 4.4. A system (X, f) exhibits *finite (resp. finite positive) \mathcal{T}_α -inverse shadowing* if for any $\varepsilon > 0$ there exists $\delta > 0$ such that for any $x \in X$ and any $\varphi \in \mathcal{T}_0(f, \delta)$ and any $n \in \mathbb{N}$ there exists $y_n \in X$ such that for any $k \in \{-n, \dots, 0, \dots, n\}$ (resp. $k \in \{0, \dots, n\}$) $d(f^k(x), \varphi(y_n)_k) < \varepsilon$.

The proof of the lemma below is very similar to that of Theorem 3.1 and is therefore omitted.

Lemma 4.5. *A map $f: X \rightarrow X$ has finite (resp. finite positive) \mathcal{T}_0 -inverse shadowing if and only if for any $\varepsilon > 0$ there exists $\delta > 0$ such that for any $x \in X$ and any $n \in \mathbb{N}$ there exists $y_n \in X$ such that for any $\varphi \in \mathcal{T}_0(f, \delta)$ and any $k \in \{-n, \dots, 0, \dots, n\}$ (resp. $k \in \{0, \dots, n\}$) $d(f^k(x), \varphi(y_n)_k) < \varepsilon$.*

Proposition 4.6. *For any $\alpha \in \{0, c, h\}$, a homeomorphism (resp. continuous map) $f: X \rightarrow X$ has finite (positive) \mathcal{T}_α -inverse shadowing if and only if it has (positive) \mathcal{T}_α -inverse shadowing.*

Proof. We will prove the cases when f is a homeomorphism, with reference to \mathcal{T}_α -inverse shadowing. The cases when f is a continuous map, with reference to positive \mathcal{T}_α -inverse shadowing, are similar.

Clearly, for any $\alpha \in \{0, c, h\}$, if f has it has (positive) \mathcal{T}_α -inverse shadowing then it has finite (positive) \mathcal{T}_α -inverse shadowing.

Suppose that $f: X \rightarrow X$ has finite \mathcal{T}_c -inverse shadowing. Let $\varepsilon > 0$ be given and let $\delta > 0$ correspond to $\frac{\varepsilon}{2}$ in finite inverse shadowing. Let $x \in X$ and take $\varphi \in \mathcal{T}_c(f, \delta)$. For any $n \in \mathbb{N}$, there exists $y_n \in X$ such that $d(\varphi(y_n)_i, f^i(x)) < \frac{\varepsilon}{2}$ for all $i \in \{-n, \dots, 0, \dots, n\}$. By sequential compactness, (y_n) has a convergent subsequence; call it (y_n) again. Let $y = \lim_{n \rightarrow \infty} y_n$; $y \in \overline{B_{\frac{\varepsilon}{2}}(x)}$. Since $\varphi: X \rightarrow X^{\mathbb{Z}}$ is continuous,

$$\lim_{n \rightarrow \infty} \varphi(y_n) = \varphi \left(\lim_{n \rightarrow \infty} y_n \right) = \varphi(y).$$

Moreover it follows that, for any $k \in \mathbb{N}$ there exists $M \in \mathbb{N}$ such that for any $m > M$ we have $d(\varphi(y)_i, \varphi(y_m)_i) < \frac{\varepsilon}{2}$ for all $i \in \{-k, \dots, 0, \dots, k\}$. For a fixed k and corresponding M , we must also have, for all $m > M$ that, $d(\varphi(y_m)_i, f^i(x)) < \frac{\varepsilon}{2}$ for all $i \in \{-k, \dots, 0, \dots, k\}$. Then, by the triangle inequality,

²The authors of [1] prove that compact metric minimal systems are either uniformly equicontinuous or sensitive. Without the presence of compactness it is still an easy exercise to show that the system is either equicontinuous or sensitive.

$$\forall k \in \mathbb{N} \exists M \in \mathbb{N} : \forall m > M \forall i \in \{-k, \dots, 0, \dots, k\}, d(\varphi(y)_i, f^i(x)) < \varepsilon.$$

From this it follows that for every $k \in \mathbb{N}$ we have $d(\varphi(y)_k, f^k(x)) < \varepsilon$. Hence $f: X \rightarrow X$ has \mathcal{T}_c -inverse shadowing. The proof for the class \mathcal{T}_h is similar to the above (as are the positive versions for the classes \mathcal{T}_c and \mathcal{T}_h); these are therefore omitted.

Now suppose that $f: X \rightarrow X$ has finite \mathcal{T}_0 -inverse shadowing. Let $\varepsilon > 0$ be given and let $2\delta > 0$ correspond to $\frac{\varepsilon}{2}$ in finite inverse shadowing; without loss of generality taking $\delta < \frac{\varepsilon}{4}$. Then, by Lemma 3.1, for any $n \in \mathbb{N}$ there exists $y_n \in X$ such that for any $\varphi \in \mathcal{T}_0(f, 2\delta)$, and any $i \in \{-n, \dots, 0, \dots, n\}$, $d(\varphi(y_n)_i, f^i(x)) < \frac{\varepsilon}{2}$. By sequential compactness, (y_n) has a convergent subsequence; call it (y_n) again. Let $y = \lim_{n \rightarrow \infty} y_n$; $y \in \overline{B_{\frac{\varepsilon}{2}}(x)}$. Because f and f^{-1} are continuous, there exists $N \in \mathbb{N}$ such that $d(f(y_n), f(y)) < \delta$ and $d(f^{-1}(y_n), f^{-1}(y)) < \delta$ for all $n > N$. Then for any $n > N$ we have

$$B_\delta(f(y)) \subseteq B_{2\delta}(f(y_n)) \subseteq B_{\frac{\varepsilon}{2}}(f(x)),$$

and

$$B_\delta(f^{-1}(y)) \subseteq B_{2\delta}(f^{-1}(y_n)) \subseteq B_{\frac{\varepsilon}{2}}(f^{-1}(x)).$$

Now let $(z_k)_{k \in \mathbb{Z}}$ be a δ -pseudo-orbit through y (so $z_0 = y$), we will show this ε -shadows x . Fix $n > N$. Then

$$(\dots, \dots, z_{-k}, \dots, z_{-2}, z_{-1}, y_n, z_1, z_2, \dots, z_k, \dots),$$

is a 2δ -pseudo-orbit through y_n ; by finite shadowing $d(f^i(x), z_i) \leq \frac{\varepsilon}{2}$ for all $i \in \{-n, \dots, -1, 1, \dots, n\}$. Additionally, $d(x, z_0) \leq \frac{\varepsilon}{2} < \varepsilon$, as $z_0 = y$. Since $n > N$ was arbitrary it follows that, for any $n > N$, $d(f^i(x), z_i) \leq \frac{\varepsilon}{2}$ for all $i \in \{-n, \dots, -1, 1, \dots, n\}$. This in turn entails that $d(f^i(x), z_i) \leq \frac{\varepsilon}{2}$ for all $i \in \mathbb{Z}$. As this was an arbitrary δ -pseudo-orbit through y we are done. \square

References

- [1] Joseph Auslander and James A. Yorke. Interval maps, factors of maps, and chaos. *Tôhoku Math. J. (2)*, 32(2):177–188, 1980.
- [2] Andrew D. Barwell, Gareth Davies, and Chris Good. On the ω -limit sets of tent maps. *Fund. Math.*, 217(1):35–54, 2012.
- [3] Andrew D. Barwell, Chris Good, Piotr Oprocha, and Brian E. Raines. Characterizations of ω -limit sets in topologically hyperbolic systems. *Discrete Contin. Dyn. Syst.*, 33(5):1819–1833, 2013.
- [4] Andrew D. Barwell, Jonathan Meddaugh, and Brian E. Raines. Shadowing and ω -limit sets of circular Julia sets. *Ergodic Theory Dynam. Systems*, 35(4):1045–1055, 2015.

- [5] Andrew D. Barwell and Brian E. Raines. The ω -limit sets of quadratic Julia sets. *Ergodic Theory Dynam. Systems*, 35(2):337–358, 2015.
- [6] Rufus Bowen. Markov partitions for Axiom A diffeomorphisms. *Amer. J. Math.*, 92:725–747, 1970.
- [7] Rufus Bowen. ω -limit sets for axiom A diffeomorphisms. *J. Differential Equations*, 18(2):333–339, 1975.
- [8] Taeyoung Choi, Keonhee Lee, and Yong Zhang. Characterisations of Ω -stability and structural stability via inverse shadowing. *Bull. Austral. Math. Soc.*, 74(2):185–196, 2006.
- [9] Robert M. Corless. Defect-controlled numerical methods and shadowing for chaotic differential equations. *Phys. D*, 60(1-4):323–334, 1992. Experimental mathematics: computational issues in nonlinear science (Los Alamos, NM, 1991).
- [10] Robert M. Corless and S. Yu. Pilyugin. Approximate and real trajectories for generic dynamical systems. *J. Math. Anal. Appl.*, 189(2):409–423, 1995.
- [11] Ethan M. Coven, Ittai Kan, and James A. Yorke. Pseudo-orbit shadowing in the family of tent maps. *Trans. Amer. Math. Soc.*, 308(1):227–241, 1988.
- [12] Phil Diamond, Victor Kozyakin, Peter Kloeden, and Alexei Pokrovskii. Computer robustness of semi-hyperbolic mappings. *Random Comput. Dynam.*, 3(1-2):57–70, 1995.
- [13] Leobardo Fernández and Chris Good. Shadowing for induced maps of hyperspaces. *Fund. Math.*, 235(3):277–286, 2016.
- [14] Chris Good, Robert Leek, and Joel Mitchell. Equicontinuity, transitivity and sensitivity: The Auslander-Yorke dichotomy revisited. *Discrete Contin. Dyn. Syst.*, 40(4):2441–2474, 2020.
- [15] Chris Good, Sergio Macías, Jonathan Meddaugh, Joel Mitchell, and Joe Thomas. Expansivity and unique shadowing, 2020.
- [16] Chris Good and Jonathan Meddaugh. Shifts of finite type as fundamental objects in the theory of shadowing. *Inventiones mathematicae*, Dec 2019.
- [17] Chris Good, Jonathan Meddaugh, and Joel Mitchell. Shadowing, internal chain transitivity and α -limit sets, 2019.
- [18] Chris Good, Joel Mitchell, and Joe Thomas. Preservation of shadowing in discrete dynamical systems. *J. Math. Anal. Appl.*, 485(1):123767, 39, 2020.
- [19] B. Honary and Alireza Zamani Bahabadi. Inverse shadowing and weak inverse shadowing property. *Appl. Math. (Irvine)*, 3(5):478–483, 2012.
- [20] P. Kloeden, J. Ombach, and A. Pokrovskii. Continuous and inverse shadowing. *Funct. Differ. Equ.*, 6(1-2):137–153, 1999.
- [21] P. E. Kloeden and J. Ombach. Hyperbolic homeomorphisms and bishadowing. *Ann. Polon. Math.*, 65(2):171–177, 1997.
- [22] Keonhee Lee. Continuous inverse shadowing and hyperbolicity. *Bull. Austral. Math. Soc.*, 67(1):15–26, 2003.
- [23] Keonhee Lee and Kazuhiro Sakai. Various shadowing properties and their equivalence. *Discrete Contin. Dyn. Syst.*, 13(2):533–540, 2005.
- [24] Jorge Lewowicz. Persistence in expansive systems. *Ergodic Theory Dynam. Systems*, 3(4):567–578, 1983.
- [25] Jonathan Meddaugh and Brian E. Raines. Shadowing and internal chain transitivity. *Fund. Math.*, 222(3):279–287, 2013.
- [26] Joel Mitchell. Orbital shadowing, ω -limit sets and minimality. *Topology Appl.*, 268:106903, 7, 2019.
- [27] Helena E. Nusse and James A. Yorke. Is every approximate trajectory of

- some process near an exact trajectory of a nearby process? *Comm. Math. Phys.*, 114(3):363–379, 1988.
- [28] Kenneth James Palmer. *Shadowing in dynamical systems: theory and applications*, volume 501. Springer Science & Business Media, 2013.
 - [29] Jong-Jin Park and Keonhee Lee. Inverse shadowing of circle maps. *Bull. Austral. Math. Soc.*, 69(3):353–359, 2004.
 - [30] D. W. Pearson. Shadowing and prediction of dynamical systems. *Math. Comput. Modelling*, 34(7-8):813–820, 2001.
 - [31] Timothy Pennings and Jeffrey Van Eeuwen. Pseudo-orbit shadowing on the unit interval. *Real Anal. Exchange*, 16(1):238–244, 1990/91.
 - [32] Sergei Yu. Pilyugin. *Shadowing in dynamical systems*, volume 1706 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1999.
 - [33] Sergei Yu. Pilyugin. Inverse shadowing by continuous methods. *Discrete Contin. Dyn. Syst.*, 8(1):29–38, 2002.
 - [34] Clark Robinson. Stability theorems and hyperbolicity in dynamical systems. In *Proceedings of the Regional Conference on the Application of Topological Methods in Differential Equations (Boulder, Colo., 1976)*, volume 7, pages 425–437, 1977.
 - [35] Kazuhiro Sakai. Diffeomorphisms with persistency. *Proc. Amer. Math. Soc.*, 124(7):2249–2254, 1996.
 - [36] Kazuhiro Sakai. Various shadowing properties for positively expansive maps. *Topology Appl.*, 131(1):15–31, 2003.
 - [37] Peter Walters. On the pseudo-orbit tracing property and its relationship to stability. In *The structure of attractors in dynamical systems (Proc. Conf., North Dakota State Univ., Fargo, N.D., 1977)*, volume 668 of *Lecture Notes in Math.*, pages 231–244. Springer, Berlin, 1978.