

SARS-CoV-2 pandemic and repercussions for male infertility patients

Esteves, Sandro C.; Kirkman-Brown, Jackson

DOI:

[10.1111/andr.12809](https://doi.org/10.1111/andr.12809)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Esteves, SC & Kirkman-Brown, J 2020, 'SARS-CoV-2 pandemic and repercussions for male infertility patients: a proposal for the individualized provision of andrological services', *Andrology*. <https://doi.org/10.1111/andr.12809>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is the peer reviewed version of the following article: Esteves, S.C., Lombardo, F., Garrido, N., Alvarez, J., Zini, A., Colpi, G.M., Kirkman-Brown, J., Lewis, S.E., Björndahl, L., Majzoub, A., Cho, C.-L., Vendeira, P., Hallak, J., Amar, E., Cocuzza, M., Bento, F.C., Figueira, R.C., Sciorio, R., Laursen, R.J., Metwalley, A.M., Jindal, S.K., Parekattil, S., Ramasamy, R., Alviggi, C., Humaidan, P., Yovich, J.L. and Agarwal, A. (2020), SARS-CoV-2 pandemic and repercussions for male infertility patients: a proposal for the individualized provision of andrological services. *Andrologia*. Accepted Author Manuscript. doi:10.1111/andr.12809 , which has been published in final form at <https://doi.org/10.1111/andr.12809>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

SARS-CoV-2 pandemic and repercussions for male infertility patients: a proposal for the individualized provision of andrological services

Journal:	<i>Andrology</i>
Manuscript ID	ANDR-2020-0206.R1
Manuscript Type:	Opinion Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Esteves, Sandro; Androfert, Lombardo, Francesco; 1st University of Rome La Sapienza Rome, University Laboratory of Seminology and Immunology of Reproduction Garrido, Nicolás ; IVI Foundation, Health Research Institute La Fe alvarez, Juan; Centro de Infertilidad Masculina Zini, Armand; McGill University, Colpi, G. M.; Univ Milan, Kirkman-Brown, Jackson; University of Birmingham College of Medical and Dental Sciences, Center for Human Reproduction Lewis, Sheena; Queens University Belfast, Medicine Björndahl, Lars; Karolinska University Hospital and Karolinska Institutet, Centre for Andrology and Sexual Medicine Majzoub, Ahmad; Hamad Medical Corporation Department of Medical Education, Urology Cho, Chak Lam; Kwong Wah Hospital, Surgery (Urology) Vendeira, Pedro; Clínica do Dragão, Andrology Unit Hallak, Jorge; University of Sao Paulo, Dept of Urology Amar, Edouard ; American Hospital of Paris, Cabinet D'Andrologie Cocuzza, Marcello; University of Sao Paulo brazil, Urology Bento, Fabiola ; Androfert Male Reproductive Reference Centre Figueira, Rita; Androfert Male Reproductive Reference Centre Sciorio, Romualdo; Royal Infirmary of Edinburgh, Edinburgh Assisted Conception Programme Laursen, Rita ; Skive Regional Hospital, Fertility Clinic Metwalley, Ahmad; Al Saad Specialist Hospitals, IVF Laboratory Jindal, Sunil; Jindal Hospital, Division of Reproductive Medicine and Andrology Parekattil , Sijo; University of Central Florida College of Medicine Ramasamy, Ranjith; University of Miami Miller School of Medicine Alviggi, Carlo; University of Naples Federico II, Department of Neuroscience, Reproductive Science and Odontostomatology Humaidan, Peter; The fertility Clinic, Viborg Hospital (Skive) Yovich, John; PIVET Medical Centre, Reproductive Medicine Agarwal, Ashok; Case Western Reserve University, Cleveland Clinic Lerner College of Medicine; The Cleveland Clinic Foundation, Urology</p>
Major Area:	Clinical
Clinical:	Urology, Assisted Reproduction, Laboratory Methodologies

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Basic Science:	
Specific Topics:	Semen analyses, Spermatozoa, Reproductive tract



OPINION

SARS-CoV-2 pandemic and repercussions for male infertility patients: a proposal for the individualized provision of andrological services

Authors and Institutions:

§Sandro C. Esteves^{1,2,3}, Francesco Lombardo⁴, Nicolás Garrido⁵, Juan Alvarez⁶, Armand Zini⁷, Giovanni M. Colpi^{8,9}, Jackson Kirkman-Brown^{10,11}, Sheena EM Lewis^{12,13}, Lars Björndahl¹⁴, Ahmad Majzoub^{15,16}, Chak-Lam Cho^{17,18}, Pedro Vendeira¹⁹, Jorge Hallak^{20,21,22}, Edouard Amar²³, Marcello Cocuzza²⁴, Fabiola C. Bento¹, Rita C. Figueira¹, Romualdo Sciorio²⁵, Rita J. Laursen²⁶, Ahmad M. Metwalley^{27,28}, Sunil K. Jindal²⁹, Sijo Parekattil³⁰, Ranjith Ramasamy³¹, Carlo Alviggi³², Peter Humaidan^{3,26}, John L. Yovich^{33,34,35}, Ashok Agarwal³⁶.

¹ANDROFERT, Andrology and Human Reproduction Clinic, Campinas, Brazil
(s.esteves@androfert.com.br; fabiola.bento@androfert.com.br; rita.labfiv@androfert.com.br)

²Department of Surgery (Division of Urology), University of Campinas (UNICAMP), Campinas, Brazil

³Faculty of Health, Aarhus University, Aarhus, Denmark (peter.humaidan@midt.rm.dk)

⁴Department of Medical Physiopathology, University of Rome "La Sapienza", Rome, Italy
(francesco.lombardo@uniroma1.it)

⁵IVI Foundation, Health Research Institute La Fe, Valencia, Spain
(Nicolas.Garrido@ivirma.com)

⁶Centro ANDROGEN, La Coruña, Spain (jalvarez@androgen.es)

⁷Division of Urology, Department of Surgery, St. Mary's Hospital, McGill University, Montreal, Québec, Canada (ziniarmand@yahoo.com)

⁸Andrology Service, Procrea Swiss Fertility Center, Lugano, Switzerland
(gmcolpi@yahoo.com)

⁹Andrology and IVF Department, San Carlo Cinic, Paderno Dugnano / Milano, Italy

¹⁰ Centre for Human Reproductive Science, IMSR, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK (j.kirkman-brown@nhs.net)

¹¹Birmingham Women's Fertility Centre, Birmingham Women's & Children's NHS Foundation Trust, Birmingham, UK

- 1
2
3 33 ¹²Queens University Belfast, Northern Ireland, United Kingdom (s.e.lewis@qub.ac.uk)
4
5 34 ¹³Examenlab Ltd., Weavers Court, Belfast, Northern Ireland, UK
6
7 35 ¹⁴ANOVA, Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden
8
9 36 (lars.bjorndahl@ki.se)
10 37 ¹⁵Department of Urology, Hamad Medical Corporation, Doha, Qatar
11
12 38 (dr.amajzoub@gmail.com)
13
14 39 ¹⁶Weill Cornell Medicine-Qatar, Doha, Qatar
15
16 40 ¹⁷Department of Surgery, Union Hospital, Hong Kong (chochaklam@yahoo.com.hk)
17
18 41 ¹⁸S.H. Ho Urology Centre, Department of Surgery, The Chinese University of Hong Kong,
19 Hong Kong
20
21 43 ¹⁹Urology/Andrology Unit, Saúde Atlântica, Clínica do Dragão, Porto, Portugal
22
23 44 (pvendeira@gmail.com)
24
25 45 ²⁰Androscience, Science and Innovation Center in Andrology and High-Complex Clinical and
26 Andrology Laboratory, São Paulo, SP, Brazil (hallakj@androscience.com.br)
27
28 47 ²¹Division of Urology, University of São Paulo, SP, Brazil;
29 Men's Health Study Group, Institute for Advanced Studies, University of São Paulo, SP,
30 Brazil
31
32 50 ²²Reproductive Toxicology Unit, Department of Pathology, University of São Paulo, SP,
33 Brazil
34
35 51 ²³Cabinet D'Andrologie Victor Hugo, American Hospital of Paris Reproductive Center,
36 Paris, France (tedamar@gmail.com)
37
38 53 ²⁴Human Reproduction Center, Division of Urology, University of São Paulo, SP, Brazil
39
40 54 (mcocuzza@uol.com.br)
41
42 55 ²⁵Edinburgh Assisted Conception Programme, EFREC, Royal Infirmary of Edinburgh,
43 Edinburgh, Scotland, UK (sciorioromualdo@hotmail.com)
44
45 57 ²⁶Fertility Clinic Skive, Skive Regional Hospital, Skive, Denmark
46
47 58 (Rita.Laursen@midt.rm.dk)
48
49 59 ²⁷IVF laboratory, Al Saad Specialist Hospitals, Jeddah, KSA (ahmadmetwalley@yahoo.com)
50
51 61 ²⁸Reproductive Medicine and Genetic Unit, GEBRI, Sadat City, Egypt
52
53 62 ²⁹Division of Reproductive Medicine and Andrology, Jindal Hospital, Meerut, UP, India
54
55 63 (drsunkjindal@gmail.com)
56
57 64 ³⁰PUR Clinic and University of Central Florida, Clermont, FL, USA (sijoip@gmail.com)
58
59
60

³¹Department of Urology, Miller School of Medicine, University of Miami, Miami, Florida, USA (ranjithrama@gmail.com)

³²Department of Neuroscience, Reproductive Science and Odontostomatology, University of Naples Federico II, Naples, Italy (alviggi@unina.it)

³³PIVET Medical Centre, Perth, Australia (jlyovich@pivet.com.au)

³⁴Curtin University, Perth, Australia

³⁵Cairns Fertility Centre, Cairns, Australia

³⁶Andrology Center, Department of Urology, Cleveland Clinic, Cleveland, Ohio, USA (agarwaa@ccf.org)

Corresponding author:

Sandro C. Esteves, MD., PhD. ANDROFERT, Andrology and Human Reproduction Clinic, Av. Dr. Heitor Penteado, 1463, Campinas 13075-460, SP, Brazil, Email: s.esteves@androfert.com.br, Tel. +55(19)3295-8877; <https://orcid.org/0000-0002-1313-9680>

Word count: 3465

ORCID numbers:

Sandro C. Esteves: <https://orcid.org/0000-0002-1313-9680>

Francesco Lombardo: <https://orcid.org/0000-0002-6943-2966>

Nicolás Garrido: <https://orcid.org/0000-0001-8271-5218>

Armand Zini: <https://orcid.org/0000-0002-2194-5578>

Giovanni M. Colpi: <https://orcid.org/0000-0003-1431-1777>

Jackson Kirkman-Brow: <https://orcid.org/0000-0003-2833-8970>

Sheena EM Lewis: <https://orcid.org/0000-0001-5665-1572>

Lars Björndahl: <http://orcid.org/0000-0002-4709-5807>

Ahmad Majzoub: <https://orcid.org/0000-0001-7423-6241>

Cho Chak-Lam: <https://orcid.org/0000-0003-2520-6833>

Jorge Hallak: <http://orcid.org/0000-0002-6452-0502>

Marcello Cocuzza: <https://orcid.org/0000-0002-5465-4304>

Fabiola Bento: <https://orcid.org/0000-0003-3772-1357>

Romualdo Sciorio: <https://orcid.org/0000-0002-7698-8823>

Ahmad M. Metwalley: <https://orcid.org/0000-0001-5751-9506>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

98 Sunil Jindal: <https://orcid.org/0000-0002-8088-8710>
99 Sijo Parekattil: <https://orcid.org/0000-0001-6847-8826>
100 Ranjith Ramasamy: <https://orcid.org/0000-0003-1387-7904>
101 Carlo Alviggi: <https://orcid.org/0000-0001-9116-020X>
102 Peter Humaidan: <https://orcid.org/0000-0001-6884-5366>
103 John Yovich: <https://orcid.org/0000-0002-9583-3683>
104 Ashok Agarwal: <http://orcid.org/0000-0003-0585-1026>

For Peer Review

Abstract

The prolonged lockdown of health facilities providing non-urgent gamete cryopreservation – as currently recommended by many reproductive medicine entities and regulatory authorities due to the SARS-CoV-2 pandemic will be detrimental for subgroups of male infertility patients. We believe the existing recommendations should be promptly modified and propose that the same permissive approach for sperm banking granted for men with cancer is expanded to other groups of vulnerable patients. These groups include male infertility patients (e.g., azoospermic men and cryptozoospermic) undergoing medical or surgical treatment to improve sperm quantity and quality, as well as males of reproductive age affected by inflammatory and systemic auto-immune diseases who are about to start treatment with gonadotoxic drugs or who are under remission. In both scenarios, the ‘fertility window’ may be transitory; postponing diagnostic semen analysis and sperm banking in these men could compromise the prospects of biological parenthood. Moreover, we provide recommendations on how to continue the provision of andrological services in a considered manner and a safe environment. Our opinion is timely and relevant given the fact that fertility services are currently rated as of low priority in most countries.

Keywords: SARS-CoV-2, Male infertility, Sperm banking, Semen analysis, Azoospermia, Systemic auto-immune diseases, Opinion.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Severe acute respiratory syndrome-coronavirus 2 (SARS-CoV-2) is a novel coronavirus and causative agent of COVID-19, a disease with potentially dangerous implications for human health. The remarkable increase in the number of infections by SARS-CoV-2 around the world raises the prospect of massive hospitalizations that few healthcare systems would be able to deal with. On this basis, governments around the world have announced the most far-reaching restrictions on personal freedom in modern history. The urgent need to avoid a collapse in the healthcare system is the justification for the implemented measures, and reproductive medicine societies, as well as regulatory authorities, decisively followed by issuing guidance based on expert best judgment. The key recommendations for practitioners include suspension of initiation of new fertility treatment and non-urgent gamete cryopreservation, as well as suspension of elective surgery and non-urgent diagnostic procedures.^{1,2} Sperm banking has been rated as of low priority, indicating that clinical harm is very unlikely if postponed for six months.³ Exceptions are oncological patients who require urgent fertility preservation.

Taking the above mentioned into account, we would like to raise a viewpoint hardly voiced so far. Our concerns are that, first of all, a prolonged lockdown of andrological services will be detrimental to subgroups of male infertility patients. Secondly, the andrological community is uneasy about how to provide optimal care to our patients without compromising safety. We, therefore, propose remedies to mitigate the consequences of a prolonged cessation of andrological services. The aim is to help authorities and healthcare providers identify which patients might be prioritized for the continuation of andrological services in a safe environment.

The Pandemic Facts

At the time of writing (April 21), the global deaths caused by SARS-CoV-2 represent approximately one percent of total deaths expected to occur worldwide over the first three months of the current year, with a wide variation in the reported death rates per country (www.worldometers.info/coronavirus). In total, more than 2.5 million infections by SARS-CoV-2 have been reported, 95% of which have been defined as mild. Among the severe or critical cases, the overwhelming majority affects people aged 50 and above. By contrast, the

reported death rate among individuals of reproductive age ranges remains low, ranging from 0.2% in China to 0.8% in the United States, with an estimated 1.5:1 male to female ratio, mainly affecting those individuals with pre-existing conditions, including cardiovascular disease, diabetes, chronic respiratory disease, hypertension, obesity, and cancer.⁴

The impact of SARS-CoV-2 for males in need of sperm banking

While it is prudent to advocate temporary social distancing and closure of non-emergency health services, we do not know how long this pandemic will last. Estimates ranging from 3 to 12 months have been projected, depending on how effective governments implement quarantine measures and how long it takes to achieve herd immunity. Thus, we would like to consider what a prolonged lockdown of clinics providing andrological services might mean for infertility patients. This consideration will focus primarily on priority recommendations for sperm banking and diagnostic semen analysis for patients seeking fertility rather than donors.

The ‘time’ variable is crucial in specific subgroups of infertile males. Besides reproductive-age oncological patients, loss of time is particularly consequential among patients under medical treatment aimed at improving sperm quantity or quality and in those with inflammatory or auto-immune diseases who will either start treatment –with potentially gonadotoxic drugs– or are under the ‘remission window’ of such treatment, as explained in more detail below. In both scenarios, the ‘fertility window’ may be transitory and, therefore, the implications of postponing diagnostic semen analysis and sperm banking in these men could permanently compromise the prospects of biological parenthood. Hence, the provision of andrological services cannot be considered a low priority. Our opinion is particularly important given the fact that healthcare providers are reluctant to recommend assisted conception in most cases –using either fresh or frozen-thawed sperm– as pregnancy might act as a comorbidity in women affected by SARS-CoV-2.^{5,6}

Cancer patients

Up to 30% of male cancer survivors lose their fertility potential after anti-cancer therapy.⁷ Chemotherapy, radiotherapy, and radical surgical procedures might irreversibly impair spermatogenesis and/or ejaculation. Cancer itself can also affect fertility directly (e.g.,

testicular cancer, Hodgkin’s lymphoma).⁸ Currently, the only reliable method of fertility preservation in reproductive-aged men with cancer is sperm banking.⁹ Sperm banking must be ideally completed before the start of gonadotoxic therapy. Specimens are usually collected by masturbation and ejaculated sperm are cryopreserved using slow or rapid freezing protocols. Before cryopreservation, the semen sample undergoes semen analysis, which is used to both assess the baseline sperm variables (e.g., count, motility, morphology) and to plan banking. After thawing, it is inevitable that sperm parameters are overall reduced, and such samples have to be used with intrauterine insemination (IUI) or assisted reproductive technology (ART) to allow these patients to have biological children.¹⁰ The costs associated with sperm banking are relatively low and most cancer patients who banked sperm were found to be pleased by having taken that decision.¹¹

Azoospermic/cryptozoospermic males

The most vulnerable male infertility patient during the SARS-CoV-2 pandemic is probably the non-obstructive azoospermic (NOA) or cryptozoospermic patient being medically treated to restore or improve spermatogenesis. An example is a patient with hypogonadotropic hypogonadism (HH), in whom azoospermia results from the lack of adequate testicular stimulation by pituitary gonadotropins.¹² In males with pre-pubertal and post-pubertal HH, gonadotropin treatment increases testicular size, promotes virilization, and restores spermatogenesis (to varying degrees) in up to 90% of patients, with reported pregnancy rates –either by natural intercourse or with the aid of IUI or ART– in up to 65%.¹²⁻¹⁴ However, the treatment duration is long –typically six months or longer– and expensive as well. Moreover, follow-up during treatment requires monitoring serum levels of pituitary and sexual hormones, as well as semen analyses. Sperm banking might be considered in men with HH who responded to therapy, i.e., have viable sperm in the ejaculate, in particular, when the continuation of gonadotropin therapy is neither possible (e.g., due to economic or logistic issues), nor desired. In patients who have not responded yet but have economic constraints to continue therapy (e.g., coexisting female factors), the medication dose and regimen could be adjusted (e.g., decrease hCG dose, suspend FSH injections) to keep intratesticular as well as serum testosterone levels within lower normal limits.

Another example refers to males with NOA due to spermatogenic failure, including men with rare numbers of sperm occasionally found in the ejaculate (cryptozoospermia), accounting for 60% of the azoospermia cases.¹⁵ Although the condition is untreatable, medical therapy has been explored as a way to optimize or induce spermatogenesis and, thus, increase the likelihood of having sperm retrieved surgically or ejaculated. A few cohort studies have shown that sperm can be occasionally found in the ejaculate after the use of medication for boosting intratesticular testosterone production, like hCG injections –alone or combined with FSH injections–, and estrogen receptor modulators, such as tamoxifen.¹⁶⁻¹⁹ Similar to HH patients, the continuation of gonadotropin therapy is not always possible nor desired in men with NOA due to spermatogenic failure who require sperm utilization or banking after the course of medical therapy. Moreover, immediate ART might not be an option in some countries with strict lockdown measures during the current SARS-CoV-2 pandemic. Thus, sperm banking is urged in patients with ejaculated sperm as a way to preserve fertility and allow future ART with ejaculated sperm. Naturally, semen analyses are required to monitor treatment results and identify who is eligible for sperm banking. Patients achieving cryptozoospermia or severe oligozoospermia after treatment may have a short window for sperm cryopreservation as their semen quality might deteriorate.²⁰ In such events, surgical sperm retrieval will be required, which could inflict both clinical and financial burdens on patients. Nevertheless, sperm retrieval and cryopreservation of testicular sperm should be considered in specific situations when a narrow window of opportunity exists. Sperm retrieval can be performed on an outpatient basis under local/intravenous anesthesia and is associated with minimal postoperative complications.²¹

Along the same lines, varicocelectomy has been used as an attempt to improve spermatogenesis in NOA men with a coexistent varicocele. Spermatogonia type B, pachytene spermatocytes, and early spermatids are vulnerable to heat stress associated with varicocele.²² In a systematic review comprising 468 patients with varicocele and NOA, 44% of treated patients had viable ejaculated sperm postoperatively, suitable for ICSI or cryopreservation.²³ These patients should also be monitored with semen analyses, and sperm cryopreservation recommended for those with ejaculate sperm due to the risk of relapse.²⁴

1
2
3
4 240 Lastly, loss of fertility and late obstruction have been reported in up to 12% and 50% of men
5
6 241 with obstructive azoospermia subjected to vasovasostomy and vasoepididymostomy,
7
8 242 respectively.^{25,26} Similar findings can also occur after the transrectal resection of the
9
10 243 ejaculatory ducts. Semen analysis is used to monitor patency status, and sperm banking could
11
12 244 be offered to those patients who experience a continuous decrease in sperm count/quality
13
14 245 during the follow-up as a way to avoid future sperm retrievals.^{26,27}

15 246 *Infertile men of advanced paternal age*

16
17 247 Infertile men of advanced paternal age (e.g., >50 years) have occasionally used sperm banking
18
19 248 for planning of medically assisted reproduction.^{28,29} Given that advanced age is a risk factor
20
21 249 for SARS-CoV-2 complications, and severe SARS-CoV-2 illness might be treated with non-
22
23 250 specific anti-viral drugs with possible gonadotoxic effects³⁰, sperm banking could be offered
24
25 251 to male infertility patients who are concerned about acquiring the infection.

26 252 *Inflammatory and systemic auto-immune diseases*

27
28 253 At present, the prevailing consensus is to allow gamete cryopreservation to continue for
29
30 254 oncological patients. However, males at reproductive age affected by non-oncological
31
32 255 conditions (i.e., inflammatory bowel diseases, autoimmune disorders) may also need
33
34 256 immediate sperm banking.³¹ Gonadotoxic drugs (e.g., cyclophosphamide, methotrexate,
35
36 257 mycophenolate mofetil, and mTOR inhibitors) are commonly used to control the
37
38 258 inflammatory process in such patients.

39
40
41 259 Inflammatory bowel disease (e.g., Crohn's disease, ulcerative colitis) mainly affects young
42
43 260 adults, and drugs used for treatment (e.g., sulfasalazine, azathioprine, methotrexate) appear to
44
45 261 harm sperm quality.³² The sulfapyridine metabolite of sulfasalazine impairs semen
46
47 262 parameters and increases the production of reactive oxygen species.^{32,33} Moreover,
48
49 263 pregnancy-related complications and the risk of congenital abnormalities might increase
50
51 264 when the father had used azathioprine before conception.³⁴⁻³⁶ Also, methotrexate (MTX) is an
52
53 265 immunosuppressive agent used to treat inflammatory and auto-immune diseases with known
54
55 266 teratogenic effects. The antifolate mechanism of MTX decreases DNA synthesis and inhibits
56
57 267 cellular proliferation, possibly resulting in oligozoospermia.³⁷

Likewise, young men may be affected by **systemic autoimmune diseases** (SADs) (e.g., systemic lupus erythematosus, rheumatoid arthritis, systemic sclerosis, ankylosing spondylitis, dermatomyositis, Behçet disease, psoriasis, among others).³⁸ In these patients, the chronic inflammation could adversely affect the hypothalamic-pituitary-testicular axis and the testicles directly, causing impairment of semen quality and quantity. However, gonadal dysfunction is primarily related to the effects of immunosuppressive therapy (e.g., alkylating agents, methotrexate, mycophenolate mofetil).³⁹ Among patients with inflammatory bowel disease or SAD considering fertility preservation, sperm banking might be conditioned to temporary discontinuation of therapy for at least 3-4 months or before initiation of therapy.³⁵ Several patients might have been planning for this 'fertility window' for an extended time, which unfortunately occurred during the **SARS-CoV-2** pandemic. Sperm banking is, therefore, an option for patients concerned about establishing a pregnancy during the **SARS-CoV-2** pandemic, in particular, those with semen abnormalities candidates for ART. On this basis, we would argue that the same permissive approach that has been granted for men with cancer to enable gamete preservation should be extended to male patients with inflammatory and autoimmune diseases.

What are the possible remedies?

We need to consider the health and psychological consequences of not offering the above patients andrological services. The lockdown of andrological services may have a devastating psychological impact on men undergoing fertility-related treatment. Like women, men undergoing fertility treatment may also experience anxiety and stress.^{40,41} This psychological distress can aggravate the feeling of fear and uncertainty imposed by the **SARS-CoV-2** pandemic⁴², which might have negative consequences for the reproductive outcome. The damage to the affected patients is difficult to measure, and it will take months, perhaps years before we can assess the broader implications of the current restrictive measures for patients as well as healthcare providers. While we believe that the various lockdowns will slow the spread of **SARS-CoV-2**, a strict lockdown is unlikely to last too long due to its practicality and pitfalls on other aspects of society, mainly economical. Thus, a certain level of risk of infections by **SARS-CoV-2** is expected because there will be new cases when

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

measures are relaxed, and no vaccine is likely to be available soon. Therefore, not only urgent short-term responses, but also long-term measures are essential. Hence, in this time of uncertainty, denying andrological services from those who need it most might be even worse than the risks of providing them. We, therefore, propose some remedies that we believe might offer fertility providers and patients alike greater autonomy, and that could be used to alleviate the adverse impact of the coronavirus pandemic in the months to come (**Figure 1**).

1. Before any service is provided, active SARS-CoV-2 infections and suspected cases should be excluded. Testing patients with the use of polymerase chain reaction (PCR) and/or blood antibody testing is recommended before starting sperm banking. Ideally, only samples from patients with negative results or who have acquired herd immunity should be cryopreserved.

2. Andrological services (e.g., diagnostic semen analysis and sperm cryopreservation) should not only be available for oncological patients, but also for the group of patients listed below.

i. Men with severe male factor infertility under medical or surgical treatment aiming at improving sperm quantity or quality (e.g., patients with NOA or cryptozoospermia/severe oligozoospermia, including post-varicocele repair, and those with evidence of loss of patency after successful surgical reconstruction of the reproductive tract).

ii. Men at reproductive age affected by inflammatory diseases or SADs, i.e., before initiation of gonadotoxic therapy or if under the ‘fertility window’ achieved after temporary (at least three months) discontinuation of therapy.

iii. Infertile men older than 50 years, in particular those with comorbidities who are candidates for IUI or ART and are concerned about the risk of acquiring SARS-CoV-2 and the possibility of anti-viral therapy causing gonadotoxic effects.

3. Surgical sperm retrieval and cryopreservation of testicular sperm should be considered in specific situations involving men with NOA undergoing medical therapy to improve spermatogenesis. In this setting, procedures should be performed, if possible, on an outpatient basis under local anesthesia. Moreover, the use of electrocautery should be avoided as the surgical smoke might carry the virus if a patient is infected but asymptomatic. Only essential staff should stay in the operating theater, and personal protection measures should be strictly followed as determined by the local healthcare authorities. In closed-controlled air systems, the airflow might produce an increase in the viral spread from potential asymptomatic

patients. Thus, special attention should be given to air quality control, including the use of air filtration systems, particularly in surgical and laboratory areas.⁴³

4. Encourage telemedicine and phone counseling for providing instructions about testing and sperm banking.

5. Adherence to infection prevention recommendations is of utmost importance for patients and health practitioners alike. This advice includes the use of appropriate personal protective equipment (PPE) by healthcare staff, adherence to social distancing measures for healthcare staff and patients, and space out appointments so that no patients are waiting together in the clinic waiting area. We stress the importance of training staff (receptionists, nurses, technicians, doctors) on PPE needs and usage (please see

<https://www.cdc.gov/coronavirus/2019-ncov/hcp/clinic-preparedness.html>).

6. Good laboratory practices should be strictly applied when handling the seminal fluid in the andrology laboratory.⁴⁴ This advice includes (i) use of class II safety cabinets⁴⁵, which gives protection to the specimen handled as well as the operator performing the work, (ii) use of high-security straws for sperm cryopreservation, as routinely used in most sperm banks, and (iii) additional measures to protect the specimens from laboratory staff (e.g., use of goggles, N95 mask, gown/coverall, and gloves) –who might be asymptomatic for SARS-CoV-2.

7. Technicians/biologists should, ideally, be tested by PCR and/or blood antibody testing before resuming activities, and only staff with negative results or who have acquired herd immunity should perform laboratory duties. If the staff that manipulated specimens get infected, an aliquot of cryopreserved semen samples should be tested (e.g., by PCR) because semen samples, cryopreservation media, straws, and pipette tips could have been contaminated by asymptomatic PCR-positive biologists and technicians.

8. A thorough discussion between patients and healthcare providers should be made for responsible shared decisions. This advice includes the development and use of dedicated informed consent, detailing the risks of attending the facility and banking of sperm during the SARS-CoV-2 pandemic. Furthermore, psychological support and financial aid might be offered to those in need. The latter might be particularly relevant to patients under economic pressure due to the pandemic who need to afford the costs of semen analysis and sperm banking.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

9. Advanced planning should guide the continuation of andrological services. Working groups and quality managers should determine which patients to prioritize and how working lists should be filled, including staff scheduling.

Practical considerations

During the coming weeks, we should continue to look critically and objectively at the SARS-CoV-2 evidence. Although our recommendations are unlikely to create any further burden to the already overwhelmed medical infrastructure, we acknowledge that patients might be reluctant to use andrological services on the basis of fear of being infected or economic reasons. We also realize that much is unknown about SARS-CoV-2 and its implication on male reproductive health. The existing data indicate that a subject can be infectious 3-5 days before the onset of actual symptoms of the viral infection, and the risk of such cases spreading the infection has not been rigorously researched.⁴⁶ While testing patients and staff with the use of PCR and/or antibody kits is recommended, the majority of clinics lack prompt access to these tests. Moreover, some countries face a short supply of test kits, which have been made available for symptomatic patients and frontline health providers only. Besides, the accuracy of these tests has been questioned, with some reports suggesting that many of the SARS-CoV-2 kits in the market have a false negative rate of 30-40%.⁴⁷ Thus, it remains to be determined how clinics can screen patients and healthcare providers optimally. Likewise, it remains to be decided who –patients or clinics– will assimilate the testing related-costs, PPE, and reduced patient volume due to extra measures instituted to avoid infections. Along these lines, clinics and hospitals providing andrological services have to determine ways of protecting themselves from potential liability issues. Although the overall mortality rate among men at reproductive age remains low, it should be considered that contamination of patients and staff could occur with SARS-CoV-2 in the context of asymptomatic shedding. For this reason, it seems sound to advise postponing medical therapy in azoospermic men who had planned to initiate medical therapy and who have no pressing concerns (e.g., no maternal factors such as advanced maternal age) until it is deemed safe to obtain regular semen analyses, hormone profiles, and banking of sperm. The same reasoning applies to semen analysis and sperm banking in men under therapy who opt to continue on medication till the pandemic ends.

At present, limited data exist about potential routes of SARS-CoV-2 infection in respiratory, cardiovascular, digestive, urinary, and reproductive systems. In this regard, no evidence of virus load in semen or testicular biopsies of SARS-CoV-2 infected patients has been reported, but data is minimal.⁴⁸⁻⁵⁰ Nevertheless, angiotensin-converting enzyme 2 (ACE2) receptors, used by the virus to enter host cells, might exist in spermatogonia, Sertoli cells, and Leydig cells^{51,52}, as well as sperm cells.⁵³ Also, previous reports suggested that other coronaviruses, like the SARS coronavirus, could cause orchitis.⁵⁴

As for pregnancy with the use of banked or fresh ejaculate sperm during the SARS-CoV-2 pandemic, it has been suggested that pregnant women might be at a higher risk of developing complications, including miscarriage, preeclampsia, and preterm birth.^{5,6} However, the evidence is still limited, and we, therefore, abstain from making recommendations about the use of fresh or banked sperm for assisted conception during the pandemic until more data are available. Naturally, the use of sperm for assisted conception –either fresh or frozen-thawed– would not be recommended in most cases if it is confirmed that pregnancy acts as an important comorbidity factor. Notwithstanding these observations, it should be acknowledged that serology testing, once properly validated and widely available, will be helpful to identify immune patients that could be allowed for treatment.⁵⁵ These patients have little risk of either pregnancy complications or propagating the disease when attending fertility clinics. Nevertheless, the provision of andrological services should only be undertaken if the medical infrastructure can support them. We reiterate the above recommendations that care should only be restarted if social distancing can be maintained, areas regularly disinfected, and screening for signs and symptoms of the infection undertaken before allowing patients into the facility in accordance with guidance issued by health regulatory authorities.

Conclusions

We propose remedies to mitigate the consequences of a prolonged cessation of andrological services due to the SARS-CoV-2 pandemic to vulnerable subgroups of male infertility patients. In a moment when the reorganization of healthcare services is focused on supporting SARS-CoV-2 patients who might need critical care, limiting burdens for national health systems could still represent a relevant issue. We advocate that correct identification of the more “time-sensitive” cases is crucial for regulating the continuation of andrological services,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

including diagnostic semen analysis and sperm banking. Moreover, we provide recommendations on how to most optimally provide care to our patients –without compromising safety– once andrological services are resumed. The aim is to help authorities and healthcare providers identify which patients might be prioritized during the SARS-CoV-2 pandemic for the continuation of andrological services in a safe environment.

Acknowledgments: None

Author Contributions

SCE contributed to the conception, designed the manuscript, and wrote the first draft. JA, NG, FB, GP, AM, C-LC, PV, JH, MC, JY, and AA wrote sections of the manuscript. All authors contributed to manuscript revision and critical analysis, read and approved the submitted version. The corresponding author takes the final responsibility for the decision to submit the manuscript for publication.

Declaration of interests

SCE and CA declares the receipt of unrestricted research grants and lecture fees from Merck outside the submitted work. SEMML is an employee of Examenlab Ltd., a university spin-out company with a commercial interest in sperm DNA damage. FL, NG, JA, AZ, GMC, JK-B, LB, AH, C-LC, PV, JH, EA, MC, FCB, RCF, RS, RL, AMM, SKJ, SP, RR, PH, JLY, and AA declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding: None

References

1. American Society for Reproductive Medicine (ASRM). American Society for Reproductive Medicine Patient Management And Clinical Recommendations during The Coronavirus (Covid-19) Pandemic. 2020. Available from: <https://www.asrm.org/news-and-publications/covid-19/statements/patient-management-and-clinical-recommendations-during-the-coronavirus-covid-19-pandemic/> [cited 2020, April 14].
2. European Society for Human Reproduction and Embryology (ESHRE). Coronavirus Covid-19: ESHRE statement on pregnancy and conception. 2020.
3. European Association of Urology (EAU) COVID-19 Recommendations. EAU Guidelines Office Rapid Reaction Group: An organization-wide collaborative effort to adapt the EAU guidelines recommendations to the COVID era. Available from:

- 448 [https://uroweb.org/wp-content/uploads/Combined-non-oncology-COVID-](https://uroweb.org/wp-content/uploads/Combined-non-oncology-COVID-recommendations.pdf)
449 [recommendations.pdf](https://uroweb.org/wp-content/uploads/Combined-non-oncology-COVID-recommendations.pdf) [cited 2020, April 18].
- 450 4. COVID-10 Coronavirus Pandemic. Available from:
451 <https://www.worldometers.info/coronavirus/> [cited 2020, April 16].
- 452 5. Yu N, Li W, Kang Q, et al. Clinical features and obstetric and neonatal outcomes of
453 pregnant patients with COVID-19 in Wuhan, China: a retrospective, single-centre,
454 descriptive study. *Lancet Infect Dis*. 2020 Mar 24 [Epub ahead of print].
- 455 6. Chen H, Guo J, Wang C, et al. Clinical characteristics and intrauterine vertical
456 transmission potential of COVID-19 infection in nine pregnant women: a
457 retrospective review of medical records. *Lancet*. 2020;395(10226):809-15.
- 458 7. Johnson MD, Cooper AR, Jungheim ES, Lanzendorf SE, Odem RR, Ratts VS. Sperm
459 banking for fertility preservation: a 20-year experience. *Eur J Obstet Gynecol*
460 *Reprod Biol*. 2013;170:177-82.
- 461 8. Xu R, Centola GM, Tanrikut C. Genitourinary cancer patients have worse
462 baseline semen parameters than healthy sperm bankers. *Andrology*. 2019;7:449-453.
- 463 9. Agarwal A, Ong C, Durairajanayagam D. Contemporary and future insights into fertility
464 preservation in male cancer patients. *Transl Androl Urol*. 2014;3:27-40.
- 465 10. Freour T, Mirallie S, Jean M, Barriere P. Sperm banking and assisted reproductive
466 outcome in men with cancer: a 10 years' experience. *Int J Clin Oncol*. 2012;17:598-603.
- 467 11. Williams DH. Sperm banking and the cancer patient. *Ther Adv Urol*. 2010;2:19-34.
- 468 12. Fraietta R, Zylberstejn DS, Esteves SC. Hypogonadotropic hypogonadism revisited.
469 *Clinics (Sao Paulo)* 2013;68 Suppl 1:81-88.
- 470 13. Yang L, Zhang SX, Dong Q, Xiong ZB, Li X. Application of hormonal treatment in
471 hypogonadotropic hypogonadism: more than ten years experience. *Int Urol Nephrol*.
472 2012;44:393-9.
- 473 14. Behre HM. Clinical Use of FSH in Male Infertility. *Front Endocrinol*
474 (Lausanne). 2019;10:322.
- 475 15. Esteves SC. Clinical management of infertile men with nonobstructive azoospermia.
476 *Asian J Androl*. 2015;17:459-470.
- 477 16. Moein MR, Tabibnejad N, Ghasemzadeh J. Beneficial effect of tamoxifen on sperm
478 recovery in infertile men with nonobstructive azoospermia. *Andrologia* 2012;44 Suppl 1:194-
479 198.

17. Shinjo E, Shiraishi K, Matsuyama H. The effect of human chorionic gonadotropin-based hormonal therapy on intratesticular testosterone levels and spermatogonial DNA synthesis in men with non-obstructive azoospermia. *Andrology* 2013;1:929-935.
18. Hussein A, Ozgok Y, Ross L, Rao P, Niederberger C. Optimization of spermatogenesis-regulating hormones in patients with non-obstructive azoospermia and its impact on sperm retrieval: a multicentre study. *BJU Int.* 2013;111 (3 Pt B):E110-114.
19. Laursen RJ, Elbaek HO, Povlsen BB, Lykkegaard J, Jensen KBS, Esteves SC, et al. Hormonal stimulation of spermatogenesis: a new way to treat the infertile male with non-obstructive azoospermia? *Int Urol Nephrol.* 2019;51:453-456.
20. Koscinski I, Wittemer C, Lefebvre-Khalil V, Marcelli F, Defossez A, Rigot JM. Optimal management of extreme oligozoospermia by an appropriate cryopreservation programme. *Hum Reprod.* 2007;22:2679-84.
21. Miyaoka R, Orosz JE, Achermann AP, Esteves SC. Methods of surgical sperm extraction and implications for assisted reproductive technology success. *Panminerva Med.* 2019;61:164-177.
22. Agarwal A, Hamada A, Esteves SC. Insight into oxidative stress in varicocele-associated male infertility: part 1. *Nat Rev Urol.* 2012;9:678-90.
23. Esteves SC, Miyaoka R, Roque M, Agarwal A. Outcome of varicocele repair in men with nonobstructive azoospermia: systematic review and meta-analysis. *Asian J Androl.* 2016;18:246-253.
24. Pasqualotto FF, Sobreiro BP, Hallak J, Pasqualotto EB, Lucon AM. Induction of spermatogenesis in azoospermic men after varicoectomy repair: an update. *Fertil Steril.* 2006;85:635-9.
25. Carbone DJ Jr, Shah A, Thomas AJ Jr, Agarwal A. Partial obstruction, not antisperm antibodies, causing infertility after vasovasostomy. *J Urol.* 1998;159:827-30.
26. Farber NJ, Flannigan R, Li P, Li PS, Goldstein M. The Kinetics of Sperm Return and Late Failure Following Vasovasostomy or Vasoepididymostomy: A Systematic Review. *J Urol.* 2019;201:241-250.
27. Esteves SC, Miyaoka R, Agarwal A. Surgical treatment of male infertility in the era of intracytoplasmic sperm injection - new insights. *Clinics (Sao Paulo).* 2011;66:1463-78.
28. Jennings MO, Owen RC, Keefe D, Kim ED. Management and counseling of the male with advanced paternal age. *Fertil Steril.* 2017;107:324-328.

- 513 29. Bertoncelli Tanaka M, Agarwal A, Esteves SC. Paternal age and assisted
 514 reproductive technology: problem solver or trouble maker? *Panminerva Med.* 2019;61:138-
 515 151.
- 516 30. Drobnis EZ, Nangia AK. Antivirals and Male Reproduction. *Adv Exp Med Biol.*
 517 2017;1034:163-178.
- 518 31. Choux C, Cavalieri M, Barberet J, Samson M, Bonnotte B, Fauque P, Sagot P.
 519 [Immunosuppressive therapy and fertility preservation: Indications and methods].
 520 *Rev Med Interne.* 2018;39:557-565.
- 521 32. Shin T, Kobori Y, Suzuki K, Iwahata T, Yagi H, Soh S, et al. Inflammatory bowel disease
 522 in subfertile men and the effect of mesalazine on fertility. *Syst Biol Reprod*
 523 *Med.* 2014;60:373–376.
- 524 33. Alonso V, Linares V, Bellés M, Albina ML, Sirvent JJ, Domingo JL, Sánchez DJ.
 525 Sulfasalazine induced oxidative stress: a possible mechanism of male infertility. *Reprod*
 526 *Toxicol.* 2009;27:35–40.
- 527 34. Rajapakse RO, Korelitz BI, Zlatanich J, Baiocco PJ, Gleim GW. Outcome of pregnancies
 528 when fathers are treated with 6-mercaptopurine for inflammatory bowel disease.
 529 *Am J Gastroenterol.* 2000;95:684-8.
- 530 35. Palomba S, Sereni G, Falbo A, Beltrami M, Lombardini S, Boni MC, et al. Inflammatory
 531 bowel diseases and human reproduction: a comprehensive evidence-based review. *World J*
 532 *Gastroenterol.* 2014; 20:7123-36.
- 533 36. Nørgård B, Pedersen L, Jacobsen J, Rasmussen SN, Sørensen HT. The risk of congenital
 534 abnormalities in children fathered by men treated with azathioprine or mercaptopurine before
 535 conception. *Aliment Pharmacol Ther.* 2004;19:679-85.
- 536 37. Sands K, Jansen R, Zaslau S, Greenwald D. Review article: the safety of therapeutic
 537 drugs in male inflammatory bowel disease patients wishing to conceive. *Aliment Pharmacol*
 538 *Ther.* 2015;41(9):821-34.
- 539 38. Brubaker WD, Li S, Baker LC, Eisenberg ML. Increased risk of autoimmune disorders in
 540 infertile men: analysis of US claims data. *Andrology.* 2018;6:94-98.
- 541 39. Tiseo BC, Cocuzza M, Bonfa E, Srougi M, Silva CA. Male fertility potential alteration in
 542 rheumatic diseases: a systematic review. *Int Braz J Urol.* 2016;42:11-21.
- 543 40. Peterson BD, Newton CR, Feingold T. Anxiety and sexual stress in men and women
 544 undergoing infertility treatment. *Fertil Steril.* 2007;88:911-4.

41. Boivin J, Schmidt L. Infertility-related stress in men and women predicts treatment outcome 1 year later. *Fertil Steril*. 2005;83:1745-52.
42. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int J Environ Res Public Health*. 2020;17(5).
43. Esteves SC, Bento FC. Implementation of air quality control in reproductive laboratories in full compliance with the Brazilian Cells and Germinative Tissue Directive. *Reprod Biomed Online*. 2013;26:9-21.
44. Esteves SC, Agarwal A. Explaining how reproductive laboratories work. In: Bento FC, Esteves SC, Agarwal A (eds). *Quality management in ART Clinics: a practical guide*. Springer, New York, 1st. ed. 2013, pp. 79-127.
45. Ramstorp M. What is a clean room? In: Esteves SC, Varghese A, Worilow KC (eds.). *Clean room technology in ART clinics: a practical guide*. CRC Press, Boca Raton, 1st. ed. 2017, pp. 3-18.
46. Yuen KS, Ye ZW, Fung SY, Chan CP, Jin DY. SARS-CoV-2 and COVID-19: The most important research questions. *Cell Biosci*. 2020;10:40.
47. Colin P. West, Victor M. Montori, Priya Sampathkumar. COVID-19 Testing: The Threat of False-Negative Results. *Mayo Clinic Proceedings*, 2020; doi:10.1016/j.mayocp.2020.04.004
48. Song C, Wang Y, Li W, Hu B, Chen G, Xia P, et al. Detection of 2019 novel coronavirus in semen and testicular biopsy specimen of COVID-19 patients. Available from: <https://www.medrxiv.org/content/10.1101/2020.03.31.20042333v2.article-info> [cited 2020, April 18]. doi: <https://doi.org/10.1101/2020.03.31.20042333>
49. Pan F, Xiao X, Guo J, Song Y, Li H, Patel DP, et al. No evidence of SARS-CoV-2 in semen of males recovering from COVID-1. *Fertil Steril*. Ahead of print April 17, <https://doi.org/10.1016/j.fertnstert.2020.04.024>.
50. Paoli D, Pallotti F, Colangelo S, Basilico F, Mazzuti L, Turriziani O, et al. Study of SARS-CoV-2 in semen and urine samples of a volunteer with positive rino-pharyngeal swab. *J Endocrinol Invest*. 2020 (in press). <https://doi.org/10.1007/s40618-020-01261-1>.
51. Pan PP, Zhan QT, Le F, Zheng YM, Jin F. Angiotensin-converting enzymes play a dominant role in fertility. *Int J Mol Sci*. 2013;14:21071-86.
52. Wang Z, Xu X. scRNA-seq Profiling of Human Testes Reveals the Presence of the


- 1
2
3 578 ACE2 Receptor, A Target for SARS-CoV-2 Infection in Spermatogonia, Leydig and
4
5 579 Sertoli Cells. *Cells*. 2020;9(4).
6
7 580 53. Gianzo M, Muñoa-Hoyos I, Urizar-Arenaza I, Larreategui Z, Quintana F, Garrido
8
9 581 N, et al. Angiotensin II type 2 receptor is expressed in human
10
11 582 sperm cells and is involved in sperm motility. *Fertil Steril*. 2016;105:608-616.
12
13 583 54. Xu J, Qi L, Chi X, Yang J, Wei X, Gong E, et al. Orchitis: a complication of severe acute
14
15 584 respiratory syndrome (SARS). *Biol Reprod*. 2006;74:410-6.
16
17 585 55. Petherick A. Developing antibody tests for SARS-CoV-2. *Lancet* 2020;395:1101-02.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Legend to Figure

Figure 1. SARS-CoV-2 pandemic and provision of andrological services: proposal for individualized management.

For Peer Review



*HH: hypogonadotropic hypogonadism; NOA: non-obstructive azoospermia; IUI: intrauterine insemination; ART: assisted reproductive technology

Figure 1. SARS-CoV-2 pandemic and provision of andrological services: proposal for individualized management.

677x381mm (300 x 300 DPI)