### UNIVERSITY<sup>OF</sup> BIRMINGHAM University of Birmingham Research at Birmingham

# Expression of SOAT1 in adrenocortical carcinoma and response to mitotane monotherapy

Weigand, Isabel; Altieri, Barbara; Lacombe, Amanda; Basile, Vittoria ; Kircher, Stefan; Landwehr, Laura-Sophie; Schreiner, Jochen; Zerbini, Maria; Ronchi, Cristina; Megerle, Felix; Berruti, Alfredo; Canu, Letizia; Volante, Marco; Paiva, Isabel; Della Casa, Silvia; Sbiera, Silviu; Fassnacht, Martin; Fragoso, Maria ; Terzolo, Massimo; Kroiss, Matthias *DOI:* 

10.1210/clinem/dgaa293

License: None: All rights reserved

Document Version Peer reviewed version

Citation for published version (Harvard):

Weigand, I, Altieri, B, Lacombe, A, Basile, V, Kircher, S, Landwehr, L-S, Schreiner, J, Zerbini, M, Ronchi, C, Megerle, F, Berruti, A, Canu, L, Volante, M, Paiva, I, Della Casa, S, Sbiera, S, Fassnacht, M, Fragoso, M, Terzolo, M & Kroiss, M 2020, 'Expression of SOAT1 in adrenocortical carcinoma and response to mitotane monotherapy: an ENSAT multicenter study', *The Journal of clinical endocrinology and metabolism*, vol. 105, no. 8. https://doi.org/10.1210/clinem/dgaa293

Link to publication on Research at Birmingham portal

#### **Publisher Rights Statement:**

Tubis is a pre-copyedited, author-produced version of an article accepted for publication in Journal of Clinical Endocrinology and Metabolism following peer review. The version of recordIsabel Weigand, Barbara Altieri, Amanda M F Lacombe, Vittoria Basile, Stefan Kircher, Laura-Sophie Landwehr, Jochen Schreiner, Maria C N Zerbini, Cristina L Ronchi, Felix Megerle, Alfredo Berruti, Letizia Canu, Marco Volante, Isabel Paiva, Silvia Della Casa, Silviu Sbiera, Martin Fassnacht, Maria Candida B V Fragoso, Massimo Terzolo, Matthias Kroiss, Expression of SOAT1 in adrenocortical carcinoma and response to mitotane monotherapy: an ENSAT multicenter study, The Journal of Clinical Endocrinology & Metabolism, is available online at: https://doi.org/10.1210/clinem/dgaa293

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

### Expression of sterol-O-acyl transferase 1 (SOAT1) in adrenocortical carcinoma and response to mitotane monotherapy: an ENSAT multicenter study

3

Isabel Weigand<sup>1</sup>, Barbara Altieri<sup>1</sup>, Amanda M. F. Lacombe<sup>2</sup>, Vittoria Basile<sup>3</sup>, Stefan Kircher<sup>4</sup>,
Laura-Sophie Landwehr<sup>1</sup>, Jochen Schreiner<sup>1</sup>, Maria C. N. Zerbini<sup>2</sup>, Cristina L. Ronchi<sup>1,5,6</sup>,
Felix Megerle<sup>1</sup>, Alfredo Berruti<sup>7</sup>, Letizia Canu<sup>8</sup>, Marco Volante<sup>9</sup>, Isabel Paiva<sup>10</sup>, Silvia Della
Casa<sup>11</sup>, Silviu Sbiera<sup>1</sup>, Martin Fassnacht<sup>1,12</sup>, Maria Candida B. V. Fragoso<sup>2</sup>, Massimo
Terzolo<sup>3</sup>, Matthias Kroiss<sup>1,12</sup>

<sup>1</sup>Department of Internal Medicine I, Division of Endocrinology and Diabetes, University
 Hospital, University of Würzburg, Würzburg, Germany

<sup>2</sup> Unidade de Suprarrenal, Laboratorio de Hormônios e Genetica Molecular LIM42, Serviço
de Endocrinologi e Metaboligia, Hospital de Clínicas; Departamento de Patologia, Faculdade
de Medicina da Universidade de São Paulo, Brazil, Instituto do Câncer do Estado de São

14 Paulo (ICESP), Faculdade de Medicina da Universidade de São Paulo, São Paul, Brazil

- <sup>3</sup>Department of Clinical and Biological Sciences, University of Turin at San Luigi Hospital,
   Turin, Italy
- <sup>4</sup>Institute of Pathology, University of Würzburg, Würzburg, Germany
- <sup>5</sup>Institute of Metabolism and System Research, University of Birmingham, Birmingham, UK
- <sup>6</sup>Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners
   Birmingham, UK
- <sup>7</sup>Medical Oncology Unit, Department of Medical and Surgical Specialties, Radiological
   Sciences and Public Health, University of Brescia at ASST Spedali Civili, Brescia, Italy
- <sup>8</sup>Endocrinology Unit, Department of Experimental and Clinical Biomedical Sciences,
  University of Florence, Florence, Italy
- <sup>9</sup>Department of Oncology, University of Turin, Turin, Italy
- 26 <sup>10</sup> University Hospital of Coimbra, Coimbra, Portugal
- <sup>11</sup> Division of Endocrinology and Metabolism, Fondazione Policlinico Gemelli, Catholic
   University, Rome, Italy
- <sup>12</sup>Comprehensive Cancer Center Mainfranken, University of Würzburg, Würzburg, Germany

- 30 Short title: SOAT1 expression and mitotane
- 31 Key words: adrenal cancer, chemotherapy, treatment, prognosis
- 32
- 33 Correspondence and request for reprints to:
- 34 Matthias Kroiss, MD, PhD
- 35 Division of Endocrinology and Diabetology,
- 36 University Hospital of Würzburg,
- 37 Oberdürrbacher Str. 6
- 38 97080 Würzburg (Germany)
- 39 Telephone: +49-931-201-39740
- 40 Email: Kroiss m@ukw.de
- 41

#### 42 **Disclosure**:

43 The authors have nothing to disclose.

#### 44 Financial support:

This work has been supported by grants from the DFG German Research Foundation (grant KR-4371/1-2 to M.K, FA-466/4-2 to M.F. and RO-5435-3-3 to C.L.R.), the DFG German Research Foundation Project 314061271- TRR 205 (M.F. & M.K.) from Fundação de Amparo à Pesquisa do Estado de São Paulo (grant 2017/26345-5 to M.C.B.V.F.), Else Kröner-Fresenius-Stiftung (Grant 2016\_A96 to S.S. and M.K) and from Deutsche Krebshilfe (grant 70112969 to C.L.R. & M. F.).

- 51
- 52 **Precis** (max 200 Zeichen)
- 53 Mitotane is a cornerstone of adrenal cancer treatment. In this international study, expression 54 of putative mitotane target SOAT1 in tissue did not predict treatment response to mitotane 55 monotherapy.
- 56
- 57

#### 58 Abstract (max 250 words)

59 **Context** Objective response rate to mitotane in advanced adrenocortical carcinoma (ACC) is 60 approximately 20% and adverse drug effects are frequent. To date there is no marker 61 established that predicts treatment response. Mitotane has been shown to inhibit sterol-O-62 acyl transferase 1 (SOAT1) which leads to endoplasmic reticulum stress and cell death in 63 ACC cells.

64 **Objective** To investigate SOAT1 protein expression as a marker of treatment response to 65 mitotane.

66 **Patients** 231 ACC patients treated with single agent mitotane as adjuvant (n=158) or 67 advanced disease therapy (n=73) from twelve ENSAT centers were included. SOAT1 protein 68 expression was determined by immunohistochemistry on formalin-fixed paraffin-embedded 69 (FFPE) specimens.

Main outcome measure: recurrence-free survival (RFS), progression-free survival (PFS),
 disease-specific survival (DSS)

72 Results 61/135 patients (45 %) with adjuvant mitotane treatment had recurrences and 45/68 73 patients (66 %) with mitotane treatment for advanced disease had progressive disease. After 74 multivariate adjustment for sex, age, hormone secretion, tumour stage and Ki67 index, RFS 75 (HR=1.07, 95% CI 0.61-1.85, p=0.82) and DSS (HR=1.30, 95% CI 0.58-2.93, p=0.53) in 76 adjuvantly treated ACC patients did not differ significantly between tumors with high and low 77 SOAT1 expression. Similarly, in the advanced stage setting, PFS (HR=1.34, 95% CI 0.63-78 2.84, p=0.45) and DSS (HR=0.72, 95% CI 0.31-1.70, p=0.45) were comparable and 79 response rates not significantly different.

80 **Conclusions** SOAT1 expression was not correlated with clinical endpoints RFS, PFS and 81 DSS in ACC patients with mitotane monotherapy. Other factors appear to be relevant for 82 mitotane treatment response and ACC patient survival.

83

#### 84 Introduction

Adrenocortical carcinoma (ACC) is a rare malignancy with a generally poor prognosis 85 [1] and limited effective treatment options [1, 2]. Mitotane is the only approved drug 86 for metastatic disease [3] but efficacy is very limited and the observed objective 87 response rate is only approximately 20 % [4-6]. Controversy exists regarding 88 89 adjuvant use which is supported by a large retrospective study [6, 7] and advocated by current guidelines [2] in patients at moderate or high risk of recurrence after 90 complete resection. Adverse drug effects like adrenal insufficiency, diarrhea, nausea 91 and other gastrointestinal symptoms but also central nervous symptoms such as 92 dizziness and speech disturbance may be severe and disabling [8-10] and must be 93 balanced against potential treatment benefits. Mitotane efficacy is correlated with 94 95 plasma concentrations above 14 mg/l [11]. Therapeutic drug monitoring (TDM) is therefore recommended [2]. Some patients for unknown reasons fail to achieve 96 97 mitotane plasma concentrations within the therapeutic window which is associated 98 with decreased efficacy [5, 12]. To date, few markers have been suggested for the prediction of response [13, 14], but they have not been validated in a large series. 99 Establishment of such a marker would be a major advancement in ACC treatment 100 101 and enable tailored treatment of potential responders and avoidance of unnecessary 102 mitotane exposure in non-responders.

We have provided evidence that mitotane inhibits sterol-O-acyl transferase 1 (SOAT1) also known as ACAT1 [15] (not to be mistaken with acetyl-CoA acetyltransferase known under the same name), an enzyme catalyzing the esterification of cholesterol in the adrenal cortex [16]. This leads to the accumulation of toxic lipids and endoplasmic reticulum (ER) stress which results in apoptosis of adrenocortical cells [15]. Accordingly, a SOAT1 inhibitor has been tested in a phase I clinical trial as a treatment for advanced ACC [17].

SOAT1 is strongly expressed in adrenocortical cell lines, normal adrenal glands and different adrenocortical tumor entities, with the highest variation among ACC, while it is only weakly to moderately expressed in non-adrenal tissues [15]. Despite strong evidence of an inhibitory effect on SOAT1, other mechanisms such as impaired mitochondrial respiration and function [18-20] may contribute to the relatively tissuespecific toxicity of mitotane. In a small cohort of patients with advanced ACC, it has been shown that SOAT1expression was correlated with the response to mitotane treatment [15].

Here, we aimed to validate in a large multicenter study whether SOAT1 expression is a predictive marker for mitotane efficacy by investigating the association of SOAT1 tissue expression with recurrence free survival (RFS) in patients with adjuvant mitotane treatment, progression-free survival (PFS) after mitotane monotherapy administered to patients with advanced disease and disease-specific survival (DSS) for both cases.

#### 124 Patients and Methods

#### 125 Setting and data acquisition

126 Formalin-fixed paraffin-embedded (FFPE) tumor specimens of 231 ACC were included from 127 12 centers belonging to the European Network for the Study of Adrenocortical Tumors 128 (ENSAT; www.ensat.org). Only adult patients with histologically confirmed ACC were 129 included [21]. Patients that have been included in our previous analyses of SOAT1 130 expression [15] have been excluded from this analysis. All patients started mitotane 131 treatment as first medical therapy no later than 3 months after complete resection in the 132 adjuvant setting (n=158) or diagnosis of irresectable or recurrent or metastatic ACC in the 133 advanced stage setting (n=73). The study was conducted as part of the ENSAT registry, has 134 been approved by the ethics committee at each participating institution and was conducted in 135 accordance with the principles of the Declaration of Helsinki. All patients gave informed 136 written consent.

137 Clinical and pathological data, including sex, age at diagnosis, date of diagnosis, tumor stage 138 according to the ENSAT staging system [22], hormone secretion, Weiss score [21], Ki67 139 proliferation index, mitotane plasma concentrations after three and six months and response 140 to treatment during follow-up were either provided by the participant center or collected 141 through the ENSAT registry (https://registry.ensat.org).

#### 142 Chromogenic immunohistochemistry

Full FFPE sections mounted on slides were deparaffinised, rehydrated and antigen retrieval was performed in 10mM citric acid monohydrate buffer (pH 6.5) under pressure for 13 min. Blocking of unspecific binding sites occurred with 20% human AB serum at room temperature (RT) for 1 h and the primary antibody (SOAT1; ab39327 Abcam) was incubated in a 1:1000 dilution for 1h at RT as previously described [15]. The N-Universal negative control anti-rabbit (Dako) was used and signal amplification was achieved by the Advance HRP Link Kit for 40 min and developed for 10 min with the DAB+ Liquid Kit (Dako). Nuclei were counterstained using Mayer's hematoxylin for 3 min and blued for 5 min in running tap water. To ensure specificity of the antibody used [23], we overexpressed human SOAT1 in ACC cells which resulted in an increase of both detected SOAT1 bands and SOAT1 WB of 5 normal adrenal glands also resulted only in the two specific bands (Fig. S1).

#### 154 Semi-quantitative analysis of SOAT1 immunoreactivity

155 Chromogenic staining intensities were determined by two independent investigators (I.W. 156 and B.A. or L.-S.L.) and graded as 0 (negative), 1 (low), 2 (medium) and 3 (high). The 157 proportion of positive tumor cells was calculated for each slide and scored 0 if 0% were 158 positive, 0.1 if 1-9% were positive, 0.5 if 10-49% were positive and 1 if ≥50% were positive 159 [24, 25]. A semi guantitative H-Score was then calculated by multiplying the staining intensity 160 grading score with the proportion score. Where discrepancies were observed, results were 161 jointly assessed by both investigators and the final score was formed by consensus. The 162 Spearman's correlation for inter-observer agreement for each staining was high (r>0.85).

#### 163 Statistical analysis

164 RFS and PFS were considered as the time between diagnosis and documented recurrence 165 and progression (based on cross sectional imaging), respectively. DSS was calculated from 166 the time of diagnosis until disease-related death or censored at last follow-up. RFS, PFS and 167 DSS were analysed using the Kaplan-Meier method and groups were compared by using 168 the log-rank test. Assessment of prognostic factors (ENSAT stage, ki67, age, sex, hormone 169 secretion and for the group with advanced disease additionally: preM-TTP (pre mitotane time 170 to progression= time between diagnosis and progress before initiation of mitotane treatment) 171 was performed with the Cox proportional hazard regression model. The Chi-square test was 172 used to investigate dichotomic variables, whereas non-parametric Kruskal-Wallis s test was 173 used for comparison among groups for non-normal distributed variables. Correlations 174 between H-Score and prognostic factors were evaluated by Spearman's correlation. P 175 values <0.05 were considered statistically significant. Statistical analyses were performed 176 with IBM SPSS Version 23 and GraphPad Prism Version 6.

#### 177 Results

#### 178 Patient characteristics

179 Clinical characteristics of 231 ACC patients are summarised in Table 1. Median age at 180 diagnosis was 54.2 years (range 17-83) in the adjuvant group and 51 years (range 16-80) in 181 the group with disease. In both groups, approximately 60% of the patients were female and 182 40% were male. At diagnosis, the majority of patients treated with mitotane monotherapy in 183 the adjuvant setting had an ENSAT tumor stage of I-II (62.3%), whereas, in the advanced 184 stage setting, most of the patients had a tumor stage of IV (55.6%). The remaining patients 185 with advanced disease had a localized tumor at diagnosis and started mitotane therapy only 186 after developing local recurrence or metastases. Data regarding Ki67 index were available in 187 91.2% and 83.5% of patients in the adjuvant and advanced stage setting, respectively. 31 188 patients (21%) of the adjuvant group and 18 patients (27.3%) of the advanced stage group 189 had Ki67 index staining below 10% (p=0.35, chi-square=0.88). Median Weiss score was 6 190 (range 1-9) in both groups. In both arms, about 70% of the tumors were hormonally active. 191 Median time to start mitotane were one month in the adjuvant group and less than one month 192 in the group with advanced disease. Median mitotane plasma levels at three months of 193 therapy were 9.3 mg/l and 10 mg/l, after six months 13.5 mg/l and 12.8 mg/l in the adjuvant 194 and advanced stage cohort, respectively. In the advanced stage group, preM-TTP was <365 195 days in 51/63 patients (81%) for DSS and <365 days in 52/67 patients (78%) for PFS.

196 No recurrence was observed in 74/135 patients within a median follow-up of 18.5 months 197 (range 1-216 months) in patients treated in adjuvant setting. Best response to advanced 198 stage mitotane was complete (n=1) or partial response in 9, stable disease in 13 and 199 progressive disease in 45 patients. Median follow up of patients still alive (n=18) was 19.5 200 months (range 2-180 months) in this setting.

#### 201 SOAT1 expression and correlation with known prognostic factors of ACC

202 Tissue SOAT1 expression differed widely in tumors of both the adjuvant and the group with 203 advanced disease and exhibited different intra-tumoral patterns between homogeneous and 204 heterogeneous staining intensity (Fig. 1). Semiquantitative H-score accounts for this 205 heterogeneity as it takes into account both the staining intensity and percentage of cells 206 being stained and ranged from 0 to 3. Scores from 0 to <2 were designated low expression 207 (Fig. 1J-L) while scores ≥2 were indicative of high expression (Fig. 1A-I). No difference in 208 SOAT1 expression was found between hormone producing and endocrine inactive ACC with 209 mean staining intensities of  $1.53 \pm 0.9$  in inactive vs.  $1.48 \pm 0.9$  in hormonally active ACC, 210 p=0.76. No correlation of SOAT1 H-score was observed with Ki67, ENSAT stage, Weiss 211 score and age at diagnosis neither in the adjuvant, nor in the advanced stage setting.

#### 212 SOAT1 expression as factor of survival and response to mitotane treatment in ACC

213 In the adjuvant setting (Fig. 2A), we did not observe significant differences of RFS between 214 ACC patients with low SOAT1 expression in comparison to those with high SOAT1 215 expression (median 22 months, range 1-153 vs. median 12 months, range 1.5-216 log rank 216 p=0.12). When we only included patients with Ki67≥10% to analyse RFS, we did not observe 217 significant differences between SOAT1 low and high expressing ACC either (log rank 218 p=0.73). DSS (Fig. 2B) did not significantly differ between patients whose tumors expressed 219 low levels of SOAT1 compared to those with high SOAT1 expression (median 51 months, 220 range 1-252 vs. 31 months, range 2-216 log rank p=0.23). Similarly, in the group with 221 advanced disease, no significant difference in PFS (Fig. 2C) between patients with low

222 SOAT1 expression and those with high SOAT1 expression (median PFS 5 months, range 1-223 59 vs. median 4 months, range 1-25 log rank p= 0.66) was observed. Median DSS (Fig. 2D) 224 was likewise not different in tumors with low vs. high SOAT1 (median 22 months, range 4-180 vs. 21 months, 2-83 months, log rank p=0.47). When we analysed all patients together 225 226 (Fig. S2A), low SOAT1 expression was associated with a significantly longer median 227 recurrence-/progression-free survival of 13 months (range 1 -153 months vs 8 months (range 228 1-216 months, log rank p=0.049). We did not observe a significant difference in DSS (Fig. 229 S2B) between tumors with low SOAT1 vs high SOAT1 expression (median: 41 months, 230 range 1 -252 vs. median: 28 months, range 2-216, log rank p=0.41).

The proportion of tumors with low and high SOAT1 expression did not differ between patients in the adjuvant cohort without recurrence (low, n=44; high, n=30) and with recurrence (low, n=35; high, n=26) (Fig. 3A). Similarly, in the cohort with advanced disease, there were no differences between tumors with low and high SOAT1 regarding objective response to mitotane (low, n=6; high, n=4) *vs.* stable disease (low, n=6; high, n=7) and progressive disease (low, n=25; high, n=20), respectively (Fig. 3B).

We next aimed at multivariable adjustment for known clinical/histopathological ACC prognostic factors. In the adjuvant arm, univariate analysis revealed only a Ki67-Index <10% as significantly associated with improved DSS and RFS (Table 2). In patients with advanced disease the following factors were significantly associated with improved DSS: male sex, Ki67-Index <10% and preM-TTP >365 days. After multivariate analysis of all factors, including SOAT1 expression, only preM-TTP >365 days retained statistical significance (Table 3).

#### 244 **SOAT1** expression is not related to mitotane plasma concentrations

245 We next examined the potential association of SOAT1 expression with mitotane plasma 246 concentrations. Mitotane plasma levels after three months of treatment did not significantly 247 differ between patients whose tumors showed high vs low expression of SOAT1 both in the 248 adjuvant (median mitotane levels:10.3 mg/l vs 9.1 mg/l) and in the advanced disease setting 249 (median mitotane levels: 11.7 mg/l vs 9.1 mg/l) (Fig. 4A). SOAT1 expression was not 250 associated with mitotane plasma concentrations above 14 mg/l neither in the adjuvant (Fig. 251 4B) nor in the advanced disease arm (Fig. 4C). Similar results were observed after six 252 months of mitotane treatment (median mitotane levels 14.2 mg/l vs 13 mg/l in the adjuvant 253 group and 11.9 mg/l vs 12.8 mg/l in the group with advanced disease). When analyzing only 254 patients reaching the mitotane target level of 14 mg/l after three months, significantly fewer 255 patients with high SOAT1 expression responded to therapy (Fig. 4D) while this difference 256 was no longer observed when considering the six months time point (Fig. 4E).

257 Median dose of mitotane intake was 4 g/daily (range 1-12 g) in the adjuvant arm and 5 258 g/daily (range 2-12 g/daily) in patients treated for advanced disease and did not significantly 259 differ between the SOAT1 high and low expressing group (p=0.6 (adjuvant) and p=0.4260 (advanced disease)).

#### 261 Discussion

Mitotane is the only approved drug for the treatment of ACC, however, objective response rates are only approximately 20% [5, 6]. In addition to its limited therapeutic potential, adverse events occur frequently and reliable markers predicting response to therapy are currently not established. Therefore, it is crucial to define a particular subgroup of patients that will take advantage from treatment and to avoid toxicity in patients unlikely to respond.

267 At present, this topic has been addressed only in a limited number of patients [13, 14] and 268 very recently a study demonstrated mitotane sensitivity only in a very specific sub-group of 269 patients [26]. Although mitotane has been used in the clinic for decades, its precise 270 mechanism of action and molecular target remained unknown for decades, despite intense 271 research including several different "omics" approaches [18-20, 27]. We demonstrated that 272 mitotane inhibits SOAT1, leading to ER-stress and cell death of adrenocortical cells [15]. It 273 was also shown that SOAT1 is predominantly expressed in adrenocortical cells, compared to 274 cells of non-adrenal origin [15], possibly explaining the specific adrenolytic toxicity of 275 mitotane. In addition, in glioblastoma, inhibition of SOAT1 has been proposed as a novel 276 treatment [28, 29].

277 In hepatocellular carcinoma high SOAT1 expression was associated with a worse prognosis 278 [30] and has previously been described in prostate cancer as well [31]. An adverse outcome 279 of SOAT1 expression in ACC was recently demonstrated [32]. These results suggest that the 280 elevated expression of SOAT1 could be a prognostic feature of diverse cancers. In a small 281 single center series of patients (n=25) with advanced ACC [15], we had previously shown 282 that SOAT1 expression is associated with improved progression-free survival. This ENSAT 283 multicenter retrospective study aimed at validating the value of SOAT1 as a histologic marker 284 for mitotane response. Our results disprove our initial hypothesis, as no significant 285 differences in response to mitotane treatment could be observed between ACC tissue 286 samples with high and low levels of SOAT1 protein neither in an adjuvant setting nor in 287 patients treated with advanced disease.

Our study has the strength of a large collection of tissue samples from specialized ACC centers. SOAT1 expression was histologically determined in a centralized manner. All patients received mitotane monotherapy, no additional therapies were used during mitotane treatment. However, our study has several limitations. First, the clinical data and samples 292 collection were retrospectively retrieved from twelve different ENSAT centers (11 European 293 and one from Brazil) which likely is associated with different treatment strategies. This not 294 only comprises surgery and medical treatment but also documentation and follow-up. 295 Second, mitotane treatment itself is cumbersome and different dosing regimens are in use at 296 different centers [33-35]. In addition, patient-specific factors that are only partially understood 297 lead to a high heterogeneity of mitotane plasma concentrations [36-38]. Accordingly, 298 mitotane plasma concentrations in our cohort after three and six months of treatment were 299 highly variable. When considering only patients who reached mitotane plasma 300 concentrations of >14 mg/L at three or six months, SOAT1 expression was not correlated to 301 clinical response.

302 The lack of an association of SOAT1 expression with survival endpoints and response 303 implicates that additional target molecules different from SOAT1 may be relevant for its toxic 304 effect in adrenal cortical cells. In vitro, SOAT1 expression was shown to not be a predictor as 305 demonstrated in few ACC primary cultures [23] which would support the theory that 306 additional targets might be of greater importance. One such potential mechanism includes 307 inhibition of mitochondrial respiratory chain. The novel compound nevanimibe (previously 308 known as ATR101) which has been developed as a new treatment for ACC has been shown 309 to be a potent SOAT1 inhibitor by one group [39] but was also shown to inhibit mitochondrial 310 respiration by a different group [40] similar to mitotane.

Importantly, we found pronounced heterogeneity of SOAT1 expression in approximately 20 % of tumor samples. It is conceivable that this tissue heterogeneity was not completely accounted for in the monocentric study by Ferreira Lacombe *et al.* [32] in which a tissue microarrays were used to evaluate SOAT1 expression whereas we used full sections. Relationship of SOAT1 with Ki67 index and cortisol secretion was demonstrated in the previous study but not in ours. However, in our study ki67 value was provided by the various participating centers and thus a uniform analysis of this index is not guaranteed.

318 In an adjuvant setting, several other known factors such as resection status or Ki67 index 319 [41], are important to predict tumor recurrence, since even after complete resection, 320 recurrence rates are high [42-44]. In line with previous studies, Ki67-index below 10% (Table 321 2) was significantly associated with a better DSS and TTP in our cohort of patients treated 322 with mitotane in this setting. Similarly, in advanced ACC, Ki67 index, mutational burden [45] 323 but also clinical factors like age or presence of symptoms, have been identified [46, 47] to 324 predict patient outcome independently of mitotane treatment [48]. In our cohort of patients 325 with advanced disease, mitotane monotherapy, Ki67-Index below 10% was also associated 326 with a better DSS (Table 3), which retained significance after multivariate adjustment but was 327 not observed for TTP in a univariate analysis (Table 3). This may be due to the relatively

- small cohort but is in line with a previous study in which only the DSS, but not the TTPcorrelated with a Ki67-Index below 10% in advanced ACC [5].
- In conclusion, in this multicenter study, we could not confirm SOAT1 expression to be aclinically useful marker to predict treatment response to mitotane.
- 332

#### 333 Contributors:

- The following scientists contributed tissue samples and clinical data in addition to those listed as co-authors:
- 336 Marcus Quinkler (Berlin), Masanori Murakami (Munich), Felix Beuschlein (Munich, Zurich),
- Harm R. Haak (Eindhoven), Marcin Motyka (Krakow), Vasileois Chortis (Birmingham) and
- 338 Annamaria Colao (Naples).

#### 339 Acknowledgment

- 340 We are grateful to Michaela Haaf and Martina Zink for sample and data administration and
- 341 Kerstin Lenz for expert technical assistance.

342

#### 343 References

- 3441.Fassnacht, M., M. Kroiss, and B. Allolio, Update in adrenocortical carcinoma. J Clin Endocrinol345Metab, 2013. **98**(12): p. 4551-64.
- Fassnacht, M., et al., European Society of Endocrinology Clinical Practice Guidelines on the Management of Adrenocortical Carcinoma in Adults, in collaboration with the European Network for the Study of Adrenal Tumors. Eur J Endocrinol, 2018.
- 3493.Schteingart, D.E., et al., Management of patients with adrenal cancer: recommendations of350an international consensus conference. Endocr Relat Cancer, 2005. 12(3): p. 667-80.
- Reidy-Lagunes, D.L., et al., *Complete Responses to Mitotane in Metastatic Adrenocortical Carcinoma-A New Look at an Old Drug.* Oncologist, 2017. 22(9): p. 1102-1106.
- 353 5. Megerle, F., et al., *Mitotane Monotherapy in Patients With Advanced Adrenocortical*354 *Carcinoma.* J Clin Endocrinol Metab, 2018. **103**(4): p. 1686-1695.
- 3556.Terzolo, M., et al., Adjuvant mitotane treatment for adrenocortical carcinoma. N Engl J Med,3562007. **356**(23): p. 2372-80.
- Berruti, A., et al., Long-Term Outcomes of Adjuvant Mitotane Therapy in Patients With
   Radically Resected Adrenocortical Carcinoma. J Clin Endocrinol Metab, 2017. 102(4): p. 1358 1365.
- 360 8. Fassnacht, M. and B. Allolio, *Clinical management of adrenocortical carcinoma*. Best Pract
  361 Res Clin Endocrinol Metab, 2009. 23(2): p. 273-89.
- 362 9. Else, T., et al., *Adrenocortical carcinoma*. Endocr Rev, 2014. **35**(2): p. 282-326.
- 36310.Daffara, F., et al., Prospective evaluation of mitotane toxicity in adrenocortical cancer364patients treated adjuvantly. Endocr Relat Cancer, 2008. **15**(4): p. 1043-53.
- Hermsen, I.G., et al., *Plasma concentrations of o,p'DDD, o,p'DDA, and o,p'DDE as predictors of tumor response to mitotane in adrenocortical carcinoma: results of a retrospective ENS@T multicenter study.* J Clin Endocrinol Metab, 2011. **96**(6): p. 1844-51.
- Terzolo, M., et al., *Mitotane levels predict the outcome of patients with adrenocortical carcinoma treated adjuvantly following radical resection.* Eur J Endocrinol, 2013. 169(3): p.
   263-70.
- 37113.Ronchi, C.L., et al., CYP2W1 is highly expressed in adrenal glands and is positively associated372with the response to mitotane in adrenocortical carcinoma. PLoS One, 2014. **9**(8): p. e105855.
- 373 14. Volante, M., et al., *Ribonucleotide reductase large subunit (RRM1) gene expression may*374 *predict efficacy of adjuvant mitotane in adrenocortical cancer*. Clin Cancer Res, 2012. 18(12):
  375 p. 3452-61.
- Sbiera, S., et al., *Mitotane Inhibits Sterol-O-Acyl Transferase 1 Triggering Lipid-Mediated Endoplasmic Reticulum Stress and Apoptosis in Adrenocortical Carcinoma Cells.* Endocrinology, 2015. **156**(11): p. 3895-908.
- 379 16. Lee, H.T., et al., Inhibitors of acyl-CoA:cholesterol O-acyltransferase (ACAT) as
  380 hypocholesterolemic agents: synthesis and structure-activity relationships of novel series of
  381 sulfonamides, acylphosphonamides and acylphosphoramidates. Bioorg Med Chem Lett, 1998.
  382 8(3): p. 289-94.
- 38317.Smith, D.C., et al., A phase 1 study of nevanimibe HCl, a novel adrenal-specific sterol O-384acyltransferase 1 (SOAT1) inhibitor, in adrenocortical carcinoma. Invest New Drugs, 2020.
- Hescot, S., et al., *Mitotane alters mitochondrial respiratory chain activity by inducing cytochrome c oxidase defect in human adrenocortical cells.* Endocr Relat Cancer, 2013. 20(3):
  p. 371-81.
- 388 19. Poli, G., et al., *Morphofunctional effects of mitotane on mitochondria in human*389 adrenocortical cancer cells. Endocrine-Related Cancer, 2013. 20(4): p. 537-550.
- Hescot, S., et al., *Identifying mitotane-induced mitochondria-associated membranes dysfunctions: metabolomic and lipidomic approaches.* Oncotarget, 2017. 8(66): p. 109924 109940.
- 39321.Weiss, L.M., L.J. Medeiros, and A.L. Vickery, Jr., Pathologic features of prognostic significance394in adrenocortical carcinoma. Am J Surg Pathol, 1989. 13(3): p. 202-6.

395 22. Fassnacht, M., et al., Limited prognostic value of the 2004 International Union Against Cancer 396 staging classification for adrenocortical carcinoma: proposal for a Revised TNM Classification. 397 Cancer, 2009. 115(2): p. 243-50. 398 23. van Koetsveld, P.M., et al., The Efficacy of Mitotane in Human Primary Adrenocortical 399 Carcinoma Cultures. J Clin Endocrinol Metab, 2019. 400 24. Olaussen, K.A., et al., DNA repair by ERCC1 in non-small-cell lung cancer and cisplatin-based 401 adjuvant chemotherapy. N Engl J Med, 2006. 355(10): p. 983-91. 402 25. Weigand, I., et al., Differential expression of the protein kinase A subunits in normal adrenal 403 glands and adrenocortical adenomas. Sci Rep, 2017. 7(1): p. 49. 404 26. Altieri, B., et al., Effects of Germline CYP2W1\*6 and CYP2B6\*6 Single Nucleotide 405 Polymorphisms on Mitotane Treatment in Adrenocortical Carcinoma: A Multicenter ENSAT 406 Study. Cancers (Basel), 2020. 12(2). 407 27. Zsippai, A., et al., Effects of mitotane on gene expression in the adrenocortical cell line NCI-408 H295R: a microarray study. Pharmacogenomics, 2012. 13(12): p. 1351-61. 409 28. Geng, F., et al., Inhibition of SOAT1 Suppresses Glioblastoma Growth via Blocking SREBP-1-410 Mediated Lipogenesis. Clin Cancer Res, 2016. 22(21): p. 5337-5348. 411 29. Geng, F. and D. Guo, Lipid droplets, potential biomarker and metabolic target in 412 glioblastoma. Intern Med Rev (Wash D C), 2017. 3(5). 413 Jiang, Y., et al., Proteomics identifies new therapeutic targets of early-stage hepatocellular 30. 414 carcinoma. Nature, 2019. 567(7747): p. 257-261. 415 Stopsack, K.H., et al., Cholesterol uptake and regulation in high-grade and lethal prostate 31. 416 cancers. Carcinogenesis, 2017. 38(8): p. 806-811. 417 32. Lacombe, A.M.F., et al., Sterol O-Acyl Transferase 1 as a Prognostic Marker of Adrenocortical 418 Carcinoma. Cancers (Basel), 2020. 12(1). 419 33. Kerkhofs, T.M., et al., Comparison of two mitotane starting dose regimens in patients with 420 advanced adrenocortical carcinoma. J Clin Endocrinol Metab, 2013. 98(12): p. 4759-67. 421 34. Mauclere-Denost, S., et al., High-dose mitotane strategy in adrenocortical carcinoma: 422 prospective analysis of plasma mitotane measurement during the first 3 months of follow-up. 423 Eur J Endocrinol, 2012. 166(2): p. 261-8. 424 35. Baudin, E., et al., Impact of monitoring plasma 1,1-dichlorodiphenildichloroethane (o,p'DDD) 425 levels on the treatment of patients with adrenocortical carcinoma. Cancer, 2001. 92(6): p. 426 1385-92. 427 36. Kerkhofs, T.M., et al., Development of a pharmacokinetic model of mitotane: toward 428 personalized dosing in adrenocortical carcinoma. Ther Drug Monit, 2015. **37**(1): p. 58-65. 429 37. Arshad, U., et al., *Enzyme autoinduction by mitotane supported by population* 430 pharmacokinetic modelling in a large cohort of adrenocortical carcinoma patients. Eur J 431 Endocrinol, 2018. 179(5): p. 287-297. 432 38. Puglisi, S., et al., Mitotane Concentrations Influence the Risk of Recurrence in Adrenocortical 433 Carcinoma Patients on Adjuvant Treatment. J Clin Med, 2019. 8(11). 434 39. LaPensee, C.R., et al., ATR-101, a Selective and Potent Inhibitor of Acyl-CoA Acyltransferase 1, 435 Induces Apoptosis in H295R Adrenocortical Cells and in the Adrenal Cortex of Dogs. 436 Endocrinology, 2016. 157(5): p. 1775-88. 437 40. Cheng, Y., R.E. Kerppola, and T.K. Kerppola, ATR-101 disrupts mitochondrial functions in 438 adrenocortical carcinoma cells and in vivo. Endocr Relat Cancer, 2016. 23(4): p. 1-19. 439 41. Beuschlein, F., et al., Major prognostic role of Ki67 in localized adrenocortical carcinoma after 440 complete resection. J Clin Endocrinol Metab, 2015. 100(3): p. 841-9. 441 42. Luton, J.P., et al., Clinical features of adrenocortical carcinoma, prognostic factors, and the 442 effect of mitotane therapy. N Engl J Med, 1990. 322(17): p. 1195-201. 443 43. Fassnacht, M., et al., Adrenocortical carcinoma: a clinician's update. Nat Rev Endocrinol, 444 2011. 7(6): p. 323-35. 445 44. Scollo, C., et al., Prognostic Factors for Adrenocortical Carcinoma Outcomes. Front Endocrinol 446 (Lausanne), 2016. 7: p. 99.

447	45.	Lippert, J., et al., Targeted Molecular Analysis in Adrenocortical Carcinomas: A Strategy
448		Toward Improved Personalized Prognostication. J Clin Endocrinol Metab, 2018. 103(12): p.
449		4511-4523.
450	46.	Zhang, F., et al., Prognostic Role of Ki-67 in Adrenocortical Carcinoma After Primary
451		Resection: A Retrospective Mono-Institutional Study. Adv Ther, 2019. 36(10): p. 2756-2768.
452	47.	Libe, R., Clinical and molecular prognostic factors in adrenocortical carcinoma. Minerva
453		Endocrinol, 2019. <b>44</b> (1): p. 58-69.
454	48.	Libe, R., et al., Prognostic factors in stage III-IV adrenocortical carcinomas (ACC): an European
455		Network for the Study of Adrenal Tumor (ENSAT) study. Ann Oncol, 2015. 26(10): p. 2119-25.
456		

#### **Table 1: Patient characteristics.**

	adjuvant
Patients n	158
Age in years: median (range)	54 2 (17-83)
Sex - n (%)	01.2 (11 00)
female	97 (61)
male	61 (39)
Tumor stage - n (%)	
I	10 (6.5)
II	88 (56)
III	52 (33)
IV	7 (4.5)
R status (n=156)	
0	121 (77.6)
Х	30 (19.2)
1	5 (3.2)
Ki67-Index (n=144) - n (%)	
<10%	30 (21)
≥10%	114 (79)
Weiss score (n=142): median (range)	6 (1-9)
Endocrine activity (n=143) - n (%)	
hormone secretion	96 (67)
no over-secretion	47 (33)
Months to mitotane start (median (IQR)) Mitotane plasma concentration at 3 months (median ±SD) (n=132)	1 (2)
	9.05 ±5.9
Mitotane plasma concentration at 6 months (median $\pm$ SD) (n=125)	13 ±5.7

	advanced disease
Patients n	72
Age in years: median (range)	51 (16-80)
female	42 (58)
male	30 (42)
Tumor stage- n (%)	
I	2 (2.8)
II	19 (26.4)
III	11 (15.3)
IV	40 (55.6)
R status (n=68)	
0	28 (41)
Х	15 (22)
1	13 (19)
2	12 (18)
Ki67-Index (n=66) - n (%)	
<10%	18 (27.3)
≥10%	48 (66.4)
Weiss score (n=55): median (range) Endocrine activity (n=62) - n (%)	6 (1-9)
hormone secretion	45 (72.6)
no over-secretion	17 (27.4)
Months to mitotane start (median (IQR)) Mitotane level at 3 months (median ±SD)	0 (2)
(n=61) Mitotane level at 6 months (median ±SD)	10±6.3
(n=51)	12.5±7

## 461Table 2: Impact of SOAT1 expression and known prognostic parameters on RFS and462DSS in the adjuvant (R0 or RX) cohort.

<u>RFS</u>	univariate analysis				multivariate analysis				
variables	HR	95% CI	р		HR	95% CI	р		
Sex									
female (n=91)									
male (n=56)	1.13	0.72-1.76	0.60		1.30	0.77-2.19	0.33		
Age									
<50 (n=90)									
≥50 (n=58)	0.78	0.49-1.23	0.29		0.72	0.40-1.29	0.27		

Hormone over- secretion Yes (n=90) No (n=43)	1.39	0.84-2.32	0.20	1.55	0.86-2.77	0.14
Tumor stage						
l+ II (n=90)						
III + IV (n=56)	1.51	0.97-2.34	0.07	1.54	0.90-2.62	0.11
Ki67						
<10 (n=28)						
≥10 (n=107) Mitotane levels 3 months (median:9.1 mg/l) n=122	3.810	1.64-8.84	0.002*	2.86	1.18-6.96	0.02
<9.1		0.92-				
≥9.1 Mitotane levels 6 months (median:13 mg/l) n=116	1.50	2.44	0.11	-	-	-
<13		0.62 -				
≥13	1.02	1.67	0.95	-	-	-
SOAT1 H-Score low: <2 (n=89) H-Score high: >2						
(n=59)	1.42	0.91-2.21	0.12	1.07	0.61-1.85	0.82
DSS	un	ivariate analy	sis	r	nultivariate ar	nalysis
<u>DSS</u>	un HR	ivariate analy 95% Cl	sis	r HR	nultivariate ar 95% Cl	nalysis p
DSS variables Sex	un HR	ivariate analy 95% Cl	sis	r HR	nultivariate ar 95% Cl	nalysis p
DSS variables Sex female (n=81)	un HR	ivariate analy 95% Cl	sis	r HR	nultivariate ar 95% Cl	nalysis p
<u>DSS</u> variables Sex female (n=81) male (n=53)	un HR 1.19	ivariate analy 95% Cl 0.462- 2.28	<u>sis p</u> 0.61	<u>HR</u>	nultivariate ar 95% Cl 0.74-3.67	nalysis p 0.22
<u>DSS</u> variables Sex female (n=81) male (n=53) Age <50 (n=80)	un HR 1.19	ivariate analy 95% Cl 0.462- 2.28	<u>sis</u>  0.61	r  1.65	nultivariate ar 95% Cl 0.74-3.67	nalysis p 0.22
<u>DSS</u> variables Sex female (n=81) male (n=53) Age <50 (n=80) ≥50 (n=55)	un HR 1.19	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29	<u>sis p</u> 0.61	<u>HR</u>	nultivariate ar 95% Cl 0.74-3.67 0.24-1.55	nalysis p 0.22 0.29
<u>DSS</u> variables Sex female (n=81) male (n=53) Age <50 (n=80) ≥50 (n=55)	<u>un</u> HR 1.19 0.64	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29	<u>sis</u> <u>p</u> 0.61 0.21	<u>HR</u> 1.65 0.60	nultivariate ar 95% Cl 0.74-3.67 0.24-1.55	nalysis p 0.22 0.29
DSS variables Sex female (n=81) male (n=53) Age <50 (n=80) ≥50 (n=55) Hormone over- secretion	un HR 1.19 0.64	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29	<u>sis</u> <u>p</u> 0.61 0.21	<u>HR</u> 1.65 0.60	nultivariate ar 95% Cl 0.74-3.67 0.24-1.55	nalysis p 0.22 0.29
$\frac{DSS}{variables}$ Sex female (n=81) male (n=53) Age <50 (n=80) $\geq$ 50 (n=55) Hormone over- secretion Yes (n=85)	un HR 1.19 0.64	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29	sis p 0.61 0.21	<u>HR</u> 1.65 0.60	nultivariate ar 95% Cl 0.74-3.67 0.24-1.55	nalysis p 0.22 0.29
$\frac{DSS}{variables}$ Sex female (n=81) male (n=53) Age <50 (n=80) $\geq$ 50 (n=55) Hormone over- secretion Yes (n=85) No (n=36)	un HR 1.19 0.64 1.52	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29 0.68-3.40	<u>sis</u> <u>p</u> 0.61 0.21 0.31	<u>HR</u> 1.65 0.60 1.48	nultivariate ar 95% CI 0.74-3.67 0.24-1.55 0.58-3.79	nalysis p 0.22 0.29 0.42
$\frac{DSS}{variables}$ Sex female (n=81) male (n=53) Age <50 (n=80) ≥50 (n=55) Hormone over- secretion Yes (n=85) No (n=36) Tumor stage	un HR 1.19 0.64 1.52	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29 0.68-3.40	<u>sis</u> <u>p</u> 0.61 0.21 0.31	<u>HR</u> 1.65 0.60 1.48	nultivariate ar 95% Cl 0.74-3.67 0.24-1.55 0.58-3.79	nalysis p 0.22 0.29 0.42
$\frac{DSS}{variables}$ Sex female (n=81) male (n=53) Age <50 (n=80) $\geq$ 50 (n=55) Hormone over- secretion Yes (n=85) No (n=36) Tumor stage I + II (n=85)	un HR 1.19 0.64 1.52	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29 0.68-3.40	sis p 0.61 0.21 0.31	<u>HR</u> 1.65 0.60 1.48	nultivariate ar 95% Cl 0.74-3.67 0.24-1.55 0.58-3.79	nalysis p 0.22 0.29 0.42
$\frac{DSS}{variables}$ Sex female (n=81) male (n=53) Age <50 (n=80) $\geq$ 50 (n=55) Hormone over- secretion Yes (n=85) No (n=36) Tumor stage I + II (n=85) III + IV (n=48)	un HR 1.19 0.64 1.52 1.43	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29 0.68-3.40 0.74-2.76	<u>sis</u> <u>p</u> 0.61 0.21 0.31 0.28	<u>HR</u> 1.65 0.60 1.48 1.23	nultivariate ar 95% CI 0.74-3.67 0.24-1.55 0.58-3.79 0.54-2.78	nalysis p 0.22 0.29 0.42 0.63
$\frac{DSS}{variables}$ Sex female (n=81) male (n=53) Age <50 (n=80) $\geq$ 50 (n=55) Hormone over- secretion Yes (n=85) No (n=36) Tumor stage I + II (n=85) III + IV (n=48) Ki67	un HR 1.19 0.64 1.52 1.43	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29 0.68-3.40 0.74-2.76	<u>sis</u> <u>p</u> 0.61 0.21 0.31 0.28	r HR 1.65 0.60 1.48 1.23	nultivariate ar 95% Cl 0.74-3.67 0.24-1.55 0.58-3.79 0.54-2.78	nalysis p 0.22 0.29 0.42 0.63
$\frac{DSS}{variables}$ Sex female (n=81) male (n=53) Age <50 (n=80) $\geq$ 50 (n=55) Hormone over- secretion Yes (n=85) No (n=36) Tumor stage I + II (n=85) III + IV (n=48) Ki67 <10 (n=24)	un HR 1.19 0.64 1.52 1.43	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29 0.68-3.40 0.74-2.76	<u>sis</u> p 0.61 0.21 0.31 0.28	r HR 1.65 0.60 1.48 1.23	nultivariate ar 95% Cl 0.74-3.67 0.24-1.55 0.58-3.79 0.54-2.78	nalysis p 0.22 0.29 0.42 0.63
$\frac{DSS}{variables}$ Sex female (n=81) male (n=53) Age <50 (n=80) $\geq$ 50 (n=55) Hormone over- secretion Yes (n=85) No (n=36) Tumor stage I + II (n=85) III + IV (n=48) Ki67 <10 (n=24)	un HR 1.19 0.64 1.52 1.43	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29 0.68-3.40 0.74-2.76	sis p 0.61 0.21 0.31 0.28	r HR 1.65 0.60 1.48 1.23	nultivariate ar 95% Cl 0.74-3.67 0.24-1.55 0.58-3.79 0.54-2.78	nalysis p 0.22 0.29 0.42 0.63
$\frac{DSS}{variables}$ Sex female (n=81) male (n=53) Age <50 (n=80) $\geq$ 50 (n=55) Hormone over- secretion Yes (n=85) No (n=36) Tumor stage I + II (n=85) III + IV (n=48) Ki67 <10 (n=24) $\geq$ 10 (n=99)	un HR 1.19 0.64 1.52 1.43 4.91	ivariate analy 95% Cl 0.462- 2.28 0.32-1.29 0.68-3.40 0.74-2.76 1.17- 20.67	sis p 0.61 0.21 0.31 0.28 0.03*	 HR 1.65 0.60 1.48 1.23 3.60	nultivariate ar 95% Cl 0.74-3.67 0.24-1.55 0.58-3.79 0.54-2.78 0.80- 16.24	nalysis p 0.22 0.29 0.42 0.63 0.10

months (median:9.1						
<9.1		0.76 –				
≥9.1	1.52	3.06	0.24	-	-	-
Mitotane levels 6 months (median:13 mg/l) n=103						
≥13	0.74	0.34-1.60	0.44	-	-	-
SOAT1 H-Score low: <2 (n=81) H-Score high: ≥2						
(n=54)	1.49	0.77-2.86	0.24	1.30	0.58-2.93	0.53

#### **Table 3: Impact of SOAT1 expression and known prognostic parameters on PFS and**

**DSS** in the cohort with advanced disease. preM-TTP: pre mitotane time to progression

PFS	u	nivariate anal	ysis	multivariate analysis			
variables	HR	95% CI	р	HR	95% CI	р	
Sex							
female (n=40)							
male (n=27)	0.75	0.44-1.27	0.28	0.81	0.43-1.53	0.51	
Age							
<50 (n=29)							
≥50 (n=38)	0.84	0.50-1.42	0.52	0.73	0.36-1.50	0.40	
Hormone over-							
secretion							
Yes $(n=43)$	1.46	0 77 0 70	0.05	1 00	0.07.4.02	0.00	
No $(n=14)$	1.40	0.77-2.78	0.25	1.98	0.97-4.03	0.06-	
preM-TTP							
<365 days							
,							
≥365 days	0.37	0.18-0.72	0.004*	0.49	0.21-1.11	0.09	
Ki67							
< 10 (n - 17)							
10(n-17)	1 10	0 66 0 14	0 55	0.02	0 46 1 92	0.01	
≥10 (l1=45) Mitotane levels 3	1.19	0.00-2.14	0.55	0.92	0.40-1.83	0.81	
months							
(median:10 mg/l)							
n=58							
<10	0.70	0 40 4 00	0.04				
≥10 Mitotane levels 6	0.70	0.40-1.22	0.21	-	-	-	
months							
(median:12.5							
mg/l) n=48							

<12.5						
≥12.5	0.61	0.33-1.14	0.12	-	-	-
SOAT1						
H-Score low: <2						
(n=37)						
H-Score high: ≥2						
(n=30)	1.11	0.68-1.86	0.68	1.34	0.63-2.84	0.45

DSS	univariate analysis				multivariate analysis			
variables	HR	95% CI	р		HR	95% CI	р	
Sex								
female (n=36)								
male (n=27)	0.48	0.26-0.92	0.026*		0.92	0.40-2.11	0.83	
Age								
<50 (n=27)								
≥50 (n=36)	0.82	0.46-1.48	0.52		1.39	0.63-3.04	0.42	
Hormone over-								
secretion								
Yes (n=40)			0.00		4.00		0.07	
No (n=13)	1.04	0.52-2.07	0.92		1.20	0.52-2.80	0.67	
KI67								
<10 (n=14)								
≥10 (n=45)	2.47	1.14-5.32	0.021*		1.83	0.73-4.60	0.20	
preM-TTP		0.044						
<365 days	0.60	0.014-	~0 001*		0.10	0.02.0.40	0 004*	
~303 uays	0.00	0.257	<0.001		0.10	0.02-0.49	0.004	
Mitotane levels 3								
months								
(median:10 mg/l)								
n=54								
<10	1 05	0.56-	0.00					
∠IU Mitotane levels 6	1.05	1.98	0.88		-	-	-	
months								
(median:12.5								
mg/l) n=45								
<12.5	0.60	0.00.4.07	0.40					
C.212	0.62	0.30-1.27	0.19		-	-	-	
SUATT H-Score low: <2								
(n=35)								
H-Score high: ≥2								
(n=28)	0.81	0.44-1.46	0.48		0.72	0.31-1.70	0.45	

#### 465 Figure legends

#### 466 Figure 1: SOAT1 immunohistochemistry staining of full ACC FFPE sections. First

467 column shows an overview of SOAT1 staining intensities within the same tumors (scale bars:

- 468 3mm). Second column shows 3x magnification of the representative slide in first column
- 469 (scale bars: 700μm) and third column shows 20x magnification of the slide shown in column
- 470 A (scale bars: 200µm) (A-C: SOAT1 H-score 3, inhomogeneous staining; D-F: SOAT1 H-
- 471 Score 3, inhomogeneous staining; G-I: SOAT1 H-score 2, homogeneous staining, J-L:
- 472 SOAT1 H-score 0, homogenous staining).
- 473 Figure 2: Kaplan-Meier plots of SOAT1 low and high expressing ACC. (A) Recurrence-
- 474 /progression-free survival and **(B)** disease-specific survival of all ACC patients. **(C)**
- 475 Recurrence-free survival (D) and disease-specific survival of ACC patients in the adjuvant
- 476 group. (E) progression-free survival (F) and disease-specific survival of ACC patients with
- 477 advanced disease.
- 478 **Figure 3: SOAT1 expression and treatment response.** No significant differences
- 479 regarding mitotane response between SOAT1 high and SOAT1 low expressing tumors were
- 480 observed in the adjuvant arm (A), nor in advanced stages (B).
- Figure 4: Correlation of SOAT1 expression and mitotane plasma concentrations. (A) In both arms, high SOAT1 expression was not correlated with higher mitotane plasma levels. Patients with high SOAT1 expression are not more likely to reach mitotane plasma levels above 14 mg/l not in the adjuvant setting (B), nor in patients with advanced disease (C). When only patients reaching the mitotane target level of 14 mg/l were analysed, high SOAT1 expression was significantly correlated with higher rates of recurrences after three months (D) which did not retain significance after six months (E).