

# Predicted salivary human protease activity in experimental gingivitis revealed by endoProteo-FASP approach

Mulkern, Dimitri ; Hewitt, Amy ; Parker, Hadyn ; Batt, Joanna; Yonel, Zehra; Grant, Melissa

DOI:

[10.1111/eos.12729](https://doi.org/10.1111/eos.12729)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Mulkern, D, Hewitt, A, Parker, H, Batt, J, Yonel, Z & Grant, M 2020, 'Predicted salivary human protease activity in experimental gingivitis revealed by endoProteo-FASP approach', *European Journal of Oral Sciences*, vol. 128, no. 5, pp. 386-394. <https://doi.org/10.1111/eos.12729>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

This is the peer reviewed version of the following article: Mulkern, D, Hewitt, A, Parker, H, Batt, J, Yonel, Z, Grant, MM. Predicted salivary human protease activity in experimental gingivitis revealed by endoProteo-FASP approach. *Eur J Oral Sci* 2020; 128: 386– 394. © 2020 Eur J Oral Sci, which has been published in final form at <https://doi.org/10.1111/eos.12729>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

**Predicted Salivary Human Protease Activity In Experimental  
Gingivitis Revealed By endoProteo-FASP Approach**

Journal:	<i>European Journal of Oral Sciences</i>
Manuscript ID	EOS-11363-OA-20.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Mulkern, Dimitri; University of Birmingham, School of Dentistry Hewitt, Amy; University of Birmingham, School of Dentistry Parker, Hadyn; University of Warwick, School of Life Sciences Batt, Joanna; University of Birmingham, School of Dentistry Yonel, Zehra; University of Birmingham, School of Dentistry Grant, M; University of Birmingham, School of Dentistry
Keywords (Please write 3 to 5 keywords according to Index Medicus):	saliva, protease, gingivitis
Research Area:	Periodontology

SCHOLARONE™  
Manuscripts

1  
2  
3 **Predicted salivary human protease activity in experimental gingivitis revealed by endoProteo-**  
4 **FASP approach**  
5  
6

7 **Dimitri Mulkern 1,†, Amy Hewitt 1,†, Hadyn Parker 1,2, Joanna Batt 1, Zehra Yonel 1 and Melissa**  
8 **M. Grant 1**  
9

10  
11 <sup>1</sup> Periodontal Research Group, School of Dentistry, University of Birmingham, 5 Mill Pool  
12 Way, Edgbaston, Birmingham, B5 7EG  
13  
14

15 <sup>2</sup> School of Life Sciences, University of Warwick, Coventry, CV4 7AL, United Kingdom  
16  
17

18 <sup>†</sup> These authors contributed equally  
19  
20

21  
22  
23 **Running title:** Saliva proteases in experimental gingivitis  
24  
25  
26

27 **Corresponding author:**  
28  
29

30 Melissa Grant  
31

32 Periodontal Research Group, School of Dentistry, University of Birmingham, 5 Mill Pool Way,  
33 Edgbaston, Birmingham, B5 7EG  
34  
35

36 Telephone: +44 121 446 5520  
37  
38

39 Email: m.m.grant@bham.ac.uk  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5 *Mulkern D, Hewitt A, Parker H, Batt J, Yonel Z, Grant MM. Predicted salivary human protease*  
6 *activity in experimental gingivitis revealed by endoProteo-FASP approach. Eur J Oral Sci*  
7  
8

9 **Abstract:** Gingivitis is a highly prevalent oral condition that can be studied in humans via the 21-  
10 day experimental gingivitis model that allows for investigations into induction and resolution of  
11 gingivitis. In this study we used the autolysis of saliva as a source of peptides to predict the activity  
12 of human proteases in saliva during induction and resolution of inflammation. Healthy volunteers,  
13 with no remarkable oral or systemic conditions, were recruited into the study and stimulated  
14 saliva samples were collected at days 0, 21 and 35 of experimental gingivitis. Plaque and gingival  
15 indices were recorded to ensure clinical induction and resolution. Saliva was auto-digested at 37°C  
16 for 18 h before identification of peptides by mass spectrometry. Protease prediction was carried  
17 out using Proteasix *in silico* with the identified peptides. Comparison of day 0 to days 21 and 35  
18 showed changes in predicted protease activity. Correlation network analysis revealed that at day  
19 21 the proteases became less connected and showed a potential for a dysregulated system; by day  
20 35 the connectivity was returning towards similar conditions at day 0. This study demonstrates  
21 that changes in predicted proteases are apparent even in saliva collected from donors  
22 experiencing inflammation at three teeth.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

33  
34 **Key words:** protease, saliva, gingivitis

35  
36 **Contact author:**

37  
38 Melissa Grant,

39  
40 Periodontal Research Group, School of Dentistry, University of Birmingham, 5 Mill Pool Way,  
41 Edgbaston, Birmingham, B5 7EG

42  
43 Telephone: +44 121 446 5520

44  
45 m.m.grant@bham.ac.uk  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Periodontal disease is a very common inflammatory condition found worldwide (1). In its severest  
4 form, severe periodontitis, it affects over 10% of the population (1) and is a major cause of tooth  
5 loss. However the mild inflammation, gingivitis, is experienced by 90% of the population (2) and is  
6 easily reversible with proper tooth brushing and removal of dental plaque from the tooth surface.  
7  
8 The mouth is bathed in saliva and this fluid is rich in antimicrobial defenses such as proteins and  
9 proteases. During inflammation the composition of saliva changes (3); this is due to changes  
10 originating from the salivary glands (4) and also from the gingival tissues (5). The periodontal  
11 tissues adjacent to the plaque covered tooth surface become inflamed and this will stimulate  
12 production of local mediators, such as cytokines and chemokines, which in turn drive recruitment  
13 of leukocytes, particularly neutrophils. Fluid from these tissues enters into the mouth and thereby  
14 saliva via the gingival crevice and gingival crevicular fluid (GCF).  
15  
16

17  
18 Induction of gingivitis in the 21-day experimental gingivitis model allows for insights to be  
19 gained into human inflammation and resolution of inflammation in the oral cavity, a key mucosal  
20 surface. The first publication (6,7) of this model was in 1965 and since then it has been used to  
21 explore a wide variety of influences on inflammation, such as use of particular toothpastes (8-10).  
22 The composition of GCF has been extensively investigated for individual molecules (5,11-14) or for  
23 overarching changes using 'omic' technologies (3,15). Less information has been gathered about  
24 the changes in saliva (16) as often the model is used with only a few selected teeth being shielded  
25 from brushing, thereby allowing for contralateral teeth to be used in comparison.  
26  
27

28  
29 There has been abundant interest in the role of proteases in periodontal disease. The  
30 induction of an inflammatory response requires proteases to facilitate access of leukocytes to the  
31 site of infection. The protease network is crucial for effective responses against infection, where  
32 swift deactivation of microbial proteins or activation of host proteins by proteases can be critical,  
33 but also has a key role in wound healing (17). They are essential instigators and executioners of  
34 responses and dysregulation may lead to excessive or non-resolving inflammation. With the  
35 experimental gingivitis model the change in abundance of cathepsins (6,18), elastase (5) and  
36 matrix metalloproteases (MMPs) (10,12) have been investigated.  
37  
38

39  
40 There are more than 500 proteases (19) encoded in the human genome, making them the  
41 largest family of proteins (19), and study of their presence or activity requires a system-wide  
42 approach through degradomics. The prediction of protease sites requires knowledge of the  
43 substrates and products of protease action. There has been extensive research in this area and  
44 numerous databases exist that provide resources on proteolytic events (e.g., TopFIND (20),  
45 MEROPS (21), CutDB (22)). KLEIN *et al.* (23) developed the Proteasix tool to provide a method to  
46 predict protease activity from peptides derived from mass spectrometry data and TRINDADE *et al.*  
47 (24) developed the endoProteo-FASP technique to explore the protease activity in fluid samples,  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 such as saliva. Using the filter aided sample preparation (FASP) technique developed by  
4 WIŚNIEWSKI *et al.* (25), TRINDADE *et al.* (24) made use of the endogenous proteolytic activity of saliva  
5 samples to generate peptides that were then searched using a no enzyme search. This  
6 dramatically increases search time due to the increase in search complexity but can yield  
7 information about the non-traditional proteomics peptides in the sample. TRINDADE *et al.* explored  
8 saliva from chronic periodontitis patients vs controls donors (26) and also two healthy male donors  
9 (24). They demonstrated that degradomes were generated by autolysis of saliva over 18 or 115h at  
10 37°C and that these peptides mapped across a wide range of proteins known to be present in  
11 saliva, including those not normally identified using trypsin mediated proteomics and that  
12 protease activity could be predicted using the beta version of Proteasix. In chronic periodontitis  
13 (26) patient samples they showed that there was more predicted activity of cathepsin L and  
14 Meprin 1a but lower activities predicted for MMP12, MMP7, MMP9 and Presequence protease  
15 (PREP or PITRM1) at 115h. TRINDADE *et al.* (24) also used zymography to explore the detectable  
16 protease activity. This complementary technique separates proteins in the sample by non-reducing  
17 SDS PAGE using gelatin (or other protein) containing acrylamide gels. After electrophoresis the  
18 separated proteins are allowed to renature and if protease activity is present that can degrade  
19 gelatin areas of gel with this activity will not be stained by conventional protein staining  
20 techniques, such as Coomassie blue staining. This is due to the degradation of the gelatin  
21 incorporated within the acrylamide gel: the rest of the gel stains due to the presence of the  
22 gelatin. TRINDADE *et al.* (24) demonstrated different levels of protease activity via gelatin  
23 zymography in chronic periodontitis patients.

24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

In this study the aim was to explore, in a small number of saliva donors, the protease activity of saliva samples donated during the induction and resolution of inflammation in experimental gingivitis. The endoProteo-FASP technique was used to explore holistically the activity of proteases and shed light on the interconnected functions of proteases. This would not be possible with techniques that try to target individual proteases.

## Material and Methods

### *Recruitment and protocol*

The experimental gingivitis model previously described by CHAPPLE *et al.* (27) was used, and the study was approved by the South Birmingham Local Research Ethical Committee (LREC 2004/074). Nine non-smoking volunteers (mean age 20 yr; range 19–22, 3 males and 6 females) who had unremarkable medical histories, no known history of periodontal disease, were not undergoing orthodontic or prosthodontic appliance therapy or taking medication that may have affected results, were enrolled into the study, and informed consent was obtained. To induce gingivitis participants shielded three teeth on the opposite side from their dominant hand during brushing. These sites were the maxillary left 4, 5, and 6 (first and second premolars and first molar on the left-hand side of the upper jaw) in right-handed individuals (n = 8) and the equivalent teeth on the maxillary right side in left-handed individuals (n = 1). Contralateral sites were monitored at the corresponding teeth on the contralateral side to ensure induction and resolution of inflammation. Plaque accumulation was assessed using the Turesky modification of Quigley and Hein index (28) (PI; scores 0–5) and the gingival index (GI) of Löe (29) (scores 0–3) was used to measure gingival inflammation. A soft vinyl splint was constructed to cover five teeth, with a 5 mm clearance at each free surface from the marginal tissues. The splint was inserted gently over test teeth before brushing, to ensure against any mechanical or chemical cleaning. Participants were also asked to refrain from using mouth rinse, chewing gum, interdental cleaning aids (floss / brushes) for the duration of inflammation induction between day 0 and 21. The teeth were given a thorough prophylaxis and oral hygiene was closely monitored for 2 wk prior to the study commencing to ensure pristine gingival health at baseline. On day 21 volunteers were given a full mouth prophylaxis and recommenced brushing without using the splint, in order to resolve the experimentally induced inflammation at test sites. Fourteen days later (day 35), the final set of clinical indices was recorded.

### *Saliva collection*

At days 0, 21 and 35 each participant donated a stimulated saliva sample. Samples were collected between 9–11am on the day of collection and participants were requested not to eat or drink 2 h prior to donation. Participants were asked to rinse with water 10 min prior to donation. This sample was stimulated by rolling a sterile marble around the mouth for 5 min; saliva was expectorated into a tube placed on ice throughout the collection time. The saliva was centrifuged at 1000 *g* for 10 min to remove any particulate material and then frozen at -80°C until use.

### *Saliva processing*

On defrosting saliva was kept on ice. Protein quantity was measured using bicinchoninic acid assay (30). Saliva samples containing 150 µg protein were aliquoted into a 30 kDa molecular weight cutoff filter (Amicon; Merck-Millipore, Watford, UK) and the procedure for endoProteoFASP based on TRINDADE *et al.* (24) was followed. Briefly (Fig. S1), the saliva proteins in the Amicon filter were centrifuged at 14,000 *g* for 10 min and then ammonium bicarbonate (500µl, 50mM) was added to the filter. The filter was centrifuged again and the ammonium bicarbonate (500µl, 50mM) was replaced. This was repeated for a total of five times. After the fifth wash the Amicon filter was filled with ammonium bicarbonate (500µl, 50mM) and the filter was sealed with Parafilm and incubated at 37°C for 18 h. The resulting peptides were collected by centrifuging the Amicon filter at 14,000 *g* for 10min and then washing the filter, as before, four times with ammonium bicarbonate (500µl, 50mM). These five filtrates were combined, trifluoroacetic acid (400µl, 0.1%) was added and they were dried by vacuum centrifugation. Samples were stored at -80°C before analysis. Treatment of aliquots of saliva samples with dithiothreitol (20mM) at 100°C for 45 min completely inhibited proteolytic activity, as previously described (24).

### *Zymogram technique*

Saliva (10 µg) was mixed with non-reducing SDS sample buffer (Thermo Fisher, UK) and loaded (without reduction or heating) onto Novex 10% Zymogram Plus (1 % Gelatin) Protein gel (Thermo Fisher, Gloucester, UK). Samples were electrophoresed for 90 min at 100V before incubation in Novex Renaturing buffer (30 min, Thermo Fisher). The gels were then incubated in Novex Zymogram Developing buffer (Thermo Fisher) for 30 min before Novex Zymogram Developing buffer was exchanged for fresh Novex Zymogram Developing buffer and the gels were incubated for a further 18 h at 24 °C. Gels were then rinsed (3x 5min) in distilled water before being immersed in to Imperial Blue protein stain (Expedeon, Harston, UK) for 1 h. Gels were destained for 30 min before imaging on a GS800 imaging system (BioRad, Watford, UK). Images were then processed with Image J (31).

### *Gel band protein processing*

For identification of proteins in the bands from the zymogram gels bands not stained with Imperial Blue were excised and processed for mass spectrometry following in gel digestion as previously described (32).



### *Mass spectrometry*

UltiMate 3000 HPLC series (Dionex, Sunnyvale, CA, USA) was used for peptide concentration and separation. Samples were trapped on  $\mu$ Precolumn Cartridge, Acclaim PepMap 100 C18, 5  $\mu$ m, 100Å 300 $\mu$ m i.d. x 5mm (Dionex) and separated in Nano Series Standard Columns 75  $\mu$ m i.d. x 15 cm, packed with C18 PepMap100, 3  $\mu$ m, 100Å (Dionex). The gradient used was from 3.2% to 24% solvent B (0.1% formic acid in acetonitrile) for 30 min and then increased to 60% in 15 min. The column was washed with 80% solvent B for 15 min and equilibrated with 3.2% solvent B for another 15 min. The total run time was 75 min. Peptides were eluted directly ( $\sim 350$  nL min<sup>-1</sup>) via a Triversa Nanomate nanospray source (Advion Biosciences, Ithaca, NY, USA) into a QExactive HF (QEHF) mass spectrometer (Thermo Fisher Scientific, Gloucester, UK). The data-dependent scanning acquisition was controlled by Xcalibur 4.0 software (Thermo Fisher Scientific, Gloucester, UK). The mass spectrometer alternated between a full FT-MS scan ( $m/z$  375 – 1600) and subsequent high energy collision dissociation (HCD) MS/MS scans of the 20 most abundant ions. Survey scans were acquired in the QEHF with a resolution of 120 000 at  $m/z$  200 and automatic gain control (AGC)  $3 \times 10^6$ . Precursor ions were fragmented in HCD MS/MS with resolution set up at 60,000 and a normalized collision energy of 32. ACG target for HCD MS/MS was  $1 \times 10^5$ . The width of the precursor isolation window was 1.2  $m/z$  and only multiply-charged precursor ions were selected for MS/MS. Spectra were acquired for 60 min with dynamic exclusion time of 20s. The MS and MS/MS scans were searched against Uniprot database using Proteome Discoverer 2.2 (ThermoFisher Scientific) and the Sequest algorithm with a 1% false discovery rate (FDR) for both peptide and protein criteria and controlled by Percolator. The precursor mass tolerance was 10 ppm and the MS/MS mass tolerance was 0.02 Da. For saliva samples prepared by EndoProteoFASP no enzyme restriction was used during searching and only oxidation (M) and N terminal acetylation were set as variable modifications. For in gel digested samples trypsin enzyme was used and oxidation (M) and N terminal acetylation were set as variable modifications and cysteine carbamidomethylation on lysine and N-terminal as fixed modifications. Data have been deposited in PRIDE with accession PXD015121.

### *Data processing*

Data files from Proteome Discoverer were used to search the Proteasix (23) database via the web tool (found at <http://www.proteasix.org/>). As Proteasix returns all possible proteases at a cleavage site the data were filtered for the most probable protease per peptide terminus. To generate networks graphs Pearson correlation coefficient matrices were determined by Prism (v 6.0) (Graphpad, San Diego, CA, USA) and networks were visualised by Fruchterman-Reingold algorithm

1  
2  
3 using Gephi (v 0.9.2) (33). Data for networks were selected as having p values of  $p < 0.001$ .  
4  
5 Statistical analysis was carried out using GraphPad Prism v 6.  
6  
7

## 8 **Results**

9  
10 The nine volunteers enrolled in the 21-day experimental gingivitis study provided saliva at days 0,  
11 21 and 35. At these time points the accumulation of plaque and induction of inflammation was  
12 recorded at the three shielded teeth where plaque was expected to accumulate and the three  
13 contralateral unshielded teeth. Fig. 1 shows that there was an accumulation of plaque at day 21  
14 and that this was accompanied by an increase in inflammation as measured by the GI.  
15 Recommencing of oral hygiene measures at the shielded teeth decreased the amount of plaque  
16 and inflammation, indicating a resolution of inflammation. The saliva flow rate at each day was  
17 0.98 (SD 0.2) ml/min on day 0; 0.87 (SD 0.3) ml/min on day 21; and 0.95 (SD 0.3) ml/min day 35.  
18 Additionally the protein content was measured at each day: 1.66 (SD 0.8) mg/ml on day 0, 1.73 (SD  
19 0.8) mg/ml on day 21, and 1.66 (SD 0.5) mg/ml on day 35. No significant differences were found by  
20 one way ANOVA for either measurement.  
21  
22  
23  
24  
25  
26  
27

28 Using the endoProteo-FASP technique saliva samples from the volunteers were auto-digested  
29 for 18 h at 37°C. The resulting peptides were analysed by mass spectrometry. 784 proteins were  
30 identified by a total of 73,334 peptides. The sequences were used to predict the proteases active  
31 in each of the samples using Proteasix. The profile of peptide signatures is shown in Fig. 2. Fifty-six  
32 proteases (Table 1 shows all identities, protease name abbreviations and accession numbers) were  
33 found in total. Many were at a very low level as seen in Fig. 2. Additionally, it can be seen that the  
34 detection of protease generated peptides changes from day 0; there is wide variation between the  
35 different proteases; however, statistical significance can be found between day 0 and day 35 for  
36 calpain subunit 2 (CAPN2), cathepsin S (CTSS), neutrophil elastase (ELANE) and MMP7; and  
37 between day 21 and 35 for MMP7. 36 proteases were found in 80% or more of the samples; 19  
38 were found in all samples. Of these 36 predicted proteases 15 were identified in the LC-MS/MS  
39 data (Table 2).  
40  
41  
42  
43  
44  
45  
46  
47

48 Generation of correlation networks (Fig. 3) for each day showed that the network of  
49 proteases was highly connected at day 0 but that the network fractured into smaller modules (at p  
50 value threshold  $p < 0.001$ ) at day 21. By day 35 the network was more connected again but not to  
51 the same extent as at day 0.  
52  
53  
54

55 To further investigate the activity of proteases in saliva from the volunteers undergoing 21-  
56 days of experimental gingivitis development and a further 14 days of resolution, the saliva samples  
57 were used with gelatin zymography. Bands of activity were visualized consistently at >175 kDa  
58 (band A), approximately 90 kDa (band B) and approximately 75 kDa (band C). There were also  
59  
60

1  
2  
3 occasional donor samples with activity at approximately 40 kDa (band D, n=3 donors only). An  
4 example gel is shown in Fig. 4. The band area was measured and the resulting data is shown in Fig.  
5 4: bands A, B and C show some increases at day 21 and decreases at day 35, whereas band D  
6 shows greater variability. Overall, there was a definite variation associated with individual donors,  
7 when considering all the bands together. This appears to confirm the changes seen by the peptide  
8 analysis. The bands were additionally excised from the gels and in gel digested with trypsin to  
9 identify the proteases present. The results demonstrated the presence of MMP9 in bands A, B and  
10 C but not in D.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Manuscript Copy

## Discussion

In this study the activity of a network of proteases has been investigated using a proteomics (endoProteo-FASP) (24) and bioinformatics (Proteasix) (23) approach. In total 56 proteases showed some predicted activity. The proteases with the most activity (contributing to over 73% of the activity and at least 100 peptides each on average per sample) were calpain, cathepsin L, S, K and G, pepsin A3, matrix metalloproteinase (MMP) 7, and elastase. Previously, TRINDADE *et al.* (24) and Sun *et al.* (34) have looked at the wide range of proteases activity in saliva samples. Sun *et al.* (34), using histatin containing zymograms detected 13 proteases, 8 of which were also detected in this study. When comparing data presented here and that from TRINDADE *et al.*, TRINDADE *et al.* (26) predicted 66 proteases to be active in saliva using Proteasix. These matched to 45 of the proteases predicted in our dataset and there were a further 11 not in their data. This may be due to changes in the Proteasix algorithm since their publication. Indeed, the study presented here also found very similar patterns of zymogen activity on gelatin gels as compared to TRINDADE *et al.* (26) as well. The bands excised from the zymograms revealed MMP9 to be found in all but the lowest molecular weight band (approximately 45 kDa); as MMP9 is reported to be at least 78kDa this suggests that it is present either alone or in a complex, perhaps with an inhibitor or other proteins. The lower band (band D) could be associated with activity from one of the other smaller proteases predicted.

Thirty-six proteases were found in more than 80% of the samples. These proteases were examined further for their association with induction and resolution of inflammation in experimental gingivitis. No overt pattern could be seen with the data available here. This could be due to inter-individual variation in the responses to the accumulation of plaque and quantity of inflammation induced in the donors. TROMBELLI *et al.* (35) identified two groups responding to plaque induced inflammation whilst undergoing experimental gingivitis: low-responders and high-responders. The low responders displayed lower clinical measures of inflammation (gingival index and GCF volume) even though they accumulated similar levels of plaque. OFFENBACHER *et al.* (11), examining 21 day gingivitis GCF samples, adopted a hierarchical clustering technique based on changes in inflammatory mediator levels. Using this approach they saw associations between increased MMP-8 and MMP-9 levels (not activities) in group IC1, decreases in MMP-1 in group IC2 and decreases in MMP-1, MMP-3 and MMP-13 in group IC3. These were achieved by taking into account other mediators. Recently, NASCIMENTO *et al.* (36) identified slow and fast responders to plaque induced inflammation whilst undergoing experimental gingivitis. Further investigation by NASCIMENTO *et al.* (37) measured the salivary levels of MMP-8, MPO and TIMP-1 and suggested that the faster responders had greater neutrophil involvement. A larger number of participants in

1  
2  
3 future studies may help with identification of these kinds of groups using the methods presented  
4 in this study.  
5

6 The 36 proteases predicted in more than 80% of the samples were used to create correlation  
7 networks and explore the interconnectivity of their activity. Each network, created for days 0, 21  
8 and 35, demonstrated fairly sparse networks. At day 0 the network was highly connected;  
9 however, at day 21 there was a change, showing apparent removal of the network structure. At  
10 day 35 the network was once again more connected but not to the same extent as at day 0. Less  
11 connectivity in a network means that it is less flexible and there is less redundancy. In the case of  
12 these salivary protease networks this may indicate that the proteases are working more on their  
13 own and this may generate different peptides or effects. Further work on the peptides, for  
14 example predicting if they are antimicrobial (38, 39), could elucidate this further and could help  
15 drive development of therapeutics to target increases in antimicrobial peptides  
16

17 The Proteasix tool was developed to allow for automatic retrieval of protease signatures from  
18 a variety of data sources including mass spectrometry peptide sequencing. This is the only tool of  
19 its type, though there are similarities with others that can predict substrate sequences or that  
20 store information about proteolytic events (20-22). The tool relies on an underlying cleavage site  
21 database that has been curated from information on proteases from the literature and other  
22 databases (23). These cleavage sites are represented by amino acid motifs of between 2-8 amino  
23 acids with a scissile bond between amino acids P1 and P1'. In particular the MEROPS (21) database  
24 has been used to generate probabilities, specificities and sensitivities associated between  
25 predicted protease and peptide. This is important as proteases are promiscuous and have  
26 pronounced overlaps in substrate recognition. FUCHS *et al.* (40) mapped the protease specificity of  
27 61 proteases based on the similarity of their substrate sequences using sequence logos. They  
28 demonstrated a high number of less specific proteases and that these could be clustered together  
29 largely by catalytic type: for instance the MMPs clustered together and the caspases clustered  
30 together, however the cathepsins with their differing mechanisms were spread further apart. The  
31 Proteasix tool is set with high specificity (41), i.e., the percentage of sites correctly predicted not to  
32 be cleaved by a particular protease (see Table 2 for some examples). This goes a long way to  
33 ensure that the assignments are correct, though at the expense of sensitivity, the proportion of  
34 sites correctly predicted to be cleaved by a particular protease, in some cases. At present there is  
35 no automated tool for prediction of protease activity by bacterial proteases. Thus, it is difficult to  
36 take a wide approach to understanding subversion of the protease web by bacterial protease  
37 activity, however well-known proteases important in oral health such as gingipains from  
38 *Porphyromonas gingivalis*, karilysin from *Tannerella forsythia* and dentilisin from *Treponema*  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 *denticola* are listed in the MEROPS database and have sequence logos so it may be possible to  
4 examine data for specific bacterial proteases.  
5

6  
7 In previous research individual proteases or total activity have been investigated during 21  
8 day experimental gingivitis models. Elastase presence, measured by ELISA, was found to increase  
9 from day 0 to 21 in GCF (5,14) and the activity of the enzyme was also shown to increase, by a  
10 factor of three even though the quantity of its inhibitors increased even further (5). SUN *et al.* (34)  
11 have also demonstrated the co-identification of proteases from human saliva with their protein  
12 inhibitors. Interestingly, they were using zymograms too and found proteases in multiple locations  
13 as we did for MMP9. Cathepsins B, L and G have also been detected by activity assays in GCF  
14 (13,18) with an increase across the 21-day experimental gingivitis course. MMPs have been  
15 extensively investigated in both GCF and saliva. In terms of experimental gingivitis MMP-1, -3, -7, -  
16 13, -8 and -9 were examined by multiplexed antibody based assay (Luminex) revealing little  
17 change across the 21 days of induction and 28 days of resolution of inflammation (11). The MMPs  
18 are particularly interesting as a lot of research evidence has been gathered around them and their  
19 use for the detection of periodontitis (see FRANCO *et al.* (42) and DE MORAIS *et al.* (43) for reviews).  
20 They work in a protease web to activate each other from their pro-forms and can work  
21 synergistically to breakdown the extracellular matrix. MMP-7 (44), MMP-8 (45) and MMP-9 (46)  
22 have been found in multiple forms: showing complexes, latent and active forms as well as the  
23 fragments released during activation. Here we have seen high predicted protease activity for  
24 MMP-7, a salivary MMP, whereas MMP-9 showed lower predicted activity and MMP-8 very low  
25 numbers of predicted generated peptides. These latter two are generated largely in the gingiva  
26 and may have much lower activities in saliva than for example GCF adjacent to these tissues.  
27 Furthermore, MMP-7 has a very different sequence logo compared to MMP-8 and MMP-9 (Table  
28 3). Further research using dithiothreitol treated boiled saliva as a source of proteins and isolated  
29 MMPs may be able to give further evidence for the levels of protease activity and cross-over in  
30 identification. For total protease activity BIKKER *et al.* (47) reported the use of a fluorescence  
31 resonance energy transfer (FRET) assay. They could demonstrate higher total protease activity at  
32 day 21 and demonstrated that it remained elevated at day 35, 14 days after recommencement of  
33 oral hygiene.  
34  
35

36  
37 The data presented here from protease prediction suggests little change in many proteases  
38 across the experimental gingivitis period, with some showing decreases on resolution of  
39 inflammation, whereas the zymogram data suggests an increase at day 21 compared to the other  
40 days. However, there is a wide variation in both. If it is possible to isolate individual activities of  
41 individual proteases from saliva in future experiments the similarities or disparities can be  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 explored in the future, bearing in mind the complex nature of the proteases present in terms of  
4 their specificities and that of the reporting assays.  
5

6  
7 One limitation of our study is that only three teeth underwent plaque accumulation, rather  
8 than the whole mouth as in the original experimental gingivitis set up. This will diminish the  
9 quantity of inflammation in the mouth as a whole; however, we have previously demonstrated  
10 that there are quantitative changes to the GCF protein composition under these conditions (3).  
11 This level of gingivitis may be similar to that seen in an individual consistently missing small areas  
12 within their mouth rather than not cleaning at all and could equate to similar levels of gingivitis in  
13 the wider population. That a small apparent change is seen, highlights that even mild changes in  
14 gingival inflammation could change the protease network and resilience.  
15  
16  
17  
18  
19  
20  
21  
22  
23

#### 24 **Acknowledgements**

25 This work was supported by the University of Birmingham.  
26

#### 27 **Conflicts of Interest statement**

28 The authors have no conflicts of interest.  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

1. TONETTI M, JEPSEN S, JIN L, OTOMO-CORGEL J. Impact of the global burden of periodontal diseases on health, nutrition and wellbeing of mankind: A call for global action. *J Clin Periodontol* 2017; **44**: 456-462.
2. PIHLSTROM B, MICHALOWICZ B, JOHNSON N. Periodontal diseases. *Lancet* 2005; **366**: 1809-1820.
3. GRANT M, CREESE A, BARR G, LING M, SCOTT A, MATTHEWS J, GRIFFITHS H, COOPER H, CHAPPLE I. Proteomic analysis of a noninvasive human model of acute inflammation and its resolution: the twenty-one day gingivitis model. *J Proteome Res* 2010; **9**: 4732-4744.
4. BROGDEN K, JOHNSON G, VINCENT S, ABBASI T, VALI S. Oral Inflammation, a Role for Antimicrobial Peptide Modulation of Cytokine and Chemokine Responses. *Expert Rev Anti Infect Ther.* 2013; **11**: 1097-113.
5. GIANOPOULOU C, ANDERSEN E, DEMEURISSE C, CIMASONI G. Neutrophil elastase and its inhibitors in human gingival crevicular fluid during experimental gingivitis. *JDR* 1992; **71**: 359-363.
6. LOE H, HOLM-PEDERSEN P. Absence and Presence of Fluid from Normal and Inflamed Gingivae. *Periodontics* 1965; **3**: 171-177.
7. LOE H, THEILADE E, JENSEN S. Experimental Gingivitis in Man. *J Periodontol.* 1965; **36**: 177-187.
8. PUTT M, VAN DER WEIJDEN G, KLEBER C, SAXTON C. Validation of a 21-day, partial-mouth gingivitis model for evaluating chemotherapeutic dentifrices. *J Periodontal Res* 1993; **28**: 301-307.
9. SAXTON C, HUNTINGTON E, CUMMINS D. The effect of dentifrices containing Triclosan on the development of gingivitis in a 21-day experimental gingivitis study. *Int DentJ* 1993; **43**: 423-429.
10. LANG N, ANTON E, GABRIEL Y, SALVI G, PJETURSSON B, WINSTON J, HE T. An experimental gingivitis study to evaluate the clinical effects of a stannous fluoride dentifrice. *Oral health prevdenti* 2004; **2**: 369-376.



- 1  
2  
3  
4 11. OFFENBACHER S, BARROS S, MENDOZA L, MAURIELLO S, PREISSER J, MOSS K, DE JAGER M, ASPIRAS M.  
5  
6 Changes in gingival crevicular fluid inflammatory mediator levels during the induction and  
7  
8 resolution of experimental gingivitis in humans. *J Clin Periodontol* 2010; **37**: 324-333.  
9
- 10 12. LORENCINI M, SILVA J, DE LA HOZ C, CARVALHO H, STACH-MACHADO D. Changes in MMPs and  
11  
12 inflammatory cells in experimental gingivitis. *Histol histopathol* 2009; **24**: 157-166.  
13
- 14 13. KUNIMATSU K, YAMAMOTO K, ICHIMARU E, KATO Y, KATO I. Cathepsins B, H and L activities in  
15  
16 gingival crevicular fluid from chronic adult periodontitis patients and experimental  
17  
18 gingivitis subjects. *J Periodontal Res* 1990; **25**: 69-73.  
19
- 20 14. KUNIMATSU K, ICHIMARU E, KATO I, KATO Y, SONODA Y, AOKI Y, YAMAMOTO K. Granulocyte  
21  
22 medullasin levels in gingival crevicular fluid from chronic adult periodontitis patients and  
23  
24 experimental gingivitis subjects. *J Periodontal Res* 1990; **25**: 352-357.  
25
- 26 15. BOSTANCI N, RAMBERG P, WAHLANDER A, GROSSMAN J, JONSSON D, BARNES V, PAPAPANOU P. Label-  
27  
28 free quantitative proteomics reveals differentially regulated proteins in experimental  
29  
30 gingivitis. *J Proteome Res* 2013; **12**: 657-678.  
31
- 32 16. ZHOU M, MENG H, ZHAO Y, CHEN Z. Changes of four proinflammatory proteins in whole saliva  
33  
34 during experimental gingivitis. *Chin J Dent Res* 2012; **15**: 121-127.  
35
- 36 17. MARSHALL N, FINLAY B, OVERALL C. Sharpening Host Defenses during Infection: Proteases Cut  
37  
38 to the Chase. *MCP* 2017; **16**: S161-S171.  
39
- 40 18. KUNIMATSU K, MINE N, MURAOKA Y, KATO I, HASE T, AOKI Y, YAMAMOTO K. Identification and  
41  
42 possible function of cathepsin G in gingival crevicular fluid from chronic adult periodontitis  
43  
44 patients and from experimental gingivitis subjects. *J Periodontal Res* 1995; **30**: 51-57.  
45
- 46 19. PUENTE X, SANCHEZ L, OVERALL C, LOPEZ-OTIN C. Human and mouse proteases: a comparative  
47  
48 genomic approach. *Nat rev. Genetics* 2003; **4**: 544-558.  
49
- 50 20. LANGE P, HUESGEN P, OVERALL C. TopFIND 2.0-linking protein termini with proteolytic  
51  
52 processing and modifications altering protein function. *Nucleic Acids Res* 2012; **40**: D351-  
53  
54 361.  
55
- 56 21. RAWLINGS N, BARRETT A, BATEMAN A. MEROPS: the database of proteolytic enzymes, their  
57  
58 substrates and inhibitors. *Nucleic Acids Res* 2012; **40**: D343-350.  
59  
60

- 1  
2  
3  
4 22. IGARASHI Y, EROSHKIN A, GRAMATIKOVA S, GRAMATIKOFF K, ZHANG Y, SMITH J, OSTERMAN A, GODZIK  
5 A. CutDB: a proteolytic event database. *Nucleic Acids Res* 2007; **35**: D546-5493.  
6  
7  
8 23. KLEIN J, EALES J, ZURBIG P, VLAHOU A, MISCHAK H, STEVEN, R. Proteasix: a tool for automated  
9 and large-scale prediction of proteases involved in naturally occurring peptide generation.  
10  
11 *Proteomics* 2013; **13**: 1077-1082.  
12  
13  
14 24. TRINDADE F, AMADO F, GOMES P, VITORINO R. endoProteoFASP: a novel FASP approach to  
15 profile salivary peptidome and disclose salivary proteases. *Talanta* 2015; **132**: 486-493.  
16  
17  
18 25. WISNIEWSKI J, ZOUGMAN A, NAGARAJ N, MANN M. Universal sample preparation method for  
19 proteome analysis. *Nat methods* 2009; **6**: 359-362.  
20  
21  
22 26. TRINDADE F, AMADO F, OLIVEIRA-SILVA R, DANIEL-DA-SILVA A, FERREIRA R, KLEIN J, FARIA-ALMEIDA R,  
23 GOMES P, VITORINO R. Toward the definition of a peptidome signature and protease profile  
24 in chronic periodontitis. *Proteom Clin appl* 2015; **9**: 917-927.  
25  
26  
27  
28 27. CHAPPLE I, SOCRANSKY S, DIBART S, GLENWRIGHT H, MATTHEWS J. Chemiluminescent assay of  
29 alkaline phosphatase in human gingival crevicular fluid: investigations with an  
30 experimental gingivitis model and studies on the source of the enzyme within crevicular  
31 fluid. *J Clin Perio* 1996; **23**: 587-594.  
32  
33  
34  
35  
36 28. LOBENE R, SOPARKAR P, NEWMAN M. Use of dental floss. Effect on plaque and gingivitis. *Clin*  
37 *prev dent* 1982; **4**: 5-8.  
38  
39  
40 29. LOE H. The Gingival Index, the Plaque Index and the Retention Index Systems. *J Periodontol*  
41 1967; **38**: Suppl:610-616.  
42  
43  
44 30. SMITH P, KROHN R, HERMANSON G, MALLIA A, GARTNER F, PROVENZANO M, FUJIMOTO E, GOEKE N,  
45 OLSON B, KLENK D. Measurement of protein using bicinchoninic acid. *Anal Biochem* 1985;  
46 **150**: 76-85.  
47  
48  
49 31. SCHNEIDER C, RASBAND,W, ELICEIRI K. NIH Image to ImageJ: 25 years of image analysis. *Nat*  
50 *methods* 2012; **9**: 671-675.  
51  
52  
53 32. SHEVCHENKO A, TOMAS H, HAVLIS J, OLSEN J, MANN M. In-gel digestion for mass spectrometric  
54 characterization of proteins and proteomes. *Nat Protoc* **2006**; **1**: 2856–2860  
55  
56  
57 33. BASTIAN M, HEYMAN S, JACOMY M. Gephi: an open source software for exploring and  
58 manipulating networks. *AAAI ICWSM* 2009.  
59  
60

- 1  
2  
3  
4 34. SUN X, SALIH E, OPPENHEIM F, HELMERHORS E. Activity-based mass spectrometric  
5 characterization of proteases and inhibitors in human saliva. *Proteomics. Clin appl* 2009; **3**:  
6 810-820.  
7  
8  
9  
10 35. TROMBELLI L, TATAKIS D, SCAPOLI C, BOTTEGA S, ORLANDINI E, TOSI M. Modulation of clinical  
11 expression of plaque-induced gingivitis. II. Identification of "high-responder" and "low-  
12 responder" subjects. *J Clin Periodontol* 2004; **31**: 239-252.  
13  
14  
15 36. NASCIMENTO G, DANIELSEN B, BAEUM V, LOPEZ R. Identification of inflammatory response  
16 patterns in experimental gingivitis studies. *Eur J Oral Sci* 2019; **127**: 33-39.  
17  
18  
19 37. NASCIMENTO G, BAEUM V, SORSA T, TERVAHARTIALA T, SKOTTRUP P, LOPEZ R. Salivary levels of  
20 MPO, MMP-8 and TIMP-1 are associated with gingival inflammation response patterns  
21 during experimental gingivitis. *Cytokine* 2019; **115**: 135-141.  
22  
23  
24 38. BHADRA P, YAN J, LI J, FONG S, SIU S. AmPEP: Sequence-based prediction of antimicrobial  
25 peptides using distribution patterns of amino acid properties and random forest. *Sci Rep*  
26 2018; **8**: 1697.  
27  
28  
29 39. MEHER P, SAHU T, SAINI V, RAO A. Predicting antimicrobial peptides with improved accuracy  
30 by incorporating the compositional, physico-chemical and structural features into Chou's  
31 general PseAAC. *Sci Rep* 2017; **7**: 42362.  
32  
33  
34 40. FUCHS J, VON GRAFENSTEIN S, HUBE R, KRAMER C, LIEDL K. Substrate-driven mapping of the  
35 degradome by comparison of sequence logos. *PLoS Comput Biol* 2013; **9**: e1003353.  
36  
37  
38 41. RAWLINGS N. Peptidase specificity from the substrate cleavage collection in the MEROPS  
39 database and a tool to measure cleavage site conservation. *Biochimie* 2016; **122**: 5-30.  
40  
41  
42 42. FRANCO C, HERNÁNDEZ-RÍOS P, SORSA T, BIGUETTI C, HERNÁNDEZ M. Matrix Metalloproteinases as  
43 Regulators of Periodontal Inflammation. *Int J Mol Sci* 2017; **18**: 440.  
44  
45  
46 43. DE MORAIS E, PINHEIRO J, LEITE R, SANTOS P, BARBOZA C, FREITAS R. Matrix metalloproteinase-8  
47 levels in periodontal disease patients: A systematic review. *J Periodont Res* 2018; **53**: 156-  
48 163.  
49  
50  
51 44. LUNDMARK A, JOHANNSEN G, ERIKSSON K, KATS A, JANSSON L, TERVAHARTIALA T, RATHNAYAKE N,  
52 AKERMAN S, KLINGE B, SORSA T, YUCEL-LINDBERG T. Mucin 4 and matrix metalloproteinase 7 as  
53 novel salivary biomarkers for periodontitis. *J Clin Periodontol* 2017; **44**: 247-254.  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4 45. GURSOY U, KONONEN E, TERVAHARTIALA T, GURSOY M, PITKANEN J, TORVI P, SUOMINEN A, PUSSINEN  
5 P, SORSA T. Molecular forms and fragments of salivary MMP-8 in relation to periodontitis. *J*  
6 *Clin Periodontol* 2018; **45**: 1421-1428.  
7  
8  
9  
10 46. BILDT M, BLOEMEN M, KUIJPERS-JAGTMAN A, VON DEN HOFF J. Collagenolytic fragments and  
11 active gelatinase complexes in periodontitis. *J Periodontol* 2008; **79**: 1704-1711.  
12  
13  
14 47. BIKKER F, NASCIMENTO G, NAZMI K, SILBEREISEN A, BELIBASAKIS G, KAMAN W, LOPEZ R, BOSTANCI N.  
15 Salivary Total Protease Activity Based on a Broad-Spectrum Fluorescence Resonance  
16 Energy Transfer Approach to Monitor Induction and Resolution of Gingival Inflammation.  
17  
18  
19  
20 *Mol Diagn Ther* 2019; **23**: 667-676.  
21  
22  
23  
24  
25

## 26 Supporting Information

27  
28 Additional supporting information may be found in the online version of the article:  
29

30  
31 **Fig. S1.** Saliva processing for endoProteo-FASP protocol  
32  
33  
34  
35  
36

## 37 Figure legends

38  
39 Figure 1. Change in plaque index and gingival index over the course of the 21 day experimental  
40 gingivitis induction and resolution phases. Shielded side denotes the teeth covered with the  
41 removable splint to prevent brushing and contralateral side denotes teeth cleaned as normal and  
42 monitored for comparison.  
43  
44  
45  
46  
47

48 Figure 2. Abundance of peptides detected with signatures for all proteases as predicted by  
49 Proteasix. Data represents all days and donors and is shown as mean+/- standard deviation. Data  
50 were analysed by oneway ANOVA. \* p<0.05, \*\*\*p<0.001, \*\*\*\*p<0.0001  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Figure 3. Correlation network graphs of each time point. A. day 0; B. day 21 and C. day 35. Size of  
5 node is proportional to the number of peptides predicted to be generated by that protease. Edges  
6 represent correlations with  $p < 0.001$ .  
7  
8  
9

10  
11 Figure 4. Quantitation of bands found on zymograms. Representative example of zymogram, saliva  
12 samples (10ug) separated under non-reducing conditions on a 1% gelatin polyacrylamide gel. After  
13 renaturing the gel was developed for 18h and then stained with Coomassie Blue. m represents  
14 molecular weight markers, 7,8 and 9 represent the saliva donors and D0, D21 and D35 represent  
15 days 0, 21 and 35. A, B, C and D represent the bands of interest. Quantitation of all bands shown in  
16 labelled graphs. Box and whiskers plots: boxes represent 75th percentile with upper line, median  
17 with middle line and 25th percentile with lower line of box; whiskers represent upper and lower  
18 limits of all the data. Band D shown as individual data for the 3 individuals displaying this behavior.  
19 All bands shows summed data from each band for each individual.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

33 Table legends

34  
35 Table 1. Accession numbers and descriptions for all proteases predicted by Proteasix  
36  
37

38 Table 2. Proteases identified by peptides in mass spectrometry analysis  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1. Accession numbers and descriptions for all proteases predicted by Proteasix

Protein name	Accession number	Protein description
ADAMTS4	O75173	A disintegrin and metalloproteinase with thrombospondin motifs 4
BMP1	P13497	Bone morphogenetic protein 1
CAPN1	P07384	Calpain-1 catalytic subunit
CAPN2	P17655	Calpain-2 catalytic subunit
CASP1	P29466	Caspase-1
CASP 2	P42575	Caspase-2
CASP 6	P55212	Caspase-6
CASP7	P55210	Caspase-7
CASP 8	Q14790	Caspase-8
CMA1	P23946	Chymase
CTRC	Q99895	Chymotrypsin-C
CTSB	P07858	Cathepsin B
CTSD	P07339	Cathepsin D
CTSE	P14091	Cathepsin E
CTSG	P08311	Cathepsin G
CTSK	P43235	Cathepsin K
CTSL	P07711	Cathepsin L1
CTSS	P25774	Cathepsin S
ELANE	P08246	Neutrophil elastase
F2	P00734	Prothrombin
GZMA	P12544	Granzyme A
GZMB	P10144	Granzyme B
GZMK	P49863	Granzyme K
GZMM	P51124	Granzyme M
HPN	P05981	Serine protease hepsin
HTRA2	O43464	Serine protease HTRA2, mitochondrial
KLK3	P07288	Prostate-specific antigen
KLK4	Q9Y5K2	Kallikrein-4
KLK5	Q9Y337	Kallikrein-5
KLK6	Q92876	Kallikrein-6

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

LGMMN	Q99538	Legumain
MEP1A	Q16819	Meprin A subunit alpha
MME	P08473	Neprilysin
MMP1	P03956	Interstitial collagenase
MMP12	P39900	Macrophage metalloelastase
MMP13	P45452	Collagenase 3
MMP14	P50281	Matrix metalloproteinase-14
MMP2	P08253	72 kDa type IV collagenase
MMP25	Q9NPA2	Matrix metalloproteinase-25
MMP3	P08254	Stromelysin-1
MMP7	P09237	Matrilysin
MMP8	P22894	Neutrophil collagenase
MMP9	P14780	Matrix metalloproteinase-9
PCSK2	P16519	Neuroendocrine convertase 2
PCSK5	Q92824	Proprotein convertase subtilisin/kexin type 5
PGA3	P0DJD8	Pepsin A-3
PGC	P20142	Gastricsin
PITRM1	Q5JRX3	Presequence protease, mitochondrial
PLG	P00747	Plasminogen
PRTN3	P24158	Myeloblastin
ST14	Q9Y5Y6	Suppressor of tumorigenicity 14 protein
THOP1	P52888	Thimet oligopeptidase
TMPRSS11E	Q9UL52	Transmembrane protease serine 11E
TMPRSS6	Q8IU80	Transmembrane protease serine 6
TMPRSS7	Q7RTY8	Transmembrane protease serine 7
TRY3	P35030	Trypsin-3

---

Table 2. Proteases identified by peptides in mass spectrometry analysis.

Protease symbol	Protease name	<u>Protease UniProt Accession</u>	Size (kDa)
ADAMTS4	A disintegrin and metalloproteinase with thrombospondin motifs 4	<u>Q75173</u>	90
CTSD	Cathepsin D	<u>P07339</u>	45
CTSE	Cathepsin E	<u>P14091</u>	43
CTSG	Cathepsin G	<u>P08311</u>	29
ELANE	Neutrophil elastase	<u>P08246</u>	29
GZMM	Granzyme M	<u>P51124</u>	28
HTRA2	Serine protease HTRA2	<u>O43464</u>	49
KLK4	Kallikrein-4	<u>Q9Y5K2</u>	27
KLK6	Kallikrein-6	<u>Q92876</u>	27
MMP14	Matrix metalloproteinase-14	<u>P50281</u>	66
MMP7	Matrilysin	<u>P09237</u>	30
MMP9	Matrix metalloproteinase-9	<u>P14780</u>	78
PGA3	Pepsin A-3	<u>P0DJD8</u>	42
PLG	Plasminogen	<u>P00747</u>	91
TMPRSS7	Transmembrane protease serine 7	<u>Q7RTY8</u>	94



**Table 3.** Sequence logos for MMPs. Data extracted from MEROPS [21] (<https://www.ebi.ac.uk/merops/>). The letters represent the amino acid in that position, with lower case letters indicating lower specificity and capital letters demonstrating a higher specificity for that amino acid's presence. Hyphens represent any amino acid and the + represents the scissile bond (cleavage site). Amino acids P1 and P1' are to the left and right of the scissile bond respectively.

Matrix metalloproteinase	Sequence logo	Sensitivity of detection by Proteasix	Specificity of detection by Proteasix
MMP1	-/pa/-/+li/-/-/-	31	98.45
MMP2	-/p/-/+li/-/-/-	76	86.88
MMP3	-/p/ag/-+l/-/-/-	70	79.35
MMP7	-/pa/-/+L/-/-/-	54	53.97
MMP8	g/pas/-/g+l/-/g/-	38	90.6
MMP9	g/pa/-/g+l/-/ga/-	63	63.68
MMP12	g/pag/ag/ga+l/-/ga/-	67	88.47
MMP13	g/P/-/g+l/-/ga/-	70	93.66
MMP14	-/p/-/+l/-/-/-	75	80.66
MMP25	-/a/-/+l/-/-/-	28	82.71





Figure 1 Change in plaque index and gingival index over the course of the 21 day experimental gingivitis induction and resolution phases. Shielded side denotes the teeth covered with the removable splint to prevent brushing and contralateral side denotes teeth cleaned as normal and monitored for comparison.

80x91mm (300 x 300 DPI)




Figure 2. Abundance of peptides detected with signatures for all proteases as predicted by Proteasix. Data represents all days and donors and is shown as mean+/- standard deviation. Data were analysed by oneway ANOVA. \* p<0.05, \*\*\*p<0.001, \*\*\*\*p<0.0001

170x70mm (300 x 300 DPI)




Figure 3. Correlation network graphs of each time point. A. day 0; B. day 21 and C. day 35. Size of node is proportional to the number of peptides predicted to be generated by that protease. Edges represent correlations with  $p < 0.001$ .

80x209mm (300 x 300 DPI)




Figure 4. Quantitation of bands found on zymograms. Representative example of zymogram, saliva samples (10ug) separated under non-reducing conditions on a 1% gelatin polyacrylamide gel. After renaturing the gel was developed for 18h and then stained with Coomassie Blue. m represents molecular weight markers, 7,8 and 9 represent the saliva donors and D0, D21 and D35 represent days 0, 21 and 35. A, B, C and D represent the bands of interest. Quantitation of all bands shown in labelled graphs. Box and whiskers plots: boxes represent 75th percentile with upper line, median with middle line and 25th percentile with lower line of box; whiskers represent upper and lower limits of all the data. Band D shown as individual data for the 3 individuals displaying this behavior. All bands shows summed data from each band for each individual.

170x217mm (300 x 300 DPI)

## Supporting Information

### **Predicted salivary human protease activity in experimental gingivitis revealed by endoProteo-FASP approach**

MULKERN D, HEWITT A, PARKER H, BATT J, YONEL Z, GRANT MM

School of Dentistry, University of Birmingham, Birmingham, United Kingdom

School of Life Sciences, University of Warwick, Coventry, United Kingdom

**Fig. S1.** Saliva processing for endoProteo-FASP protocol